



.1

I I
1995

. M.

. M.

- *1. *Autalia longicornis* Scherpeltz, 1947 [*impressa* Oliv.]. 3
2. *Corolalia obscura* (Gravenhorst, 1802) = *Falagria obscura* Grav. 6
3. *Falagria sulcata* (Paykull, 1789). 1
4. *Tachyusa coarctata* (Erichson, 1837). 6
5. *Bolitochara lunulata* (Paykull, 1789). 8
6. *Bolitochara obliqua* Erichson, 1837. 2
7. *Oxypoda lividipennis* Mannerheim, 1830 = *luteipennis* Erichs. 5
8. *Oxypoda alternans* (Gravenhorst, 1802). 6
9. *Oxypoda opaca* (Gravenhorst, 1802). 1
10. *Astilbus canaliculata* (Fabricius, 1787) = *Myrmedonia canaliculata* F. 5
11. *Zyras lugens* (Gravenhorst, 1802) = *Myrmedonia lugens* Grav. 6
12. *Zyras laticollis* (Markel, 1844) = *Myrmedonia laticollis* Maerkel. 1
13. *Zyras funestus* (Gravenhorst, 1806) = *Myrmedonia funesta* Grav. 5
14. *Zyras humeralis* (Gravenhorst, 1802) = *Myrmedonia humeralis* Grav. 4
15. *Zyras cognatus* (Markel, 1842) = *Myrmedonia cognata* Maerkel. 7
16. *Atemeles emarginatus* (Paykull, 1789). 1
Atemeles paradoxus (Gravenhorst, 1806).
17. *Thamiaraea cinnamomea* (Gravenhorst, 1802) = *Homalota cinnamomea* Grav. 7
18. *Dinaraea aequata* (Erichson, 1837) = *Homalota aequata* Erich. 13
- *19. *Placusa complanata* Erichson, 1839 [*pumilio* Grav.]. 1
20. *Gymnusa brevicollis* (Paykull, 1800). 1
- *21. *Thyasophila inquilina* (Markel, 1842) [*Homoeusa acuminata* Maerkel.]. 2
22. *Aleochara curtul a* (Goeze, 1777) = *fuscipes* F. 2

23. *Aleochara brevipennis* Gravenhorst,1806. 1
1 1 .. [A. fumata Gr.]
24. *Aleochara bipustulata* (Linnaeus,1761) =*nitida* Grav. 2
25. *Gyrophaena nana* (Paykull,1800). 5
26. *Tachyporus chrysomelinus* (Linnaeus,1758). 1
Tachyporus hypnorum (Fabricius,1775).
Tachyporus obtusus (Linnaeus,1767).
Tachyporus macropterus Stephens,1833 =*scitulus* Erich.
27. *Lordithon striatus* (Olivier,1795) =*Boletobius striatus* Oliv. 2
Lordithon lunulatus (Linnaeus,1761) =*Boletobius atricapillus* F.
Lordithon thoracicus (Fabricius,1777) =*Boletobius pygmaeus* F.
- *28. *Conosoma marshami* (Stephens,1832) [*Conurus pubescens* Grav.
=*Oxyporus testaceus* F.]. 1
29. *Conosoma littoreum* (Linnaeus,1758) =*Conurus littoreus* L.
1 .. *Conurus bipustulatus* Grav.
30. *Tachinus subterraneus* (Linnaeus,1758). 1 , 2
31. *Tachinus marginatus* (Fabricius,1792) =*marginatus* Gyll. 1
1 . , *T. rufipes*
- De Geer.
32. *Tachinus rufipes* (Degeer,1774). 1
33. *Tachinus lignorum* (Linnaeus,1758) =*flavipes* F. 1
T. humeralis Grav.
34. *Tachinus fimetarius* Gravenhorst,1802. 1
35. *Tachinus bipustulatus* (Fabricius,1792). 1
36. *Tachinus pallipes* (Gravenhorst,1806). 1
37. *Tachinus laticollis* Gravenhorst,1802. 2 . c 1
T. bipustulatus F.
Tachinus marginellus (Fabricius,1781) =*laticollis* v. *immaturus* Grav.
38. *Tachinus elongatus* Gyllenhal,1810. 1
39. *Cryptobium fracticorne* (Paykull,1800). 2
40. *Lathrobium elongatum* (Linnaeus,1767). 3
41. *Lathrobium quadratum* (Paykull,1789). 1
42. *Paederus riparius* (Linnaeus,1758). 5
43. *Paederus fuscipes* Curtis,1826 =*longipennis* Erich. 1
44. *Stilicus rufipes* Germar,1836. 2
45. *Stilicus similis* (Erichson,1839). 1
- *46. *Astenus pulchellus* (Heer,1839) [*Sunius angustatus* F.]. 1

79. *Nudobius lentus* (Gravenhorst,1806) =*Xantholinus lentus* Grav. 2
80. *Xantholinus tricolor* (Fabricius,1787). 1
Xantholinus linearis (Olivier,1795).
81. *Philonthus cyanipennis* (Fabricius,1792). 1
82. *Philonthus nitidus* (Fabricius,1787). 1
83. *Philonthus politus* (Linnaeus,1758) =*aeneus* Rossi. 3
84. *Philonthus albipes* (Gravenhorst,1802). 3
85. *Philonthus atratus* (Gravenhorst,1802). 4
Philonthus aerosus Kiesenwetter,1851.
86. *Philonthus marginatus* (Strom,1768) =*marginatus* [F.]. 2
87. *Philonthus rigidicornis* (Gravenhorst,1802) =*fimetarius* Grav. 1
, 1 , 3 ,
[*Philonthus sordidus* Grav.].
88. *Philonthus ebeninus* (Gravenhorst,1802). 2
89. *Philonthus sanguinolentus* (Gravenhorst,1802). 5
90. *Philonthus cruentatus* (Gmelin,1790) =*bipustulatus* Pz. 5
91. *Philonthus micans* (Gravenhorst,1802). 1
92. *Philonthus fulvipes* (Fabricius,1792). 3
93. *Philonthus tenuis* (Fabricius,1792). 4
94. *Neobisnius villosulus* (Stephens,1833). =*Philonthus elongatulus*
Erichs. 1
95. *Gabrius splendidulus* (Gravenhorst,1802) =*Philonthus splendidulus*
Grav. 2 , 2
96. *Velleus dilatatus* (Paykull,1787) =*Quedius dilatatus* [F.]. 2
- *97. *Quedius mesomelinus* (Marsham,1802) [*fulgidus* G.]. 1
98. *Quedius molochinus* (Gravenhorst,1806). 1
99. *Quedius fuliginosus* (Gravenhorst,1802). 3
100. *Quedius brevis* Erichson,1840. 5
101. *Quedius xanthopus* Erichson,1839. 1
102. *Oxyporus maxillosus* Fabricius,1792 =*angularis* Gebler. 8
103. *Oxyporus rufus* (Linnaeus,1758). 8
104. *Oxyporus mannerheimi* Gyllenhal,1827. 3
- *105. *Anthophagus angusticollis* (Mannerheim,1830) [*caraboides* L.]. 4
106. *Anthophagus caraboides* (Linnaeus,1758) =*testaceus* Grav. 5
107. *Arpedum quadrum* (Gravenhorst,1806). 1
108. *Deliphrum tectum* (Paykull,1789). 6
109. *Omalium rivulare* (Paykull,1792). 4
110. *Omalium caesum* Gravenhorst,1806. 2
111. *Phloeonomus planus* (Paykull,1792) =*Omalium planum* Grav. 1

3. *Stenus* Latr. (Coleoptera, Staphylinidae) // . - . -1991. - . 168-173.
4. // . - . -1991. - . 4-37.
5. -1905. -1024 .
6. Enumeratio Coleopterorum Fennoscandiae et Daniae / H. Silfverberg. -Helsinki:Tamprint. -1979. -79 s.

(Coleoptera . *Staphylinidae*) // . - . — . : i i , 1995. . 1. . 131—138.

XIX . () (-), 117 , 20 : 6 .

Pisanenko A.D., Alexandrovich O.R. Revision of *Staphylinidae* (Coleoptera) from N.M. Arnold's collection // Fauna and taxonomy: Proceed. Zool. museum Byel. Univ. — Minsk: Nauka i tekhnika 1995. Vol. I. P. 131—138.

Study of N. M. Arnold's collection which was collected in Mogilev region (E Belarus) in the second half of XIX Century and is preserved in Zoological Institute' AN of Russia (St. Petersburg) revealed 117 species of *Staphylinidae*. Twenty species from this collection were wrongly identified previously. List of the valid names, synonyms from Arnold's publication and number of specimens is reported.