

Report of ciliate epibionts (Ciliophora, Suctorea) on meiobenthic invertebrates from the Indian coast near Karwar, Karnataka

Tapas Chatterjee¹, Igor Dovgal² and Mandar Nanajkar³

¹ Crescent International School, Bario, Govindpur, Dhanbad 828109, Jharkhand, India

² A. O. Kovalevsky Institute of Biology of the Southern Seas RAS, Sevastopol, 299011, Russia

³ CSIR, National Institute of Oceanography, Dona Paula, Goa-403004, India

| Submitted March 16, 2020 | Accepted April 17, 2020 |

Summary

The article describes four epibiont suctorian ciliates, viz. *Thecacinetta cothurnioides* (Collin, 1909) and *Trematosoma rotunda* (Allgén, 1952) on the nematode *Tricomma* sp., *Thecacinetta calix* (Schröder, 1907) and *Acinetta tuberosa* Ehrenberg, 1834 on the harpacticoid copepod, from the Karwar coast of India (Karnataka state), the Arabian Sea. *Trematosoma rotunda* is reported here first time as epibiont on the genus *Tricomma* representative. The systematic positions of the found suctorian species, their brief descriptions, measurements, and data on their distribution, hosts and localisations at the host body are presented.

Key words: ciliate epibiont, marine meiobenthos, India, Suctorea

Introduction

The phenomenon of epibiosis is common in marine biocenoses, and suctorian ciliates belong to epibiont ciliates that most frequently occur in aquatic habitats. Suctorians can be found on a wide diversity of hosts and substrates. The majority of these ciliates are commensals of various water invertebrates or vertebrates (Dovgal, 2002, 2013).

Epibiont ciliates from the west coast of India were studied by many authors (Santhakumari 1985, 1986; Chatterjee, 1996; Dovgal et al., 2008, 2009; Chatterjee et al., 2013, 2019a, 2019b).

The present article reports four suctorian ciliate species inhabiting meiobenthic invertebrate hosts (harpacticoid copepods and nematodes) from the Karwar coast of the Arabian Sea, Karnataka state, India.

Material and methods

The study area is located 5 kilometres south of Karwar (District: Uttar Kannada, Karnataka), Latitude: 14°46'38.57" N, Longitude: 74°06'59.92"E, which is near the Goa-Karnataka border, west coast

Fig. 2. Ciliate species found near Karwar, Indian coast. A – *Thecacineta cothurnioides* (Collin, 1909) on *Tricoma* sp.; B – *Thecacineta calix* (Schröder, 1907) on harpacticoid copepod; C – *Acineta tuberosa* Ehrenberg, 1834 on furca of harpacticoid copepod; D – *Trematosoma rotunda* (Allgén, 1952) on *Tricoma* sp.

deep sea. Details of distribution and host specificity of this species are given in Chatterjee et al. (2019).

Subclass Endogenia Collin, 1012
Order Acinetida Raabe, 1964
Family Acinetidae Ehrenberg, 1834
Genus *Acineta* Ehrenberg, 1834

Acineta tuberosa Ehrenberg, 1834 (Fig. 2 C)

Material examined. Two individuals found on a harpacticoid copepod, attached near the furca region of the host.

Brief description. Suctorian ciliate with a smooth, triangular or cup-shaped, laterally flattened lorica. The lorica aperture is dumbbell-shaped. The upper half of the lorica is characterized by sharp widening. The body is attached to the bottom of the lorica. Macronucleus spherical or elongated, medial, located along the longitudinal axis of the cell. The single contractile vacuole located subapically, above the macronucleus. The stalk is long, sometimes

curved, equipped with a well-developed attachment disk. In the area of connection with the lorica, the stalk is somewhat widened, in some cases might be dipped to the base of the shell.

Measurements based on two individuals (in μm): lorica length 57-96, lorica width 22-36, lorica mouth width 38-45, stalk length 18-24, stalk diameter 4-5, macronucleus diameter 14.

Remarks. The species was reported from marine and brackish waters as periphytic or nonspecific commensals of various aquatic invertebrates (Dovgal, 2013).

In India, this species was reported from Andhra Pradesh coast (Bay of Bengal) as epiphytic on algae (Radhakrishna, 1984; Kalavati and Raman, 2008). Chatterjee et al. (2013) reported this species on a harpacticoid copepod from the Dias beach, Dona Paula, Goa, India (the Arabian sea).

Genus *Trematosoma* Batisse, 1972

Trematosoma rotunda (Allgén, 1952) (Fig. 2 D)

Material examined. Numerous individuals distributed throughout the body surface of the nematode *Tricoma* sp.

Brief description. Marine loricate suctorian. Cell body short, rounded, laterally flattened, entirely fills up the lorica and clamped to their aperture border. Lorica delicate, gently striated. Tentacles clavate, short, arranged at the upper body surface as a row in two groups. Macronucleus spherical or oval, centrally located. Stalk short, thin, slightly ribbed.

Measurements based on four individuals (in μm): lorica length 26–29, lorica width 14–22, body length 19–39, body width 11–20, stalk length 5–15, stalk diameter 3–4, macronucleus diameter 6–8, length of tentacles 3–12.

Remarks. Allgén (1952) reported this species from Falkland Islands, located near the southern tip of Argentina and Antarctic Peninsula (Graham Land) on the nematodes *Desmodora tenuispiculum* Allgén, 1928 and *D. stateni* Allgén, 1928, respectively. Dovgal et al. (2009) provided remarks on systematics and nomenclature of this species. This species was reported from Ratnagiri, Maharashtra, west coast of India, the Arabian Sea, as epibiont on the nematodes *Pseudochromadora* sp. (Dovgal et al., 2009). The above-mentioned authors also identified this species from the intertidal zone of North-Eastern Queensland, Australia, based on microphotograph of Fisher (2003). Bhattacharjee (2014) reported it at the coast of Rushikulya (Odisha State of India) on the nematode *Chromaspirina* sp. Ansari and Bhadury (2016) reported it from the mangrove ecosystem of Sagar Island of Sundarbans from the north-western Bay of Bengal, on the nematodes *Dorylaimopsis punctata*. Ghosh and Mandal (2019) reported it on the nematodes *Desmodora scaldensis* de Man, 1889 from the subtidal region of the Matla estuary, Sundarbans, Bay of Bengal. In the present record, this species was registered near Karwar, Karnataka state of the west coast of India, on the nematode *Tricoma* sp. *Trematosoma rotunda* is reported here first time as an epibiont on the host from the genus *Tricoma*.

Acknowledgements

Second author's (ID) work was fulfilled within the framework of research topic #AAAA-A19-119060690014-5 at A.O. Kovalevsky Institute of Biology of the Southern Seas RAS.

References

- Allgén C.A. 1952. Über zwei neue antarktische Suctorien aus dem Gracham Land. K. Fysiograf. Sälls. i Lund Forhändl. 21 (18), 173–176.
- Ansari K.G.M.T. and Bhadury P. 2016. Occurrence of epibionts associated with meiofaunal basibionts from the world's largest mangrove ecosystem, the Sundarbans. Mar. Biodiver. 47, 539–548.
- Bhattacharjee D. 2014. Suctorian epibionts on *Chromaspirina* sp. (Nematoda: Desmodoridae) from the shallow continental shelf of the Bay of Bengal, northern Indian Ocean. Mar. Biodivers. Rec. 7 (e49), 1–3.
- Chatterjee T. 1996. Ciliate infestation on the phytal Halacaridae (Acari) from the Kovalam Beach (Kerala coast). J. Bombay Nat. Hist. Soc. 93, 107–108.
- Chatterjee T., Fernandez-Leborans G., Ramteke D. and Ingole B.S. 2013. New records of epibiont Ciliates (Ciliophora) from Indian coast with descriptions of six new species. Cah. Biol. Mar. 54, 143–159.
- Chatterjee T., Nanajkar M. and Dovgal I. 2019a. New record of *Loricophrya stresemanni* (Ciliophora: Suctorea) as epibiont on nematodes from the Indian Ocean and notes on the genus *Loricophrya*. Cah. Biol. Mar. 60, 283–288.
- Chatterjee T., Nanajkar M., Dovgal I., Sergeeva N. and Bhav S. 2019b. New records of epibiont *Thecacineteta calix* (Ciliophora: Suctoria) from the Caspian Sea and Angria Bank, Arabian Sea. Cah. Biol. Mar. 60, 445–451.
- Collin B. 1909. Diagnoses preliminaries d'Acinetiens nouveaux ou mal connus. C.R. Hebd. Seances Acad. Sci. 149, 1094–1095.
- Dovgal I.V. 2002. Evolution, phylogeny and classification of Suctorea (Ciliophora). Protistology. 2, 194–270.
- Dovgal I.V. 2013. Suctorians (Ciliophora, Suctorea). In: Fauna of Ukraine, vol. 36. Ciliates. Issue 1. Naukova Dumka, Kiev (In Russian with English summary).
- Dovgal I., Chatterjee T. and Ingole B. 2008. An overview of suctorian ciliates (Ciliophora, Suctorea) as epibionts of halacarid mites (Acari: Halacaridae). Zootaxa. 1810, 60–68.
- Dovgal I., Chatterjee T. and Ingole B. 2009. New records of *Thecacineteta cothurnioides* and *Trematosoma rotunda* (Ciliophora, Suctorea) as epibionts on nematodes from Indian Ocean. Protistology. 6, 19–23.

Fisher R. 2003. Ciliate hitch-hikers-nematode ecto-commensals from tropical Australian sea-grass meadows. *J. Mar. Biol. Assoc. U.K.* 83, 445–446.

Ghosh M. and Mandal S. 2019. Living with nematode: an epibiont *Trematosoma rotunda* associated with basibiont *Desmodora scaldensis* from Matla Estuary, Sundarbans, India. *Thalassas.* 35, 619–624.

Kalavati C. and Raman A.V. 2008. Taxonomy and ecology of protozoan from marginal marine environments of east coast of India. *Rec. Zool. Surv. India, Occ. Paper* 282, 1–136.

Radhakrishna G. 1984. Studies on some aspects of ecology of ciliates associated with algae of Visakhapatnam coast. Ph.D. Thesis. Andhra University.

Santhakumari V. 1985. Epizoic and ectoparasitic protozoans from planktonic copepods of the southwest and southeast coasts of India with the description of a new species. *J. Mar. Biol. Ass. India.* 27, 29–38.

Santhakumari V. 1986. Two new species of the suctorians, *Acineta satyanandani* sp. nov. and *Paracineta karunakarani* sp. nov. epizoic on ostracodes. *Indian J. Fish.* 33, 471–475.

Address for correspondence: Igor Dovgal. A.O. Kovalevsky Institute of Biology of the Southern Seas RAS, Nakhimov Ave. 2, Sevastopol, 299011, Russia; e-mail: dovgal-1954@mail.ru