

List of Chinese Insects

Vol. II

Hua Li – zhong

Zhongshan (Sun Yat – sen) University Press
Guangzhou, 2002

- Mycetina emmerichi* Mader, [CHINA: Sichuan]
Mycetina fulva Chujo, 1938 [CHINA: Taiwan]
Mycetina humerosignata Nakane, [CHINA: Taiwan]
Mycetina maderi Strohecker, 1982 [CHINA]
Mycetina sasaji Strohecker, 1982 [CHINA: Taiwan]
Parindalmus sinensis Strohecker, [CHINA]
Pedanus gerstackeri Gorham, 1873 [CHINA: Taiwan; INDONESIA]
Phaeonychus rufipennis (Motschulsky, 1860) (*Endomychus*) [CHINA: Taiwan; JAPAN]
Phaeonychus similis Chujo, 1938 [CHINA: Taiwan]
Saula chujoi Sasaji, [CHINA: Taiwan]
Saula japonica Gorham, 1874 [CHINA: Taiwan; JAPAN]
Saula nigripes Gerstacker, 1858 [CHINA: Taiwan; INDIA, SRI LANKA, MYANMAR, CALEMANDAN]
Saula taiwana Chujo, 1938 [CHINA: Taiwan] Host: *Cannamomum camphora*.
Stenotarsoides maculosus Fairmaire, 1888 [CHINA: Beijing]
Stenotarsus aokii Chujo, 1942 [CHINA: Taiwan]
Stenotarsus chujoi Strohecker, [CHINA]
Stenotarsus porcellus Strohecker, [CHINA: Taiwan]
Stenotarsus ryukyensis Chujo et Kiuchi, 1974 [CHINA: Taiwan; JAPAN]
Stenotarsus yoshionis Chujo, 1938 [CHINA: Taiwan]
Trochoides desjardinsi Guerin - Meneville, 1838 [CHINA: Taiwan; INDIA, MYANMAR, THAILAND, MALAYSIA, SRI LANKA, INDONESIA, PHILIPPINES, CALEMANDAN, MADAGASCAR, MAURITIUS, E. AFRICA]

Family Merophysidae

- Holoparamesus capitatus* Wollaston, 1874 [CHINA: Henan, Hubei]
Holoparamesus caularum (Aube,) [CHINA: Guangxi] Host: rice.
Holoparamesus depressus Curtis, 1833 Syn. *Calyptridium kunzei* Aube, 1843 [CHINA: Liaoning, Inner Mongolia, Hebei, Shandong, Henan, Shanxi, Shaanxi, Gansu, Qinghai, Hubei, Anhui, Jiangsu, Jiangxi, Zhejiang, Fujian, Guangdong, Hunan, Guangxi, Guizhou, Sichuan, Yunnan; JAPAN, COSMOPOLITAN] Host: grains, Chinese medicine.
Holoparamesus ellipticus Wollaston, 1874 [CHINA: Jilin, Liaoning, Inner Mongolia, Hebei, Henan, Shandong, Shanxi, Shaanxi, Qinghai, Gansu, Hubei, Anhui, Jiangsu, Jiangxi, Zhejiang, Fujian, Guangdong, Hunan, Guangxi, Guizhou, Sichuan, Yunnan; JAPAN] Host: rice, *Zea mays*.
Holoparamesus signatus Wollaston, 1874 [CHINA: Hubei, Guangxi; JAPAN]

Family Mychothenidae

- Idiophyes niponensis* (Gorham, 1895) [CHINA (?); JAPAN]

Family Coccinellidae

- Aaages prior* Barovsky, 1926 [CHINA: Inner Mongolia, Gansu, Sichuan; MONGOLIA]
Adalia bipunctata (Linnaeus, 1758) (*Coccinella*) Syn. *Adalia fasciatopunctata* (Faldermann, 1835) [CHINA: Heilongjiang, Jilin, Liaoning, Inner Mongolia, Shaanxi, Gansu, Qinghai, Xinjiang, Ningxia, Shanxi, Hubei, Henan, Jiangsu, Jiangxi, Zhejiang, Fujian, Sichuan, Yunnan, Xizang; ASIA, ASIA MINOR, CAUCASUS, SIBERIA, EUROPE, N. AMERICA, C. & N. AFRICA] Prey on: *Aphis gossypii* Glover, *Aphis robiniae* Macchiati, *Icerya purchasi* Maskell, *Schizaphis graminum* (Rondani), *Hyalopterus pruni* (Geoffroy), *Rhopalosiphum oryzae* (Matsumura).
Adalia bipunctata duodecimpustulata Dobzansky, 1927 [CHINA: Hebei] Prey on: *Icerya purchasi* Maskell, *Aphis robiniae* Mac-

chiati.

- Adalia bipunctata* inaequalis Weise, 1879 [CHINA: Beijing]
Adalia lenticula Gorham, 1892 [CHINA: C. CHINA]
Adalia luteopicta (Mulsant, 1866) (*Coccinella*) [CHINA: Sichuan, Yunnan, Xizang; INDIA, SIKKIM, HIMALAYA]
Adalia tetraspilota (Hope, 1831) (*Coccinella*) Syn. *Adalia hopei* Mulsant, 1850 [CHINA: Xinjiang; INDIA, SIKKIM, NEPAL, AFGHANISTAN, TURKEY, IRAN, MONGOLIA. C. ASIA]
Afidenta arisana Li et Cook, 1961 [CHINA: Taiwan, Guangxi]
Afidenta misera (Weise, 1900) (*Epilachna*) Syn. *Afidenta mimetica* Dieke, 1947 [CHINA: Shandong, Hubei, Anhui, Jiangxi, Fujian, Taiwan, Guangdong, Hunan, Guangxi, Sichuan, Yunnan, Xizang; VIETNAM, SRI LANKA. INDIA, NEPAL, INDONESIA] Host: *Juglans regia*, *Pyrus*, *Glycine max*, *Rubus*.
Afidenta siamensis (Dieke, 1947) (*Afissa*) [CHINA: Yunnan; THAILAND]
Afidentula bisquadrupunctata (Gyllenhal, 1808) (*Coccinella*) [CHINA: Guangdong, Guangxi, Sichuan, Yunnan; INDIA, SRI LANKA] Host: Graminae, *Apluda mutica* Linnaeus.
Afidentula decimmaculata Cao et Wang, 1990 [CHINA: Yunnan] Host: *Colquhounia* sp.
Afidentula himalayana Kapur, 1963 [CHINA: Xizang; INDIA, SIKKIM]
Afidentula manderstjernae (Mulsant, 1853) (*Epilachna*) [CHINA: Guangxi, Sichuan, Yunnan; INDIA, SIKKIM, MYANMAR, HIMALAYA]
Afidentula quinquedecimguttata (Dieke, 1947) (*Afissa*) [CHINA: Hubei, Sichuan, Yunnan] Host: *Phyllostachys pubescens*, *Crinum asiaticum* var. *ainicum*.
Afissula expansa (Dieke, 1947) (*Afissa*) [CHINA: Sichuan, Yunnan] Host: *Elastostema*.
Afissula hydrangea Pang et Mao, 1979 Syn. *Afissula kambaitana* (Bielawski, 1966) [CHINA: Sichuan, Yunnan] Host: *Viburnum macrocephalum* f. *keteleeri*.
Afissula mysticoides (Sicard, 1912) (*Solanophila*) [CHINA: Xizang; INDIA, NEPAL]
Afissula rana Kapur, 1955 [CHINA: Xizang; NEPAL] Host: *Urtica*.
Afissula sanscrita (Crotch, 1874) (*Epilachna*) [CHINA: Shandong, Jiangxi, Xizang; INDIA, SIKKIM]
Afissula spatulata Cao et Wang, 1984 [CHINA: Yunnan]
Afissula uniformis Pang et Mao, 1979 [CHINA: Hunan, Sichuan, Yunnan, Xizang] Host: *Vitis*.
Afissula yunnanica Cao et Wang, 1990 [CHINA: Yunnan] Host: *Elsholtzia ciliata*.
Aiolocaria hexaspilota (Hope, 1831) (*Ithone*) Syn. *Aiolocaria mirabilis* (Motschulsky, 1860); *Aiolocaria mirabilis innocens* Mader, 1931 [CHINA: Heilongjiang, Jilin, Liaoning, Inner Mongolia, Hebei, Henan, Shanxi, Shaanxi, Gansu, Qinghai, Xinjiang, Hubei, Jiangxi, Zhejiang, Fujian, Taiwan, Hunan, Guizhou, Sichuan, Yunnan, Xizang; INDIA, MYANMAR, SIKKIM, HIMALAYA, NEPAL, KASHMIR, JAPAN, KOREA, SIBERIA] Prey on: mites, scale insects, aphids, *Gastrolina depressa* *depressa* Baly, *Podontia lutia* (Olivier), *Chrysomela adamsi* Baly.
Alloneda dodecaspilota (Hope, 1831) (*Coccinella*) Syn. *Alloneda dodecastigma* Cao et Pu, 1985 [CHINA: Yunnan, Xizang; MYANMAR, SIKKIM, NEPAL, SRI LANKA, VIETNAM, BHUTAN,] Host: aphids.
Alloneda osawai Sasaji, 1986 [CHINA: Taiwan]
Amida formosana Kurisaki, 1920 [CHINA: Taiwan]
Amida tricolor (Harold, 1878) Syn. *Amida tricolor* var. *formosana* Weide, 1923 [CHINA: Taiwan, Guangdong, Guangxi; JAPAN, VIETNAM]
Anatis ocellata (Linnaeus, 1758) (*Coccinella*) Syn. *Cocinella quindecimpunctata* De Geer, 1775 [CHINA: Heilongjiang, Jilin,

- Liaoning, Inner Mongolia, Hebei, Shanxi, Anhui, Zhejiang; MONGOLIA, Former USSR, EUROPE, C. ASIA, E. ASIA, JAPAN] Prey on: aphids on Pinus.
- Anisosticta kobensis* Lewis, 1896 Syn. *Coccinella novemdecim-punctata* (Linnaeus, 1758) [CHINA: Beijing, Henan, Shaanxi, Shandong, Xinjiang, Hubei, Jiangsu, Jiangxi, Hunan; KOREA, JAPAN, EUROPE, N. AMERICA, Former USSR] Prey on: *Aphis gossypii* Glover, *A. craccivora* Koch, *Macrosiphum avenae* Fabricius, *Rhopalosiphum maidis* (Fitch), aphids on *Scirpus*, eggs of corn borer.
- Anisosticta strigata* (Thunberg, 1795) [CHINA; N. EUROPE, MONGOLIA, SIBERIA, ALASKA]
- Anisosticta terminassianae* Beilawski, 1959 [CHINA: Hebei, Henan, Gansu, Shandong, Fujian; MONGOLIA, Former USSR] Prey on: aphids.
- Arawana isensis* (Kamiya, 1966) [CHINA: Sichuan, Yunnan; JAPAN]
- Asemiadalia heydeni* (Weise, 1892) (*Hippodamia*) [CHINA: Hubei, Jiangsu, Sichuan, Xizang; KASHMIR, C. ASIA, AFGHANISTAN, FERGANA, Former USSR]
- Asemiadalia lixianensis* Jing, 1986 [CHINA: Sichuan, Yunnan, Xizang]
- Asemiadalia rickmersi* (Weise, 1901) (*Hippodamia*) [CHINA: Xizang; C. ASIA]
- Asemiadalia spiculimaculata* Jing, 1986 [CHINA: Shaanxi, Hubei, Sichuan]
- Aspidimerus decemmaculatus* Pang et Mao, 1979 [CHINA: Yunnan]
- Aspidimerus esakii* Sasaji, 1968 [CHINA: Anhui, Taiwan, Guangxi] Prey on: aphids.
- Aspidimerus matsumurai* Sasaji, 1968 [CHINA: Henan, Hubei, Zhejiang, Taiwan, Guangxi, Sichuan, Yunnan; VIETNAM] Host: *Toxoptera citricida* (Kirkaldy).
- Aspidimerus ruficus* Gorham, 1895 [CHINA: Gansu, Guangxi, Sichuan, Yunnan; MYANMAR] Prey on: aphids.
- Aspidimerus sexmaculatus* Pang et Mao, 1979 [CHINA: Yunnan]
- Axinoscymnus beneficus* Kamiya, 1963 [CHINA: Taiwan, Hainan, Guangdong; JAPAN]
- Axinoscymnus nigripennis* Kamiya, 1965 [CHINA: Taiwan, Guangdong, Guangxi; JAPAN]
- Ballida brahmae* Mulsant, 1850 [CHINA]
- Bothrocalvia albolineata* (Gyllenhal, 1808) (*Coccinella*) Syn. *Halysia albolineata* Miwa, 1931; *Calvia albolineata* Liu, 1963 [CHINA: Henan, Jiangxi, Zhejiang, Fujian, Taiwan, Guangdong, Hongkong, Hunan, Guangxi, Guizhou, Sichuan, Yunnan; INDIA, JAPAN] Prey on: *Schizaphis graminum* (Rondani), *Cinara pini-formosana* Takahashi, *Lipaphis erysimi* (Kaltenbach).
- Bothrocalvia lewisi* (Crotch, 1874) (*Calvia*) [CHINA: Fujian, Guangdong, Guangxi, Yunnan; MYANMAR, CAMBODIA] Prey on *Castanea mollissima*.
- Bothrocalvia pupillata* (Swartz, 1808) (*Coccinella*) Syn. *Coleophora pusillata* (Swartz, 1808); *Coleophora pupillata* Mader, 1926 [CHINA: Guangdong, Hongkong, Hainan, Guangxi, Sichuan, Yunnan; INDIA, INDONESIA] Prey on: aphids.
- Brumoides hainanensis* Miyatake, 1970 [CHINA: Fujian, Guangdong, Hainan, Guangxi, Yunnan] Prey on: *Schizaphis graminum* (Rondani), rice planthoppers, aphids.
- Brumoides lineatus* (Weise, 1885) (*Brumus*) [CHINA: Taiwan, Hainan (Kisha I.), Guangxi; INDIA, MYANMAR, BANGLADESH, THAILAND, SRI LANKA] Prey on: *Aphis craccivora* Koch, *Longiunguis sacchari* (Zehntner), aphids.
- Brumoides maai* Miyatake, 1970 [CHINA: Fujian, Guangxi, Yunnan] Prey on: aphids.
- Brumoides ohtai* Miyatake, 1970 [CHINA: Taiwan; JAPAN]
- Brumus bifasciatus* Barovsky, 1927 [CHINA: Xinjiang]
- Brumus jacobsoni* Barovsky, 1927 [CHINA: Inner Mongolia]
- Brumus mongolicus* Fleischer, 1900 [CHINA: Inner Mongolia; TRANSBAIKALIA]
- Bulaea lichatschovi* (Hummel, 1827) (*Coccinella*) [CHINA: Xinjiang; Former USSR, POLLAND, INDIA, AFGHANISTAN, PENINSULA OF ARABIA, EUROPE, AFRICA] Host: *Beta vulgaris*.
- Bulaea nevillei* Dohrn, 1882 [CHINA; INDIA, NEPAL]
- Callicaria superba* (Mulsant, 1853) (*Caria*) Syn. *Callicaria superba japonica* Sicard, 1907; *Synonycha japonica* Kurisaki, 1915 [CHINA: Shaanxi, Gansu, Hubei, Zhejiang, Taiwan, Guizhou, Sichuan, Yunnan, Xizang; INDIA, BHUTAN, HIMALAYA, JAPAN] Prey on: *Aphanostigma iakusuiensis* Kishida, *Prociophilus kuwanai* Monzen, woolly pear aphid.
- Calvia chinensis* (Mulsant, 1850) (*Sospita*) [CHINA: Hebei, Henan, Shaanxi, Jiangsu, Jiangxi, Zhejiang, Fujian, Guangdong, Hongkong, Hainan, Guangxi, Guizhou, Sichuan, Yunnan, SW CHINA, C. CHINA] Prey on: *Rhopalosiphum oryzae* (Matsumura), aphids on *Pennis*.
- Calvia decemguttata* (Linnaeus, 1767) (*Coccinella*) [CHINA: Sichuan; JAPAN, MYANMAR, ASIA, RUSSIA, EUROPE]
- Calvia duodecimmaculata* (Gehler, 1832) (*Coccinella*) Syn. *Coccinella duodecimmaculata yasui* Nakane, 1954; *Calvia pasupati* Kapur, 1963 [CHINA: Hebei, Shaanxi, Sichuan, Xizang; INDIA, SRI LANKA, JAPAN, Former USSR, N. AMERICA] Prey on: aphids.
- Calvia hauseri* (Mader, 1930) (*Anisocalvia*) [CHINA: Gansu, Guizhou, Sichuan, Yunnan]
- Calvia krishna* (Crotch, 1874) (*Anisocalvia*) [CHINA: Yunnan; VIETNAM, SIKKIM]
- Calvia muiri* (Timberlake, 1943) (*Eocalvia*) [CHINA: Hebei, Shaanxi, Henan, Hubei, Jiangxi, Fujian, Taiwan, Guizhou, Sichuan, Yunnan; JAPAN] Prey on: *Lipaphis erysimi* (Kaltenbach), *Schizaphis graminum* (Rondani), aphids.
- Calvia quatuordecimguttata* (Linnaeus, 1758) (*Coccinella*) [CHINA: Jilin, Shaanxi, Hubei, Taiwan; JAPAN, KOREA, SIBERIA, SACHALIN, KURILE, EUROPE, N. AMERICA]
- Calvia quinquedecimguttata* (Fabricius, 1777) (*Coccinella*) Syn. *Calvia quinquedecimguttata ab. yunnan* Mader, 1931 [CHINA: Henan, Shaanxi, Gansu, Xinjiang, Shandong, Hubei, Anhui, Jiangsu, Jiangxi, Zhejiang, Fujian, Taiwan, Hunan, Guangxi, Guizhou, Sichuan, Yunnan, Xizang; INDIA, JAPAN, SIBERIA, SACHALIN, KURILE, C. ASIA, CAUCASUS, KOREA, EUROPE] Prey on: *Toxoptera citricida* (Kirkaldy), *Ceratovacuna lanigera* Zehntner.
- Calvia shirozui* (Miyatake, 1965) (*Eocaria*) Syn. *Halysia quindecimguttata septenaria* Mulsant, 1866 [CHINA: Mainland of CHINA, Taiwan]
- Calvia sicardi* (Mader, 1930) (*Anisocalvia*) [CHINA: Shaanxi, Gansu, Hubei, Hunan, Guizhou, Sichuan, Yunnan] Prey on: *Aphis gossypii* (Glover).
- Catana yasumatsui* Sasaji, 1967 [CHINA: Taiwan]
- Catanella formosana* Miyatake, 1961 [CHINA: Taiwan]
- Cheilomenes sexmaculata* (Fabricius, 1781) (*Chilomenes*) [CHINA: Shaanxi, Henan, Jiangxi, Fujian, Taiwan, Hunan, Guangdong, Guangxi, Guizhou, Sichuan, Yunnan; S. ASIA, INDIA, SE ASIA, PHILIPPINES, SRI LANKA, INDONESIA, MICRONESIA] Prey on: aphids. Natural enemy: *Homalotylus flaminus* Dalman.
- Chilocorus alishanus* Sasaji, 1968 [CHINA: Taiwan, Guizhou, Yunnan] Prey on: scale insects.
- Chilocorus bijugus* Mulsant, 1853 Syn. *Chilocorus renipustulatus* (Scriba, 1790) [CHINA: Gansu, Hubei, Jiangsu, S. CHINA,

- Guizhou, Sichuan, Yunnan, Xizang; JAPAN, INDIA, PAKISTAN, NEPAL, PALAEARCTIC] Prey on: *Aspidiotus destructor* Signoret, *Chrysomphalus aonidum* (Linnaeus), *Coccus hesperidum* Linnaeus, *Parlatoria pergandei* Comstock, *P. zizyphus* (Lucas), *Pseudaulacaspis pentagona* (Targioni - Tozzetti), *Pulvinaria citricola* Kuwana, *Unaspis yanonensis* (Kuwana).
- Chilocorus bipustulatus* (Linnaeus, 1758) (*Coccinella*) [CHINA: Gansu, Xinjiang; PAELARCTIC] Prey on: *Aphis robiniae* Macchiati, *Eriosoma lanigerum* (Hausmann), *Coccus hesperidum* Linnaeus, *Icerya purchasi* Maskell, *Lepidosaphes bechii* (Newman), *L. gloveri*.
- Chilocorus chalybeatus* Gorham, 1892 [CHINA: Shaanxi, Jiangxi, Zhejiang, Fujian, Guangdong, Hunan, Guangxi, Guizhou, Sichuan] Prey on: *Ceroplastes rubens* Maskell, scale insects.
- Chilocorus chinensis* Miyatake, 1970 [CHINA: Zhejiang, Fujian, Guangdong, Guangxi, Yunnan] Prey on: scale insects.
- Chilocorus circumdatus* (Gyllenhal, 1808) (*Coccinella*) [CHINA: Zhejiang, Fujian, Guangdong, Hainan, Guangxi, Yunnan; INDIA, SRI LANKA, INDONESIA] Prey on: scale insects.
- Chilocorus esakii* Kamiya, 1959 [CHINA: Liaoning, Shanxi, Hebei, Gansu, Shandong, Jiangsu, Sichuan; JAPAN] Prey on: *Pseudaulacaspis pentagona* (Targioni), scale insects, whiteflies.
- Chilocorus geminus* Zaslavskij, 1962 [CHINA: Inner Mongolia, Gansu, Xinjiang; MONGOLIA, TURKEY, UZBEKISTAN] Prey on: aphids, scale insects.
- Chilocorus gressitti* Miyatake, 1970 [CHINA: Fujian, Guangdong, Hainan, Guangxi, Sichuan, Yunnan; MYANMAR, INDIA, THAILAND, NEPAL, SRI LANKA, INDONESIA] Prey on: *Chrysomphalus bifasciatus* Ferris, *Ch. aonidum* (Linnaeus), *Temnaspidiotus destructor* (Signoret).
- Chilocorus hauseri* Weise, 1895 [CHINA: Shaanxi, Hubei, Fujian, Guangdong, Hainan, Guizhou, Sichuan, Yunnan; SIKKIM, MYANMAR] Prey on: *Oligonychus coffeae* (Nietner), *Quadraspidiotus perniciosus* Comstock, *Temnaspidiotus destructor* (Signoret), scale insects.
- Chilocorus hupehanus* Miyatake, 1970 [CHINA: Gansu, Hubei, Fujian, Hunan, Guangxi, Guizhou, Sichuan] Prey on: *Chrysomphalus aonidum* (Linnaeus), *Ch. bifasciatus* Ferris, *Eriococcus kaki* Kuwana, *Aulacaspis crawsi* Gkll., *Pseudaulacaspis pentagona* (Targioni), *Unaspis yanonensis* (Kuwana).
- Chilocorus kuwanae* Silvestri, 1909 Syn. *Chilocorus similis* Rossi, 1790 [CHINA: Heilongjiang, Jilin, Liaoning, Hebei, Henan, Shanxi, Shaanxi, Gansu, Shandong, Hubei, Anhui, Jiangsu, Jiangxi, Zhejiang, Hunan, Fujian, Guangdong, Guizhou, Sichuan, Yunnan, Xizang; JAPAN, KOREA, MONGOLIA, INDIA, ITALY, INDIA, introduced to U. S and ITALY] Prey on: *Ericerus pela* Chavannes.
- Chilocorus nigrinus* (Fabricius, 1798) [CHINA: Fujian, Guangdong, Guangxi, Sichuan, Yunnan] Prey on: scale insects.
- Chilocorus politus* Mulsant, 1850 [CHINA: Yunnan, Xizang; INDIA, NEPAL, THAILAND, INDONESIA] Prey on: scale insects, aphids.
- Chilocorus rubidus* Hope, 1831 [CHINA: Heilongjiang, Jilin, Liaoning, Inner Mongolia, Hebei, Shaanxi, Gansu, Ningxia, Henan, Shandong, Hubei, Jiangsu, Jiangxi, Zhejiang, Fujian, Hainan, Hunan, Guizhou, Sichuan, Yunnan, Xizang; MONGOLIA, JAPAN, KOREA, USSURI, SIBERIA, NEPAL, INDIA, CELEBES, INDONESIA, AUSTRALIA] Prey on: *Ericerus pela* Chavannes.
- Chilocorus rufitarsus* Motschulsky, 1853 [CHINA: Henan, Jiangxi, Zhejiang, Fujian, Guangdong, Guangxi, Guizhou, Sichuan, Yunnan; VIETNAM] Prey on: *Didesmococcus koreanus* Borchs., *Schizaphis piricola* matsumura, scale insects.
- Chilocorus shirozui* Sasaji, 1968 [CHINA: Taiwan]
- Chilocorus yunlongensis* Cao et Xiao, 1984 [CHINA: Yunnan] Prey on: scale insects.
- Clitostethus sternalis* (Pang et Gordon, 1986) (*Pseudoscymnus*) [CHINA: Guangdong, Hainan]
- Coccidula rufa* (Herbst, 1783) (*Chrysomela*) [CHINA: Henan, Xinjiang, Sichuan; ASIA, Former USSR, EUROPE] Prey on: aphids.
- Coccidula unicolor* Reitter, 1890 [CHINA: Henan, Xinjiang; IRAN, TURKEY] Prey on: aphids.
- Coccinella ainu* Lewis, 1896 [CHINA: Shaanxi, Gansu, Xinjiang, Hubei; JAPAN, KOREA, SIBERIA, EUROPE, N. AFRICA] Prey on: *Myzus persicae* (Sulzer).
- Coccinella divaricata* Olivier, 1808 [CHINA: Inner Mongolia, Hebei, Shandong, Shaanxi, Xinjiang, Jiangsu; EUROPE, ASIA, N. AFRICA, PALAEARCTIC] Prey on: *Schizaphis graminum* (Rondani), *Aphis robiniae* Macchiati, aphids on weeds.
- Coccinella divaricata* dohzanskii Mader, 1926 [CHINA: Gansu]
- Coccinella longifasciata* Liu, 1962 Syn. *Coccinella nigrovittata* Kapur, 1963; *Tythuspis trilineata* Weise, 1889 [CHINA: Jilin, Qinghai, Gansu, Xinjiang, Sichuan, Xizang] Prey on: aphids.
- Coccinella luteopicta* (Mulsant, 1866) (*Adalia*) [CHINA: Sichuan, Yunnan, Xizang; INDIA, SIKKIM, MYANMAR]
- Coccinella magnifica* Redtenbacher, 1843 Syn. *Coccinella lama* Kapur, 1963 [CHINA: Xizang] Prey on: *Schizaphis graminum* (Rondani), *Lipaphis erysimi* (Kaltenbach), aphids.
- Coccinella magnopunctata* Rybakow, 1889 Syn. *Coccinella undecimpunctata* var. *magnopunctata* Rybakov, 1889; *Coccinella semenovi* Weise, 1889; *Coccinella undecimpunctata* Dobzhansky, 1926; *Coccinella undecimpunctata magnopunctata* Mader, 1926 [CHINA: Gansu, Qinghai, Xinjiang, Guizhou, Sichuan, Xizang; KASHMIR, MONGOLIA, Former USSR, PALAEARCTIC] Prey on: aphids.
- Coccinella mannerheimi* Mulsant, 1850 [CHINA: NE CHINA, Inner Mongolia; Former USSR]
- Coccinella nivicola* Mulsant, 1850 [CHINA: Inner Mongolia, Xinjiang; Former USSR]
- Coccinella quinquepunctata* Linnaeus, 1758 [CHINA: Gansu; JAPAN, SIBERIA, MONGOLIA, ASIA, CAUCASUS, C. ASIA, EUROPE, AFRICA]
- Coccinella reitter*, Weise, 1891 [CHINA: Xinjiang; C. ASIA, KAZAKHSTAN]
- Coccinella sachalinensis* Ohta, 1928 [CHINA: Gansu; SACHALIN]
- Coccinella saucerottii* Mulsant, 1850 Syn. *Coccinella bodemeyeri* Weise, 1900; *Coccinella lushniki* Dobzhansky, 1917; *Coccinella lingenbergi* Kunnemann, 1917; *Coccinella tibetina* Kapur, 1963 [CHINA: Sichuan, Yunnan, Xizang; RUSSIA, EUROPE, PALAEARCTIC]
- Coccinella sauteri* klingenbergi Kunnemann, 1917 [CHINA: Xinjiang]
- Coccinella sedakovi* Mulsant, 1850 Syn. *Coccinella tianshanica* Dobzhansky, 1927 [CHINA: Xinjiang; AFGHANISTAN, MONGOLIA, C. ASIA, TURKEY, Former USSR]
- Coccinella septempunctata* Linnaeus, 1758 Syn. *Coccinella confusa* Wiedemann, 1823; *Coccinella septempunctata brucki* Mulsant, 1866 [CHINA: Heilongjiang, Jilin, Liaoning, Inner Mongolia, Hebei, Shandong, Shanxi, Shaanxi, Ningxia, Gansu, Qinghai, Xinjiang, Henan, Hubei, Anhui, Jiangsu, Jiangxi, Zhejiang, Fujian, Taiwan, Guangdong, Hunan, Guangxi, Guizhou, Sichuan, Yunnan, Xizang; JAPAN, KOREA, MONGOLIA, Former USSR, INDIA, EUROPE, U. S., AFRICA, PALAEARCTIC, COSMOPOLITAN] Prey on: *Aphis gossypii* Glover, *A. craccivora* Koch, *A. glycines* Matsumura, *A. robiniae* Macchiati, *Anomoneura mori* Schwarz, *Aphanostigma iakusuiensis* Kishida, *Lipaphis erusimi* (Kaltenbach), *Myzus persicae* (Sulzer), *M. malisuctus* Matsumura, *M. varians* Davidson, *Schizaphis piricola* Matsumura,

- Hyalopterus arundinis* Fabricius, *Sappaphis pirocola* Okamoto et Takahashi, *Brevicorne brassicae* (Linnaeus), *Schizaphis graminum* (Rondani), eggs and larvae of *Heliothis armigera* (Hubner). Natural enemy: *Hanalotylus flammiosus* Dalman.
- Coccinella takasago* Sasaji, 1982 [CHINA; Taiwan]
- Coccinella transversalis* Fabricius, 1781 Syn. *Coccinella repanda* Thunberg, 1781 [CHINA: Shaanxi, Hubei, Jiangxi, Fujian, Taiwan, Guangdong, Hongkong, Hainan, Hunan, Guangxi, Guizhou, Sichuan, Yunnan, Xizang; INDIA, VIETNAM, NEPAL, BHUTAN, SRI LANKA, THAILAND, MYANMAR, BANGLADESH, MALAYSIA, INDONESIA, PHILIPPINES, AUSTRALIA, NEW GUINEA, NEW ZEALAND, NEW CALEDONIA, TASMANIA] Prey on: *Aphis glycines* Matsumura, *Ceratovacuna lanigera* Zehntner, *Lipaphis erysimi* (Kaltenbach), *Longiumguis sacchari* (Zehntner), *Schizaphis graminum* (Rondani), larvae of *Lepidoptera* on rice, *Ioerya purchasi* Maskell.
- Coccinella transversoguttata* Faldermann, 1835 Syn. *Coccinella geminopunctata* Liu, 1962; *Coccinella transversoguttata ab. bini-terrupta* Mader, 1926; *Coccinella transversoguttata nugatoria* Mulsant, 1850; *Coccinella transversoguttata richardsoni* Brown, 1967 [CHINA: Jilin, Liaoning, Inner Mongolia, Hebei, Shanxi, Ningxia, Shaanxi, Gansu, Qinghai, Xinjiang, Hubei, Jiangxi, Zhejiang, Guangdong, Sichuan, Yunnan, Xizang; JAPAN, SIBERIA, MONGOLIA, C. ASIA, EUROPE, N. AMERICA, MEXICO] Prey on: *Schizaphis graminum* (Rondani).
- Coccinella trifasciata* Linnaeus, 1758 Syn. *Coccinella trifasciata perplexa* Mulsant, 1850; *Coccinella trifasciata subversa* LeConte, 1854 [CHINA: Heilongjiang, Jilin, Inner Mongolia, Hebei, Ningxia, Shaanxi, Qinghai, Gansu, Xinjiang, Sichuan, Xizang; MONGOLIA, Former USSR, EUROPE, N. AMERICA] Prey on: *Schizaphis graminum* (Rondani), *Aphis farinosa* Gmelin.
- Coccinella undecimpunctata* Linnaeus, 1758 Syn. *Coccinella menetriesi* Mulsant, 1850; *Coccinella pontica* Dobzhansky, 1927 [CHINA: Inner Mongolia, Hebei, Shanxi, Ningxia, Shaanxi, Gansu, Xinjiang, Henan, Shandong, Guizhou, Sichuan; ASIA, EUROPE, N. AFRICA] Prey on: *Aphis gossypii* Glover, *Schizaphis graminum* (Rondani).
- Coccinella weisei* (Rybakov, 1889) (*Micraspis*) [CHINA; TURKEY]
- Coccinella withi* Mulsant, 1850 [CHINA; NE CHINA; TURKEY]
- Coccinella elegantula* (Weise, 1890) (*Coccinella*) [CHINA: Jilin, Inner Mongolia, Qinghai, Xinjiang, Sichuan; KAZAKHSTAN, MONGOLIA, SIKKIM, C. ASIA] Prey on: aphids.
- Coccinula quatuordecimpustulata* (Linnaeus, 1758) (*Coccinella*) [CHINA: Heilongjiang, Jilin, Liaoning, Hebei, Inner Mongolia, Shanxi, Ningxia, Gansu, Xinjiang, Henan, Shandong, Jiangxi, Sichuan; JAPAN, Former USSR, EUROPE]
- Coccinula sinensis* (Weise, 1889) (*Coccinella*) [CHINA: Sichuan; MONGOLIA, JAPAN, Former USSR,] Prey on: *Aphis medicaginis* Koch, *Schizaphis graminum* (Rondani), *Rhopalosiphum maidis* Fitch.
- Coccinula sinuatmarginata* (Faldermann, 1837) [CHINA: Xinjiang; CAUCASUS, SIBERIA, C. ASIA, EUROPE]
- Coleophora cincta* Hope, 1831? Syn. *Coleophora cincta ab. pristina* Mader, 1939 [CHINA; NEPAL]
- Coleophora flavomarginata* Sasaji, 1982 [CHINA; Taiwan]
- Coleophora inaequalis* (Fabricius, 1775) (*Coccinella*) Syn. *Coccinella novemmaculata* Fabricius, 1781; *Coccinella iridea* Thunberg, 1781; *Coccinella psi* Thunberg, 1781; *Coccinella octonotata* Fabricius, 1801; *Coccinella picta* Fabricius, 1806; *Coccinella patruelis* Boisduval, 1835; *Coleophora desolata* Mulsant, 1850; *Coleophora reniplagiata* Mulsant, 1850; *Coleophora partita* Mulsant, 1850; *Coleophora mendica* Mulsant, 1850; *Coleophora symbolica* Mulsant, 1850; *Coleophora musae* Montrouzier, 1857; *Coleophora flachati* Mulsant, 1866; *Coleophora caledonica* Crotch, 1874; *Coleophora nipponi* Crotch, 1874; *Coleophora mastersi* Blackburn, 1892; *Coleophora veranoides* Blackburn, 1894; *Coleophora fasciata* Weise, 1902; *Coleophora consortula* Weise, 1902; *Coleophora rouxi* Heyden, 1911; *Coleophora kochi* Weise, 1913; *Coleophora suturella* Weise, 1913; *Coleophora comperei* Tomberlake, 1943 [CHINA: Guangdong, Taiwan; INDIA, INDONESIA, THAILAND, PHILIPPINES, MALAYSIA, CALEMANDAN, CELEBES, MICRONESIA, HAWAII, AUSTRALIA, NEW GUINEA]
- Coleophora itoi* Sasaji, 1982 [CHINA; Taiwan]
- Coleophora sexguttata* Sewise, 1910 [CHINA; Taiwan; PHILIPPINES]
- Coleophora subustulata* Gorham, 1891 [CHINA; S. CHINA]
- Coleophora tanoi* Sasaji, 1982 [CHINA; Taiwan]
- Cryptogonus angusticarinatus* Sasaji, 1968 [CHINA; Taiwan, Guangdong; VIETNAM]
- Cryptogonus bimaculatus* Kapur, 1948 [CHINA: Yunnan, Xizang; INDIA, MYANMAR] Prey on: aphids.
- Cryptogonus blandus* Mader, 1954 [CHINA: Yunnan] Prey on: aphids, scale insects.
- Cryptogonus complexus* Kapur, 1948 [CHINA: Gansu, Taiwan, Guangdong, Guangxi, Sichuan, Yunnan; INDIA, MYANMAR, VIETNAM, SIKKIM]
- Cryptogonus forcliculae* Cao et Xiao, 1984 [CHINA: Yunnan] Prey on: aphids on Pinus.
- Cryptogonus guangdongensis* Pang et Mao, 1979 [CHINA: Jiangxi, Guangdong, Hainan, Guangxi] Prey on: *Aphis craccivora* Koch, *Toxoptera citricida* (Kirkaldy).
- Cryptogonus hainanensis* Pang et Mao, 1979 [CHINA: Hainan]
- Cryptogonus himalayensis* Kapur, 1948 [CHINA: Guangdong, Yunnan; MYANMAR, INDIA] Prey on: aphids.
- Cryptogonus horishanus* (Ohta, 1929) (*Scymnus*) Syn. *Pseudaspidimerus japonensis* Nakane et Araki, 1958 [CHINA: Gansu, Hubei, Jiangxi, Zhejiang, Fujian, China; CHINA: Taiwan, Guangdong, Sichuan; JAPAN, INDONESIA] Prey on: *Toxoptera citricida* (Kirkaldy), *Myzus persicae* (Sulz).
- Cryptogonus kurosawai* Sasaji, 1968 [CHINA: Taiwan]
- Cryptogonus lijiangensis* Pang et Mao, 1979 [CHINA: Shaanxi, Gansu, Hubei, Yunnan] Prey on: aphids on *Juglans regia*.
- Cryptogonus nepalensis* Bielawski, 1972 [CHINA: Yunnan, Xizang; NEPAL] Prey on: aphids.
- Cryptogonus nigrinus* Pang et Mao, 1979 [CHINA: Yunnan]
- Cryptogonus octoguttatus* Mader, 1954 [CHINA: Sichuan, Yunnan]
- Cryptogonus ohtai* Sasaji, 1968 [CHINA; Taiwan]
- Cryptogonus orbiculus* (Gyllenhal, 1808) (*Coccinella*) Syn. *Platynaspis oculata* Motschulsky, 1858; *Cryptogonus centrogruttatus* Boheman, 1859; *Aspidimerus orbiculus* Gyllenhal, 1808; *Aspidimerus orbiculus nigripennis* Weise, 1923; *Dionus futahoshii* Ohta, 1929; *Dionus futahoshii ab. koshunus* Ohta, 1929; *Cryptogonus malasiae* Crotch, 1931; *Cryptogonus orbiculus ab. japonicus* Weise, 1900 [CHINA: Shaanxi, Gansu, Hubei, Jiangxi, Zhejiang, Fujian, Taiwan, Guangdong, Hongkong, Hainan, Hunan, Guangxi, Guizhou, Sichuan, Yunnan; JAPAN, SE ASIA, INDIA, MYANMAR, MALAYSIA, SRI LANKA, INDONESIA, MICRONESIA, CALEMANDAN] Prey on: *Toxoptera citricida* (Kirkaldy), *Aphis craccivora* Koch, *A. gossypii* Glover, scale insects.
- Cryptogonus postmedialis* Kapur, 1948 [CHINA: Hubei, Fujian, Taiwan, Guangdong, Sichuan; INDIA, MYANMAR, SIKKIM] Prey on: *Aphis gossypii* Glover.
- Cryptogonus quadriguttata* (Weise, 1895) (*Aspidimerus*) [CHINA: Taiwan, Yunnan; INDIA, MYANMAR, SIKKIM, SRI LANKA]
- Cryptogonus sagittiformis* Pang et Mao, 1979 [CHINA: Yunnan]

- Cryptogonus schraiki* Mader, 1933 Syn. *Cryptogonus schraiki* var. *urbani* Mader, 1933 [CHINA: Hubei, Sichuan, Yunnan] Prey on: aphids, scale insects.
- Cryptogonus trifurcatus* Pang et Mao, 1979 [CHINA: Yunnan] Prey on: aphids on Ficus.
- Cryptogonus trioblitus* (Gorham, 1895) (*Aspidimerus*) [CHINA: Yunnan, Xizang; MYANMAR, INDIA, NEPAL] Prey on: aphids, scale insects.
- Cryptogonus wuzhishanus* Pang et Mao, 1979 [CHINA: Shaanxi, Henan, Hainan, Sichuan] Prey on: *Unaspis yanoensis* (Kuwana).
- Cryptogonus yunnanensis* Cao et Xiao, 1984 [CHINA: Yunnan] Prey on: aphids on Pinus.
- Cryptolaemus montrouzieri* Mulsant, 1853 [CHINA: Fujian, Taiwan, Guangdong, Hongkong, Sichuan; INDIA, JAPAN, PHILIPPINES, INDONESIA, THAILAND, HAWAII, MICRONESIA, AUSTRALIA, NEW ZEALAND, AFRICA, SPAIN, EGYPT, KENYA, S. EUROPE, C. & N. AMERICA] Prey on: *Pseudococcus maritimus* (Ehrhorn), *Ps. gahani* Green, *Ps. adonidum* (Linnaeus), *Stomacoccus*.
- Delphastus pusillus* (LeCorte, 1852) [CHINA: Fujian; AMERICA (introduced from ENGLAND)] Prey on: Aleurodidae.
- Diomus brachysiphonius* Pang et Huang, 1986 [CHINA: Fujian, Hainan, Guangdong, Sichuan; VIETNAM]
- Diomus brunsturalis* Pang et Gordon, 1986 [CHINA: Shaanxi, Beijing, Zhejiang, Fujian, Taiwan, Guangdong, Sichuan; VIETNAM]
- Eoadalia indica* (Crotch, 1874) (*Adalia*) [CHINA: Sichuan, Xizang; INDIA, MYANMAR]
- Epilachna acuta* (Weise, 1899) (*Solanophila*) Syn. *Epilachna acuminata* Weise, 1889; *Solanophila acutula* Weise, 1902 [CHINA: Gansu, Jiangsu, Taiwan]
- Epilachna admirabilis* (Crotch, 1874) (*Afissa*) Syn. *Solanophila admirabilis* Mader, 1926; *Afissa admirabilis* Dieke, 1947; *Afissa admirabilis continentalis* Dieke, 1947 [CHINA: Shaanxi, Hebei, Henan, Hubei, Anhui, Jiangsu, Jiangxi, Zhejiang, Fujian, Taiwan, Guangdong, Hunan, Guangxi, Sichuan, Yunnan; JAPAN, MYANMAR, THAILAND, PHILIPPINES, SIKKIM, AUSTRALIA, INDONESIA] Host: *Solanum melongena*, *Cucumis sativus*, *Cucurbita moschata*, *Solanum melogena*, *Citrullus lanatus*, *Momordica charantia*.
- Epilachna adscita* Mader, 1930 Syn. *Solanophila adscita* ab. *priesneri*, [CHINA: Sichuan]
- Epilachna alternans* Mulsant, 1850 [CHINA: Taiwan, Sichuan; INDIA, INDONESIA]
- Epilachna ampliata* Pang et Mao, 1979 [CHINA: Yunnan]
- Epilachna angusta* Li et Cook, 1961 [CHINA: Taiwan]
- Epilachna anhweiiana* (Dieke, 1947) (*Afissa*) [CHINA: Henan, Anhui, Jiangxi, Yunnan] Host: *Styrax*.
- Epilachna bengalica* (Dieke, 1947) (*Afissa*) [CHINA: Xizang; INDIA, BANGLADESH, SIKKIM]
- Epilachna bicrescens* (Dieke, 1947) (*Afissa*) [CHINA: Hunan, Sichuan]
- Epilachna bifibra* Li et Cook, 1961 [CHINA: Taiwan]
- Epilachna bisquadripunctata herbigrada* Mulsant, 1850 [CHINA: Hongkong]
- Epilachna brivioi* (Bielawski et Fursch, 1960) (*Afissa*) [CHINA: Yunnan; MYANMAR]
- Epilachna chayuensis* Pang et Mao, 1977 [CHINA: Xizang]
- Epilachna chinensis* (Weise, 1912) (*Solanophila*) Syn. *Afissa chinensis tsushima* Nakane et Araki, 1960 [CHINA: Shaanxi, Henan, Hubei, Anhui, Jiangxi, Fujian, Taiwan, Guangdong, Hunan, Guizhou, Sichuan, Yunnan; JAPAN, KOREA] Host: *Pitosporum illicioides*, *Galium aparine* var. *terenum*, *Anemone cathayensis*.
- Epilachna circummaculata* Pang et Mao, 1977 [CHINA: Xizang]
- Epilachna clematicola* Cao et Xiao, 1984 [CHINA: Yunnan] Host: *Clematis armandii*, *Bambusa omeiensis*.
- Epilachna coalescens* Mader, 1930? [CHINA: Sichuan]
- Epilachna completa* (Dieke, 1947) (*Afissa*) [CHINA: Hunan, Sichuan]
- Epilachna complicata* (Dieke, 1947) (*Afissa*) Syn. *Afissa complicata* var. *sustenans* Dieke, 1947 [CHINA: Sichuan]
- Epilachna confusa* Li et Cook, 1961 [CHINA: Taiwan, Sichuan] Host: *Betula*.
- Epilachna convexa* (Dieke, 1947) (*Afissa*) [CHINA: Shaanxi, Hubei, Hunan, Sichuan] Host: *Anemone cathayensis*.
- Epilachna cressimala* Li et Cook, 1961 [CHINA: Taiwan, Hainan]
- Epilachna cuonaensis* Pang et Mao, 1977 [CHINA: Xizang]
- Epilachna decemguttata* (Weise, 1923) (*Solanophila*) [CHINA: Taiwan]
- Epilachna decemmaculata* Redtenbacher, 1844 [CHINA: Taiwan, Xizang; INDIA, KASHMIR, MYANMAR, SRI LANKA]
- Epilachna decipiens* Crotch, 1874 [CHINA: Taiwan; JAVA]
- Epilachna dobzhanskyi* (Dieke, 1947) (*Afissa*) [CHINA: Hubei, Jiangsu, Sichuan] Host: *Anemone cathayensis*.
- Epilachna dumerili* Mulsant, 1850 Syn. *Epilachna undecimspilota dumerili* Hope, 1831; *Solanophila dumerili discordia* Mader, 1930 [CHINA: Fujian, Taiwan, Hongkong; INDIA, MYANMAR, SIKKIM, NEPAL, THAILAND, BANGLADESH, SRI LANKA, INDONESIA]
- Epilachna elvina* Mulsant, 1850 [CHINA: Taiwan; MYANMAR, INDIA, SRI LANKA]
- Epilachna erythrotricha* Hoang, 1978 [CHINA: Guangxi; VIETNAM]
- Epilachna exoynata* Bielawski, 1965 [CHINA: Yunnan]
- Epilachna fairmairei* Frivaldsky, 1892? [CHINA]
- Epilachna fallaxi* (Weise, 1908) (*Solanophila*) [CHINA: Taiwan; INDONESIA] Host: *Boehmeria nivea*.
- Epilachna folifera* Pang et Mao, 1979 [CHINA: Jiangxi, Yunnan]
- Epilachna formosana* (Weise, 1923) (*Solanophila*) [CHINA: Taiwan] Host: *Boehmeria nivea*.
- Epilachna freyana* Bielawski, 1965 [CHINA: Shaanxi, Hubei, Jiangxi, Hainan, Hunan, Sichuan, Yunnan] Host: *Tanacetum*.
- Epilachna fugongensis* Cao et Xiao, 1984 [CHINA: Yunnan] Host: *Benincasa hispida*.
- Epilachna galerucinoidea* Korschefsky, 1934 [CHINA: Taiwan, Hainan, Yunnan; THAILAND, INDONESIA, AUSTRALIA] Host: *Vitis*.
- Epilachna gedeensis* (Dieke, 1947) (*Afissa*) [CHINA: Yunnan; JAVA]
- Epilachna glochinosa* Pang et Mao, 1979 [CHINA: Hubei, Guangxi, Sichuan, Yunnan] Host: *Akebia quinata*.
- Epilachna glochisifoliata* Pang et Mao, 1979 [CHINA: Hubei, Guangxi, Sichuan] Host: *Akebia quinata*.
- Epilachna grayi* Mulsant, 1850 [CHINA: Taiwan, Xizang; INDIA, BANGLADESH, NEPAL, INDONESIA]
- Epilachna gressitti* Li et Cook, 1961 [CHINA: Taiwan]
- Epilachna hauseri* (Mader, 1930) (*Solanophila*) [CHINA: Shandong, Guizhou, Sichuan, Yunnan]
- Epilachna hendecaspilota* (Mader, 1927) (*Solanophila*) [CHINA: Taiwan; INDIA, MYANMAR, THAILAND, SRI LANKA, PHILIPPINES, INDONESIA,]
- Epilachna hingstoni* (Kapur, 1963) (*Afissa*) [CHINA: Xizang; INDIA, NEPAL]
- Epilachna hopeiana* Miyatake, 1985 [CHINA: Guangxi; VIETNAM]
- Epilachna incauta* Mulsant, 1850 Syn. *Solanophila incauta*, Sicard, 1910 [CHINA: Taiwan; INDONESIA, MYANMAR]

- Epilachna insignis* (Corham, 1892) (*Afissa*) [CHINA: Beijing, Shaanxi, Henan, Hubei, Anhui, Jiangxi, Zhejiang, Fujian, Guangdong, Guizhou, Sichuan, Yunnan] Host: *Solanum melongena*, *S. nigrum*.
- Epilachna jianchuanensis* Cao et Xiao, 1984 [CHINA: Yunnan]
- Epilachna lata* Li et Cook, 1961 [CHINA: Taiwan]
- Epilachna lenta* (Weise, 1902) [CHINA: Gansu; VIETNAM]
- Epilachna longissima* (Dieke, 1947) (*Afissa*) [CHINA: Fujian, Taiwan, Sichuan, Yunnan]
- Epilachna lugubris* (Dieke, 1947) (*Afissa*) [CHINA: Jiangsu]
- Epilachna macularis* Mulsant, 1850 [CHINA: Jiangxi, Fujian, Taiwan, Sichuan, Yunnan, Xizang; INDIA, NEPAL, BANGLADESH, VIETNAM, THAILAND, CELEBES, JAPAN] Host: *Solanum melongena*, *S. nigrum*, *S. tuberosum*, *S. suffruticosum*.
- Epilachna macularis donckieri* Weise, 1912 [CHINA: Fujian, Yunnan]
- Epilachna maculicollis* (Sicard, 1922) (*Solanophila*) [CHINA: Taiwan, Guangxi]
- Epilachna magna* (Dieke, 1947) (*Afissa*) [CHINA: Fujian, Sichuan, Yunnan] Host: *Solanum suffruticosum*.
- Epilachna marginicollis* (Hope, 1831) (*Coccinella*) [CHINA: Xizang; MYANMAR, SIKKIM, BANGLADESH, NEPAL, HIMALAYA]
- Epilachna maxima* (Weise, 1898) (*Solanophila*) [CHINA: Taiwan, Guangxi, Yunnan; INDIA, SIKKIM] Host: *Solanum suffruticosum*.
- Epilachna media* Li et Cook, 1961 [CHINA: Taiwan]
- Epilachna microgenitalis* Li et Cook, 1061 [CHINA: Taiwan, Hunan]
- Epilachna mirabiloides* (Dieke, 1947) (*Afissa*) [CHINA: Sichuan]
- Epilachna mobilitertiae* Li et Cook, 1961 [CHINA: Taiwan]
- Epilachna mushana* Li et Cook, 1961 [CHINA: Taiwan]
- Epilachna mystica* Mulsant, 1850 [CHINA: Xizang; INDIA, MYANMAR, NEPAL, SIKKIM]
- Epilachna mysticoides* (Sicard, 1912) (*Solanophila*) [CHINA: Xizang; INDIA, NEPAL]
- Epilachna narai* Lewis, 1896 [CHINA: NE CHINA; JAPAN, SIBERIA]
- Epilachna nielamuensis* Pang et Mao, 1977 [CHINA: Xizang; NEPAL]
- Epilachna nigrica* (Weise, 1908) (*Solanophila*) Syn. *Epilachna nigrica* var. *maculicollis* Sicard, 1912 [CHINA: Taiwan; INDIA]
- Epilachna ocellata* - *maculata* (Mader, 1930) (*Solanophila*) [CHINA: Hubei, Taiwan, Guizhou, Sichuan, Yunnan; MYANMAR] Host: *Pinus*, *Dendranthema morifolium*, *Artemisia ordosica*.
- Epilachna octomaculata* (Thunberg, 1781) (*Solanophila*) [CHINA: Taiwan; MYANMAR]
- Epilachna parainsignis* Pang et Mao, 1979 [CHINA: Guangxi, Yunnan]
- Epilachna paramagna* Pang et Mao, 1979 [CHINA: Hunan, Sichuan, Yunnan] Host: *Cinnamomum camphora*.
- Epilachna pingbianensis* Pang et Mao, 1979 [CHINA: Yunnan]
- Epilachna plicata* Weise, 1889 [CHINA: Shaanxi, Gansu, Hubei, Sichuan, Yunnan] Host: *Artemisia ordosica*.
- Epilachna provisorica* (Dieke, 1947) (*Afissa*) [CHINA: Sichuan]
- Epilachna pusillanum* Mulsant, 1850 [CHINA: Taiwan]
- Epilachna quadricollis* (Dieke, 1947) (*Afissa*) [CHINA: Hebei, Shandong, Jiangsu, Zhejiang, Fujian, Guangdong, Guangxi, Sichuan] Host: *Fraxinus*.
- Epilachna rubiacis* Cao et Xiao, 1984 [CHINA: Yunnan] Host: *Rubia cordifolia*.
- Epilachna saginata* (Weise, 1902) (*Solanophila*) [CHINA: Taiwan; VIETNAM]
- Epilachna sauteri* (Weise, 1923) (*Solanophila*) [CHINA: Jiangsu, Taiwan]
- Epilachna semenowi* Weise, 1887 [CHINA: Inner Mongolia]
- Epilachna siphodenticulata* Hoang, 1983 [CHINA: Guangxi; VIETNAM]
- Epilachna sociolamina* Li et Cook, 1961 [CHINA: Taiwan]
- Epilachna spiraloidea* Cao et Xiao, 1984 [CHINA: Yunnan]
- Epilachna subacuta* (Dieke, 1947) (*Afissa*) [CHINA: Hubei, Jiangsu, Zhejiang, Fujian, Guangxi, Sichuan] Host: *Fraxinus*, *Schisandra chinensis*.
- Epilachna szechuana* (Dieke, 1947) (*Afissa*) [CHINA: Sichuan] Host: *Schisandra chinensis*.
- Epilachna tianpingiensis* Pang et Mao, 1979 [CHINA: Henan, Hubei, Hunan, Guangxi] Host: *Vitis*.
- Epilachna tridecimmaculosa* Pang et Mao, 1977 [CHINA: Xizang]
- Epilachna undecempilota* (Hope, 1831) (*Coccinella*) [CHINA: Fujian, Taiwan, Hongkong, Xizang; INDIA, THAILAND, MYANMAR, NEPAL, SRI LANKA, INDONESIA]
- Epilachna yongshanensis* Cao et Xiao, 1984 [CHINA: Yunnan]
- Epiverta chelonia* (Mader, 1932) (*Solanophila*) [CHINA: Sichuan, Yunnan, Xizang] Host: *Artemisia tomentosa*, *Anemone tomentosa*, *A. rivularis*.
- Exochomus flavipes* (Thunberg, 1781) (*Coccinella*) [CHINA: Xinjiang; INDIA, PALAEARCTIC, AFRICA, AUSTRALIA, EUROPE] Prey on: aphids, scale insects.
- Exochomus mongol* Barovsky, 1922 [CHINA: Heilongjiang, Liaoning, Inner Mongolia, Hebei, Henan, Shaanxi, Gansu, Shandong, Anhui, Jiangsu, Zhejiang; MONGOLIA, KOREA] Prey on: aphids.
- Exochomus nigromarginatus* (Miyatake, 1970) (*Epilachna*) [CHINA: Gansu, Hubei, Jiangxi, Zhejiang, Fujian, Hunan, Sichuan, Yunnan] Prey on: aphids, scale insects.
- Halyzia maculata* Jing, 1987 [CHINA: Yunnan, Xizang]
- Halyzia sanscrita* Mulsant, 1853 [CHINA: Hebei, Henan, Shaanxi, Gansu, Hubei, Jiangsu, Jiangxi, Zhejiang, Fujian, Taiwan, Hunan, Guangxi, Guizhou, Sichuan, Yunnan, Xizang; INDIA, SIKKIM, BHUTAN, YEMEN] Host: *Podosphaera leucotricha*.
- Halyzia sedecimguttata* (Linnaeus, 1758) (*Coccinella*) [CHINA: Heilongjiang, Jilin, Hebei, Shaanxi, Xinjiang; JAPAN, MONGOLIA, SIBERIA, MINOR ASIA, EUROPE]
- Halyzia shirozui* Sasaji, 1982 [CHINA: Taiwan]
- Halyzia straminea* (Hope, 1831) (*Coccinella*) [CHINA: Gansu, Yunnan, Xizang; INDIA, NEPAL, SIKKIM, BANGLADESH] Prey on: fungi.
- Harmonia arcuata* (Fabricius, 1787) (*Coccinella*) [CHINA: INDIA, ANDAMAN I.]
- Harmonia axyridis* (Pallas, 1773) (*Coccinella*) Syn. *Coccinella bissex* - *notata* Herbst, 1793; *Harmonia axyridis* var. *aulica* Faldermann, 1835; *Harmonia axyridis* ab. *binaculata* Hemmelmann, 1926; *Harmonia axyridis* ab. *conspicua* Faldermann, 1835; *Harmonia axyridis* var. *novemdecimsignata* Faldermann, 1835; *Coccinella axyridis* var. *spectabilis* Faldermann, 1835; *Coccinella axyridis* ab. *succinea* Hope, 1845; *Ptychanais yedoensis* Takizawa, 1917 [CHINA: Heilongjiang, Jilin, Liaoning, Inner Mongolia, Hebei, Henan, Shanxi, Ningxia, Shaanxi, Gansu, Xinjiang, Shandong, Hubei, Jiangsu, Jiangxi, Zhejiang, Fujian, Guangdong, Hongkong, Hunan, Guangxi, Guizhou, Sichuan, Yunnan; SIKKIM, JAPAN, KOREA, SIBERIA, MONGOLIA] Prey on: *Adelges laricis* Vall, *Ambrostoma quadrimpressum* Motschulsky, *Aphis craccivora* Koch, *A. erysimi* (Kaltenbach), *A. glycines* Matsumura, *A. gossypii* Glover, *Cinara formosana* (Takahashi), *C. pinea* Mordvilko, *Hyaloptera amygdali* Kirkaldy, *Lipaphis erysimi* (Kaltenbach), *Longiumguis sacchari* (Zehntner), *Matsucoccus massonianae* Young et Hu, *M. mat-*

- sumurae Kuwana, *Microsiphum sanborni* Gillette, *Myzus numecola* Matsumura, *M. persicae* (Sulzer), *Macrosiphum avenae* Fabricius, *M. rosivorum* Zhang, *Oregma bambusicola* Takahashi, *Rhopalosiphum padi* Linnaeus, *R. nymphacae* (Linnaeus), *R. maidis* Fitch, *Schizaphis graminum* (Rodani), *S. piriicola* (Matsumura), *Toxoptera aurantii* Boyer et Fonscolombe, *T. citricida* (Kirkaldy), *Ceratovacuna lanigera* Zehntner, *Matsucoccus matsumurae* Kuwana, eggs and larvae of *Gastrolina depressa* Baly, *Heliothis armigera* (Hubner), *Podontia lutae* (Olivier), psyllids, mealybugs, Chermidae, rice leafhoppers, planthoppers, mites.
- Harmonia conformis* (Boisduval, 1835) (*Leis*) [CHINA]
- Harmonia dimidiata* (Fabricius, 1781) (*Coccinella*) Syn. *Coccinella quindecimmaculata* Hope, 1831; *Leis quindecimmaculata* Hope, 1837; *Harmonia dimidiata* ab. *humeralis* Weise, 1912; *Harmonia dimidiata* ab. *sicardi* Mader, ; *Synonycha kikuchii* Ohta, 1929 [CHINA: Henan, Jiangxi, Zhejiang, Fujian, Taiwan, Guangdong, Hunan, Guangxi, Guizhou, Sichuan, Yunnan, Xizang; INDIA, NEPAL, INDONESIA, VIETNAM, SIKKIM, BHUTAN, KASHMIR, JAPAN] Prey on: *Aphis craccivora* Koch, *Schizaphis graminum* (Rodani), aphids.
- Harmonia eucharis* (Mulsant, 1853) (*Ballia*) Syn. *Ballia eucharis* ab. *mayeti* Mulsant, 1866; *Ballia brahmae* Mulsant, 1853; *Ballia christophori* Mulsant, 1853; *Ballia diana* Mulsant, 1953; *Ballia gustavi* Mulsant, 1853; *Ballia montivaga* Mulsant, 1853; *Ballia testacea* Mulsant, 1853; *Ballia zephirinae* Mulsant, 1866; *Pelina mayeti* Mulsant, 1866; *Neda bayaderae* Mulsant, 1866; *Halyzia roubali* Mader, 1931; *Ballia korschefskeyi* Mader, 1931 [CHINA: Sichuan, Yunnan; INDIA, JAPAN] Prey on: *Aphis pomi* De Geer.
- Harmonia fallaciosa* (Mader, 1939) (*Leis*) [CHINA]
- Harmonia frigida* (Mulsant, 1853) (*Coccinella*) [CHINA; SIBERIA]
- Harmonia octomaculata* (Fabricius, 1781) (*Coccinella*) Syn. *Coccinella arcuata* Fabricius, 1787; *Synharmonia octomaculata*, Liu, 1963 [CHINA: Henan, Hubei, Jiangxi, Zhejiang, Taiwan, Guangdong, Hainan, Hunan, Guangxi, Guizhou, Sichuan, Yunnan; INDIA, INDONESIA, PHILIPPINES, JAPAN, MICRONESIA, AUSTRALIA, FIJI, SRI LANKA, NEW GUINEA, NEW CALEDONIA] Prey on: *Aphis craccivora* Koch, *Lipahis erysimi* (Kaltenbach), *Schizaphis graminum* (Rodani), *Roxoptera citricida* (Kirkaldy), rice leafhoppers and planthoppers.
- Harmonia rephirina* Mulsant, [CHINA: Guizhou] Prey on: aphids.
- Harmonia sedecimnotata* (Fabricius, 1801) (*Coccinella*) Syn. *Neda sedecimnotata* Fabricius, 1801; *Callineda sedecimnotata*, Crotch, 1874; *Daulis girini* Mulsant, 1866; *Callineda girini*, Mader, 1926 [CHINA: Jiangxi, Zhejiang, Fujian, Guangdong, Hongkong, Hainan, Hunan, Guangxi, Guizhou, Sichuan, Yunnan, Xizang; INDONESIA, NEPAL, PHILIPPINES, CALEMANDAN, JAPAN, YEMEN] Prey on: aphids, *Mytilaspis citricola* Packard, *Rhopalosiphum oryzae* (matsumura), *Lepidosaphes ume* Kuwana, *Unaspis yanonensis* Kuwana.
- Harmonia shoichui* Sasaji, 1988 [CHINA: Taiwan]
- Harmonia trilochana* (Kapur, 1963) (*Calvia*) Syn. *Chilocorus trilochana* Kapur, 1963; *Harmonia trilochana*, Cao et Xiao, 1986 [CHINA: Sichuan, Yunnan; SIKKIM] Host: aphids on Pinus and wheat.
- Harmonia yedoensis* (Takizawa, 1919) (*Ptychanatis*) Syn. *Ballia obscurosignata* Liu, 1963 [CHINA: Hebei, Henan, Shandong, Shaanxi, Hubei, Jiangsu, Jiangxi, Zhejiang, Fujian, Taiwan, Guangdong, Hunan, Guangxi, Guizhou, Sichuan, Yunnan; JAPAN, KOREA, VIETNAM] Prey on: *Cinara piniformosana* Takahashi, *Aphis gopssypii* Glover, *Matsucoccus matsumurae* Kuwana, rice planthoppers.
- Henosepilachna boisduvali* (Mulsant, 1850) (*Epilachna*) Syn. *Henosepilachna boisduvali boisduvali* Miyatake, 1971 [CHINA: Taiwan; JAPAN, INDIA, PHILIPPINES, INDONESIA, AUSTRALIA, JAPAN]
- Henosepilachna dodecastigma* Wiedemann, 1823) (*Epilachna*) [CHINA: Xizang; INDIA, MYANMAR, NEPAL, INDONESIA] Host: *Luffa cylindrica*.
- Henosepilachna indica* (Mulsant, 1850) (*Epilachna*) Syn. *Epilachna indica* var. *ceylonica* Weise, 1900 [CHINA: Taiwan, S. CHINA; INDIA, MYANMAR, VIETNAM, MALAYSIA, SRI LANKA, INDONESIA, PHILIPPINES,] Host: *Solanum melongena*, *Lycopersicon esculentum*, *Cucurbita moschata*, *Luffa cylindrica*.
- Henosepilachna kaszabi* (Bielawski et Fursch, 1960) (*Epilachna*) Syn. *Epilachna tonkinensis* Bielawski, 1957; *Epilachna tonkinensis* Weise, 1947 [CHINA: Guangxi, Guizhou, Yunnan; MYANMAR, VIETNAM, INDIA, THAILAND, PHILIPPINES]
- Henosepilachna libera* (Dieke, 1947) (*Epilachna*) [CHINA: Taiwan, Guangxi, Sichuan, Yunnan; INDIA, MYANMAR, MALAYSIA]
- Henosepilachna ocellata* (Redtenbacher, 1844) (*Epilachna*) [CHINA: Yunnan, Xizang; INDIA, NEPAL]
- Henosepilachna operculata* (Liu, 1950) (*Epilachna*) [CHINA: Beijing, Hubei, Jiangxi, Fujian] Host: *Actinostemma labatum*.
- Henosepilachna processa* (Weise, 1908) (*Epilachna*) Syn. *Epilachna wissmanni* ab. *processa* Weise, 1908; *Epilachna wissmanni* ab. *forsteri* Weise, 1908; *Epilachna processa* Weise, 1966 [CHINA: Taiwan, Yunnan; INDIA, MYANMAR, THAILAND, MALAYSIA]
- Henosepilachna pusillanima* (Mulsant, 1850) (*Epilachna*) Syn. *Epilachna dentulata* Dieke, 1947 [CHINA: Taiwan, Guangxi, Yunnan; INDIA, VIETNAM, THAILAND, NEPAL, PHILIPPINES] Host: *Solanum nigrum*.
- Henosepilachna pytharga* (Dieke, 1947) (*Epilachna*) [CHINA: Taiwan; PHILIPPINES]
- Henosepilachna quadriplagiata* Pang et Mao, 1977 [CHINA: Xizang]
- Henosepilachna semifasciata* (Dieke, 1947) (*Epilachna*) Syn. *Epilachna indica* ab. *ceylonica* Korschefskey, 1933 [CHINA: Taiwan]
- Henosepilachna septima* (Dieke, 1947) (*Epilachna*) [CHINA: Guangdong, Hainan, Guangxi; INDOCHINA, INDIA, ANDAMAN I., INDONESIA] Host: Cucurbitaceae, *Momordica charantia*.
- Henosepilachna sexta* (Dieke, 1947) (*Epilachna*) [CHINA: Yunnan; INDONESIA] Host: *Solanum nigrum*.
- Henosepilachna subfasciata* (Weise, 1947) (*Epilachna*) [CHINA: Taiwan]
- Henosepilachna tamdaoensis* Hoang, 1977 [CHINA: Guangxi, VIETNAM]
- Henosepilachna umbonata* Pang et Mao, 1979 [CHINA: Yunnan]
- Henosepilachna verriculata* Pang et Mao, 1979 [CHINA: Yunnan]
- Henosepilachna vigintioctomaculata* (Motschulsky, 1857) (*Epilachna*) Syn. *Epilachna niponica* Lewis, 1896; *Epilachna vigintioctomaculata coalescens* Mader, 1926 [CHINA: Heilongjiang, Jilin, Liaoning, Hebei, Shandong, Henan, Shanxi, Shaanxi, Gansu, Hubei, Jiangsu, Zhejiang, Fujian, Hunan, Guizhou, Sichuan, Yunnan, Xizang; INDIA, JAPAN, KOREA, SIBERIA] Host: *Solanum melongena*, *S. tuberosum*, *Lycopersicon esculentum*, *Datura stamonium*.
- Henosepilachna vigintioctopunctata* (Fabricius, 1775) (*Coccinella*) Syn. *Coccinella sparsa* Herbst, 1786; *Epilachna sparsa orientalis* Dieke, 1947; *Epilachna sparsa* var. *cinerea* Dieke, 1947 [CHINA: Heilongjiang, Jilin, Liaoning, Hebei, Shandong, Henan,

- Shaanxi, Hubei, Anhui, Jiangsu, Jiangxi, Zhejiang, Fujian, Taiwan, Guangdong, Hongkong, Hainan, Guangxi, Hunan, Guizhou, Sichuan, Yunnan, Xizang; INDIA, MYANMAR, THAILAND, BHUTAN, NEPAL, INDONESIA, JAPAN, AUSTRALIA, NEW GUINEA] Host: Citrus, Solanum melongena, S. tuberosum, Glycine max, Luffa cylindrica, melon.
- Henosepilachna wissmanni* (Mulsant, 1850) (*Epilachna*) [CHINA: Taiwan; INDIA, THAILAND, INDONESIA, PHILIPPINES] Hikonanikona monticola Sasaji, 1967 [CHINA: Taiwan]
- Hippodamia arctica* (Schneider, 1792) (*Coccinella*) Syn. *Adonia amoena* (Faldermann, 1835) [CHINA: Inner Mongolia; SIBERIA, MONGOLIA, KAZAKHSTAN]
- Hippodamia grundmanni* (Mader, 1931) (*Adonia*) [CHINA]
- Hippodamia potanini* (Weise, 1889) (*Semiadalia*) Syn. *Asemiadalia obenbergeri* Mader, 1926; *Semiadalia obenbergeri* Barowsky, 1928; *Semiadalia obenbergeri ancora* Mader, 1933; *Semiadalia obenbergeri antecununcta* Mader, 1933; *Semiadalia obenbergeri comuna* Mader, 1933; *Semiadalia obenbergeri hauseri* Mader, 1933; *Semiadalia obenbergeri manca* Mader, 1933; *Semiadalia obenbergeri paupera* Mader, 1933; *Semiadalia obenbergeri priesneri* Mader, 1933; *Semiadalia potanini bella* Mader, 1933; *Semiadalia potanini completa* Mader, 1933 [Shaanxi, Gansu, Hubei, Jiangsu, Sichuan, Yunnan, Xizang] Host: aphids.
- Hippodamia tredecimpunctata* (Linnaeus, 1758) (*Coccinella*) Syn. *Coccinella tibialis* Say, 1924; *Coccinella signata* Faldermann, 1832; *Coccinella timberlakei* Capra, 1931 [CHINA: Heilongjiang, Jilin, Liaoning, Hebei, Henan, Shaanxi, Ningxia, Gansu, Qinghai, Xinjiang, Shandong, Anhui, Hubei, Jiangsu, Jiangxi, Zhejiang; JAPAN, KOREA, SIBERIA, MONGOLIA, ALASKA, CAUCASUS, IRAN, KAZAKHSTAN, AFGHANISTAN, EUROPE, ASIA] Prey on: Aphis gossypii Glover, A. robiniae Macchiatii, Macrosiphum avenae (Fabricius), Schizaphis graminum (Rondani), rice planthoppers.
- Hippodamia ulkei* (Crotch, 1873) (*Ceratomegilla*) Syn. *Spiladelphia barowskii* Semenov et Dobzhanskyi, 1923; *Ceratomegilla parva* Watson, 1954 [CHINA: Xizang; MONGOLIA, ALASKA, CANADA, KAZAKHSTAN]
- Hippodamia undecimnotata* (Schneider, 1792) (*Semiadalia*) Syn. *Coccinella undecimnotata* Mader, 1926 [CHINA: Yunnan; C. ASIA, EUROPE] Host: Aphis.
- Hippodamia variagata* (Goeze, 1777) (*Coccinella*) Syn. *Adonia variagata*, Rybakow, 1889; *Adonia doubledayi* Mulsant, 1850; *Adonia tredecimsignata* Mulsant, 1850; *Adonia variagata ab. thoracica* Weise, 1889 [CHINA: Heilongjiang, Jilin, Liaoning, Inner Mongolia, Hebei, Shanxi, Ningxia, Gansu, Shaanxi, Qinghai, Xinjiang, Henan, Shandong, Hubei, Jiangsu, Jiangxi, Fujian, Taiwan, Hunan, Sichuan, Yunnan, Xizang; JAPAN, INDIA, NEPAL, BHUTAN, AFGHANISTAN, C. AFRICA, ENGLAND, PALAEARCTIC] Prey on: Aphis gossypii Glover, A. craccivora Koch, A. robiniae Macchiatii, A. medicaginis Koch, A. pomi De Ceer, Aphanostigma iakusuiensis Kishida, Myzus momonis Matsumura, M. malisuctus Matsumura, Rhopalosiphum maidis Fitch, Sappaphis picicola Okamoto et Takahashi, Schizaphis graminum (Rondani), eggs of Heliothis armigera (Hubner).
- Hippodamia weisei* (Fivaldsky, 1892) (*Adonia*) [CHINA: N. CHINA]
- Hormiolus fortunatus* (Lewis, 1896) (*Scymnus*) [CHINA: Shaanxi; JAPAN]
- Hormiolus hisamatsui* Miyatake, 1976 [CHINA: Hongkong]
- Hormiolus novempunctatus* Miyatake, 1979 [CHINA: Taiwan]
- Hyperaspis amurensis* Weise, 1887 [CHINA: Henan, Zhejiang; RUSSIA]
- Hyperaspis asiatica* Lewis, 1896 [CHINA: Heilongjiang, Jilin, Liaoning, Hebei, Shandong, Shaanxi, Jiangsu, Zhejiang; JAPAN]
- Prey on: aphids, scale insects.
- Hyperaspis gyotokui* Kamiya, 1963 [CHINA: Hebei, Henan, Shaanxi, Sichuan; JAPAN] Prey on: aphids.
- Hyperaspis japonica* (Crotch, 1874) [CHINA: Jiangsu, Zhejiang; JAPAN, SIBERIA, AMUR]
- Hyperaspis leechi* Miyatake, 1961 [CHINA: Heilongjiang, Jilin, Liaoning, Hebei, Henan, Shaanxi, Hubei, Anhui, Jiangsu, Zhejiang, Fujian, Sichuan; JAPAN, KOREA, E. SIBERIA, MONGOLIA] Prey on: Aphis gossypii Glover.
- Hyperaspis repensis* (Herbst, 1873) [CHINA: Hebei, Shandong, Hubei, Jiangsu, Sichuan; PALAEARCTIC, ABYSSINIA] Prey on: Aphis gossypii Glover.
- Hyperaspis sinensis* (Crotch, 1874) (*Cryptogonus*) [CHINA: Jilin, Shaanxi, Gansu, Hebei, Henan, Hubei, Anhui, Jiangsu, Jiangxi, Zhejiang, Fujian, Guangdong, Guangxi, Guizhou, Sichuan; JAPAN, SIBERIA] Prey on: psyllids, pseudococcids, aleyrodids, leaf beetles.
- Illeis bistigmosa* (Mulsant, 1850) (*Egleis*) Syn. *Illis simplex* Mulsant, 1866 [CHINA: Guangdong, Guangxi, Yunnan; THAILAND, INDIA, VIETNAM, SRI LANKA, MALAYSIA, INDONESIA, PHILIPPINES] Host: fungi.
- Illeis chinensis* Yablokov - Kunzoryan, 1978 [CHINA: Hongkong, Guangxi] Host: fungi.
- Illeis cincta* (Fabricius, 1798) (*Coccinella*) Syn. *Thea cincta*, Crotch, 1874 [CHINA: Hebei, Henan, Hubei, Jiangsu, Jiangxi, Fujian, Taiwan, Guangdong, Hongkong, Hainan, Hunan, Guizhou, Sichuan, Yunnan; INDIA, INDONESIA, PHILIPPINES, SRI LANKA, JAPAN, KOREA] Prey on: fungi.
- Illeis confusa* Timberlake, 1943 [CHINA: Zhejiang, Guangdong, Hongkong, Yunnan; VIETNAM, AMERICA]
- Illeis indica* Timberlake, 1960 [CHINA: Yunnan; INDIA, PAKISTAN, HAWAII] Host: fungi.
- Illeis koebelei* Timberlake, 1943 Syn. *Illeis koebelei amamiana* Miyatake, 1959; *Illeis koebelei koebelei* Timberlake, 1943; *Thea cincta* Weise, 1923 [CHINA: Hebei, Shaanxi, Shaanxi, Henan, Hubei, Jiangsu, Fujian, Taiwan, Guangdong, Hunan, Guangxi, Sichuan, Yunnan; JAPAN, KOREA, PHILIPPINES, INDONESIA, HAWAII] Prey on: fungi.
- Illeis shensiensis* Timberlake, 1943 [CHINA: Hebei, Shaanxi, Fujian, Sichuan, Yunnan] Host: fungi
- Jauravia assamensis* Kapur, 1946 [CHINA: Yunnan; INDIA]
- Jauravia limbata* Motschulsky, 1858 Syn. *Paraclitostethus ovatus* Ohta, 1929; *Paraclitostethus ovatus ab. kokoronis* Ohta, 1929 [CHINA: Taiwan, Yunnan; INDIA, THAILAND, NEPAL, VIETNAM, SRI LANKA, JAPAN]
- Jauravia quadrinotata* Kapur, 1946 [CHINA: Taiwan, Yunnan; INDIA, BHUTAN, NEPAL]
- Keiscymnus koebelei* Pang et Gordon, 1986 [CHINA: Hainan]
- Keiscymnus taiwanensis* Yang, 1972 [CHINA: Fujian, Taiwan, Hainan, Guangxi, Guizhou]
- Lemnia biplagiata* (Swartz, 1808) (*Coccinella*) Syn. *Lemnia fraudulentula* Mulsant, 1850; *Lemnia melanota* Mulsant, 1850; *Coleophora nepalensis* Crotch, 1874; *Coleophora personata* Weise, 1910; *Osumia bimaculata* Kurisaki, 1923 [CHINA: Henan, Anhui, Jiangsu, Jiangxi, Zhejiang, Fujian, Taiwan, Guangdong, Hongkong, Guangxi, Sichuan, Yunnan, Xizang; JAPAN, KOREA, PHILIPPINES, INDONESIA, INDIA, MYANMAR, THAILAND, VIETNAM, NEPAL, MALAYSIA, INDOCHINA] Prey on: Schizaphis graminum (Rondani), Toxoptera citricida (Kirkaldy), Myzus persicae (Sulzer), Brevicorne brassicae (Linnaeus), Lipaphis erysimi (Kaltenbach), Aphis bambusae Fullaway, Ceratovacuna lanigera Zehntner.
- Lemnia biquadriguttata* Jing, 1988 [CHINA: Yunnan]
- Lemnia bissellata* (Mulsant, 1850) (*Coleophora*) [CHINA: Hubei,

- Jiangxi, Fujian, Taiwan, Guangdong, Hainan, Guangxi, Guizhou, Sichuan, Yunnan, Xizang; KOREA, PHILIPPINES, INDONESIA, S. ASIA, SE ASIA] Prey on: *Aphis craccivora* Koch, rice plathoppers, *Schizaphis graminum* (Rondani).
- Lernia brunniplagiata* Jing, 1988 [CHINA: Yunnan]
- Lernia callinotata* Jing, 1988 [CHINA: Yunnan]
- Lernia circumsta* (Mulsant, 1850) (*Artemis*) Syn. *Artemis mandarina* Mulsant, 1850; *Artemis rufila* Mulsant, 1850; *Coelophora westermanni* Mulsant, 1850; *Coleophora circumsta* Mulsant, 1850 [CHINA: Fujian, Taiwan, Guangdong, Hongkong, Hainan, Guangxi, Yunnan; INDIA, VIETNAM, HAWAII] Prey on: aphids.
- Lernia circumvelata* (Mulsant, 1874) (*Coleophora*) [CHINA: Shaanxi, Hubei, Anhui, Zhejiang, Fujian, Taiwan, Hainan, Guangdong, Hongkong, Hunan, Guangxi, Guizhou, Sichuan, Yunnan; INDIA] Prey on: *Aphis craccivora* Koch.
- Lernia duvauceli* (Mulsant, 1850) (*Caria*) Syn. *Alloneda novemmaculata* Cao et Pu, 1985 [CHINA: Hongkong, Guangxi; JAPAN, MYANMAR, INDIA, VIETNAM, INDONESIA, JAPAN]
- Lernia henricae* Mulsant, 1866 [CHINA]
- Lernia jianfengensis* Jing, 1988 [CHINA: Hainan]
- Lernia loi* Sasaji, 1982 [CHINA: Taiwan]
- Lernia melanaria* (Mulsant, 1850) (*Synia*) Syn. *Leis rougei* Mulsant, 1866; *Synia rougei* Mulsant, 1866 [CHINA: Shaanxi, Gansu, Henan, Hubei, Jaingxi, Fujian, Taiwan, Guangdong, Guangxi, Guizhou, Sichuan, Hunan, Yunnan; PHILIPPINES, INDIA, SRI LANKA, VIETNAM]
- Lernia newporti* (Mulsant, 1850) (*Coleophora*) Syn. *Coleophora bouringi* Crotch, 1874 [CHINA; PHILIPPINES, INDONESIA, MYANMAR]
- Lernia octopunctata* (Weise, 1910) [CHINA: Taiwan; PHILIPPINES, INDONESIA]
- Lernia saucia* (Mulsant, 1856) (*Coleophora*) Syn. *Coleophora calypso* Mulsant, 1856; *Coleophora mouhoti* Crotch, 1874; *Coleophora swinhoei* Crotch, 1874; *Chilomenes takanoni* Ohta, 1929; *Coleophora ishidai* Ohta, 1931 [CHINA: Henan, Shandong, Shaanxi, Gansu, Jiangsu, Jiangxi, Zhejiang, Fujian, Taiwan, Guangdong, Hongkong, Hunan, Guangxi, Guizhou, Sichuan, Yunnan; INDIA, VIETNAM, THAILAND, NEPAL, PHILIPPINES, INDONESIA, JAPAN, HAWAII] Prey on: aphids.
- Lernia sexguttata* (Weise, 1910) (*Coleophora*) [CHINA: Taiwan; PHILIPPINES]
- Lernia vidua* (Mulsant, 1850) (*Coleophora*) [CHINA: Taiwan; INDIA, INDONESIA, TIMOR, PHILIPPINES,]
- Lithophilus kozlovi* (Barovsky, 1909) (*Tetrabrachys*) [CHINA: Inner Mongolia, Beijing, Shanxi; MONGOLIA] Prey on: eggs of *Tuberaleyrodes machili* Takahashi, *Aleuroplatus pectiniferus* Quaintance and *Icerya purchasi* Marshall.
- Lithophilus subseriatus* (Reitter, 1883) (*Tetrabrachys*) Syn. *Tetrabrachys subseriatus* var. *unicolor* Reitter, 1883 [CHINA; MONGOLIA, EGYPT, GREECE]
- Lithophilus villosus* Faldermann, 1835 [CHINA: N. CHINA; MONGOLIA]
- Macrouleis hauseri* (Mader, 1926) (*Halysia*) [CHINA: Hubei, Shaanxi, Gansu, Fujian, Taiwan, Hainan, Hunan, Guangxi, Guizhou, Sichuan, Yunnan, Xizang; BHUTAN] Prey on: powdery mildew, fungi.
- Macronaemia hauseri* (Weise, 1905) (*Anisocalnia*) Syn. *Naemia hauseri* Weise, 1853 [CHINA: Henan, Hubei, Shaanxi, Hainan, Hunan, Guizhou, Sichuan, Yunnan, Xizang] Prey on: *Aphis craccivora* Koch, *A. nerii* Boyer et Fonscolombe, Toxoptera, *Schizaphis graminum* (Rondani), powdery mildew.
- Macronaemia hauseri* communicata Mader, 1929? [CHINA: Hubei, Sichuan, Yunnan] Prey on: fungi.
- Macronaemia paradoxa* (Mader, 1947) (*Coleomegilla*) [CHINA: Yunnan] Prey on: scale insects.
- Macronaemia yunnanensis* Cao et Xiao, 1985 [CHINA: Yunnan]
- Megalocaria dilatata* (Fabricius, 1775) (*Coccinella*) Syn. *Anisolemia dilatata* Fabricius, 1775 [CHINA: NE CHINA, Hubei, Jiangxi, Fujian, Taiwan, Guangdong, Hongkong, Guangxi, Guizhou, Sichuan, Yunnan; INDIA, INDONESIA, VIETNAM, CALEMANDAN] Prey on: *Ceratovacuna lanigera* Zehntner, *Aphis bambusae* Rullaway, *Hyalopterus arundinis* Fabricius, *Toxoptera odinae* Van der Goot.
- Megalocaria major* (Sicard, 1929) (*Anisolemia*) Syn. *Anisolemia major* ab. *yunnana* Sicard, 1929 [CHINA: Yunnan; CALEMANDAN]
- Megalocaria reichii pearsoni* (Crotch, 1874) (*Anisolemia*) Syn. *Neda reichei* Mulsant, 1950 [CHINA: Heilongjiang, Jilin, Inner Mongolia, Hebei, Shanxi, Anhui, Zhejiang, Yunnan, Xizang; INDIA, SIKKIM, INDONESIA, AUSTRALIA] Prey on: *Adelge* on *Pinus armandii*.
- Menochilus sexmaculata* (Fabricius, 1781) (*Coccinella*) Syn. *Coccinella quadriplagiata* Swartz, 1808; *Chilomenes quadriplagiata* Swartz, 1808; *Chilomenes quadriplagiata*, Lewis, 1873 [CHINA: Heilongjiang, Jilin, Liaoning, Shaanxi, Gansu, Shandong, Henan, Hubei, Jiangsu, Jiangxi, Zhejiang, Fujian, Taiwan, Guangdong, Hunan, Hainan, Guangxi, Guizhou, Sichuan, Yunnan; INDIA, CAMBODIA, THAILAND, MALAYSIA, SRI LANKA, PHILIPPINES, INDONESIA, AFGHANISTAN, IRAN, MICRONESIA, NEW GUINEA, JAPAN, CELEBES] Prey on: *Aphis craccivora* Koch, *A. gossypii* Glover, *A. medicaginis* Koch, *Diaphorina citri* (Kuwayama), *Geisha distinctissima* Walker, *Hyaloptera arundinis* Fabricius, *Lipaphis erysimi* (Kaltenbach), *Myzus persicae* (Sulzer), *Schizaphis graminum* (Rondani), *Toxoptera citricida* (Kirkaldy), rice leafhoppers, rice plathoppers, *Rhopalosiphum maidis* (Fitch).
- Micraspis allardi* (Mulsant, 1866) (*Lernia*) Syn. *Verania malaccensis* Crotch, 1874 [CHINA: Fujian, Guangdong, Yunnan; INDIA, CAMBODIA, MALAYSIA, INDONESIA, PHILIPPINES, AFGHANISTAN]
- Micraspis chinensis* (Mader, 1954) [CHINA]
- Micraspis discolor* (Fabricius, 1798) (*Coccinella*) Syn. *Alosia discolor* Fabricius, 1798; *Micraspis kurasoii* Miyatake, 1977; *Micraspis satoi* Miyatake, 1977 [CHINA: Henan, Shaanxi, Shandong, Hubei, Jiangsu, Jiangxi, Zhejiang, Fujian, Taiwan, Guangdong, Hongkong, Hainan, Hunan, Guangxi, Guizhou, Sichuan, Yunnan, Xizang; INDIA, THAILAND, SRI LANKA, INDONESIA, MALAYSIA, PHILIPPINES, CALEMANDAN, MICRONESIA, JAPAN] Prey on: *Aphis gossypii* Glover, *Lipaphis erysimi* (Kaltenbach), *Macrosiphum avenae* Fabricius, *Rhopalosiphum oryzae* (Matsumura), *Schizaphis graminum* (Rondani), *Thrips oryzae* Williams, *Toxoptera aurantii* Boyer et Fonscolombe, *Agromyza oryzae* (Munakata), *Agromyza oryzae* (Munakata), rise leafhopper, rice plathopper, pollen and stamen of rise, eggs and larvae of *Lepidoptera* on rice, *Oryza sativa*, *Zea mays*, *Brassica campestris*.
- Micraspis inops* (Mulsant, 1866) (*Chilomenes*) Syn. *Vernaria victa* Gorham, 1895; *Vernaria kiotoensis* Nakena et Araki, 1960 [CHINA: Guangdong, Yunnan; INDIA, VIETNAM, THAILAND, MYANMAR, INDONESIA, JAPAN] Prey on: aphids, pollen.
- Micraspis trilinea* Weise, 1889 [CHINA: Gansu]
- Micraspis univittata* (Hope, 1837) (*Coccinella*) [CHINA: Xizang; INDIA, NEPAL]
- Micraspis yunnanensis* Jing, 1985 [CHINA: Yunnan] Prey on: aphids.
- Microserangium okinawense* Miyatake, 1961 [CHINA: Taiwan; JAPAN]

- Nedina mirabilis* Hoang, 1984 [CHINA: Yunnan; VIETNAM] Prey on: *Aphis bambusae* Fullaway.
- Nephus bipunctatus* (Kugelann, 1794) (*Scymnus*) [CHINA: Hongkong]
- Nephus dilepismoides* Pang et Pu, 1988 Syn. *Nephus ancyrroides* Pang et Pu, 1988 [CHINA: Fujian, Taiwan, Guangdong, Hunan, Guangxi, Sichuan]
- Nephus fijiensis* (Sicard, 1922) [CHINA: Guangdong; FIJI]
- Nephus klapperichi* (Mader, 1955) [CHINA: Fujian, Guangxi]
- Nephus koltzei* (Weise, 1887) [CHINA: Hebei, Henan, Gansu, Guangxi, Guizhou, Sichuan; SIBERIA] Prey on: mealybugs on *Salix*, *Eriosoma* on *Salix*.
- Nephus koreanus* (Fursch, 1981) [CHINA: Fujian; KOREA]
- Nephus lancetapicalis* Pang et Gordon, 1986 [CHINA: Guangdong]
- Nephus obsoletus* (Weise, 1890) (*Scymnus*) [CHINA: Xizang]
- Nephus parenthesis* Weise, 1923 [CHINA: Taiwan]
- Nephus patagiatus* (Lewis, 1896) (*Scymnus*) [CHINA: Shaanxi, Taiwan; JAPAN, VIETNAM]
- Nephus phosphorus* (Lewis, 1896) (*Scymnus*) [CHINA: Taiwan, Sichuan; JAPAN, FRANCE, ITALY, GREECE, AUSTRIA]
- Nephus quadrimaculatus* (Herbst, 1783) (*Sphaeridium*) [CHINA: Taiwan; TURKEY, AUSTRIA, FRANCE, ITALY, GREECE, SYRIA]
- Nephus roepkei* (Fluiter, 1938) (*Scymnus*) [CHINA: Guangdong, Hainan; JAVA, PHILIPPINES, MICRONESIA, JAPAN]
- Nephus ryuguus* (Kamiya, 1961) (*Scymnus*) [CHINA: Shaanxi, Jiangsu, Fujian, Guangdong, Hainan, Guangxi, Guizhou, Sichuan; JAPAN] Prey on: mealybugs, *Unaspis yanoensis* (Kuwana).
- Nephus ryukyuensis* Sasaji, 1971 [CHINA: Mainland of CHINA, Taiwan; JAPAN]
- Nephus sauteri* Ohta, 1929 [CHINA: Taiwan]
- Nephus tagiapatus* (Kamiya, 1965) (*Scymnus*) [CHINA: Taiwan, Guangdong, Guangxi; JAPAN, MALAYSIA, THAILAND, INDIA]
- Nesolotis shirozui* Sasaji, 1967 [CHINA: Taiwan]
- Nesolotis tsunekii* Sasaji, 1967 [CHINA: Taiwan]
- Novius hauseri* Mader, 1930? [CHINA: Sichuan, Yunnan]
- Oenopia billieti* (Mulsant, 1853) (*Harmonia*) Syn. *Synharmonia indica* Weise, 1903 [CHINA: Sichuan, Xizang; INDIA, KASHMIR, BANGLADESH]
- Oenopia bissexnotata* (Mulsant, 1850) (*Leis*) Syn. *Synharmonia bissexnotata* (Mulsant,) [CHINA: Heilongjiang, Jilin, Liaoning, Hebei, Shaanxi, Qinghai, Gansu, Xinjiang, Hubei, Shandong, Henan, Jiangxi, Zhejiang, Hunan, Guizhou, Sichuan, Yunnan; Former USSR] Prey on: *Aphis gossypii* Glover, *A. medicaginis* Koch, *A. pomi* De Geer, *Rhopalosiphum maidis* Fitch, *Toxoptera citricada* (Kirkaldy), *Eriosoma lanigerum* (Hausman), *Rhopalosiphum oryzae* (Matsumura).
- Oenopia bluthgeni* Mader, 1933 [CHINA: Yunnan]
- Oenopia chinensis* (Weise, 1912) (*Coleophora*) Syn. *Coleophora fallaciosa* Mader, 1955; *Gyrocaria formosana* Miyatake, 1965 [CHINA: Shaanxi, Gansu, Shandong, Jiangsu, Jiangxi, Zhejiang, Fujian, Taiwan, Guangdong, Hunan, Guangxi, Guizhou, Sichuan, Yunnan] Prey on: aphids on *Pinus*.
- Oenopia conglobata* (Linnaeus, 1758) (*Coccinella*) Syn. *Coccinella conglobata contaminata* Menetries, 1849; *Coccinella conglobata buphthalmus* Mulsant, 1850 [CHINA: Heilongjiang, Inner Mongolia, Hebei, Shanxi, Shaanxi, Ningxia, Qinghai, Gansu, Xinjiang, Henan, Shandong, Anhui, Jiangsu, Zhejiang, Fujian, Sichuan, Xizang; MONGOLIA, RUSSIA, AFGHANISTAN, EUROPE, SIBERIA, PALAEARCTIC] Prey on: *Aphis gossypii* Glover, *A. saliceti* Kaltentbach, *Eriosoma lanigerum* (Hausmann), *Lipaphis erysimi* (Kaltentbach), *Longiumguis sacchari* (Zehntner), *Macrosiphum avenae* Fabricius, *Myzus persicae* (Sulzer), *Rhopalosiphum maidis* Fitch, *Rh. oryzae* (Matsumura), *Schizaphis graminum* (Rondani).
- Oenopia conglobata desertotum* (Rybakov, 1889) [CHINA: Inner Mongolia; TURKEY]
- Oenopia draconguttata* Jing, 1986 [CHINA: Xizang]
- Oenopia emmerichi* Mader, 1926 [CHINA: Hubei, Jiangxi, Sichuan, Yunnan, Xizang] Prey on: aphids.
- Oenopia flavidbrunna* Jing, 1986 [CHINA: Yunnan]
- Oenopia formosana* (Miyatake, 1965) (*Gyrocaria*) [CHINA: Shaanxi, Fujian, Taiwan, Hunan] Host: aphids.
- Oenopia kirbyi* Mulsant, 1850 [CHINA: Hubei, Taiwan, Sichuan, Yunnan, Xizang; INDIA, MYANMAR, THAILAND, HAWAII] Prey on: aphids.
- Oenopia octomaculata* (Fabricius, 1781)? (*Synharmonia*) [CHINA: Sichuan] Prey on: *Schizaphis graminum* (Rondani).
- Oenopia oncina* (Olivier, 1808) (*Coccinella*) Syn. *Coccinella oncina addicta* Mulsant, 1850 [CHINA: Hubei; IRAQ, IRAN, AFGHANISTAN, MONGOLIA, CAUCASUS, KAZAKHSTAN, EGYPT]
- Oenopia pomiensis* Jing, 1987 [CHINA: Xizang]
- Oenopia quadripunctata* Kapur, 1963 [CHINA: Xinjiang, Hubei, Jiangsu, Yunnan, Xizang; INDIA, SIKKIM, MYANMAR] Prey on: aphids.
- Oenopia sauzeti* Mulsant, 1866 [CHINA: Hubei, Gansu, Shaanxi, Anhui, Zhejiang, Fujian, Taiwan, Guangdong, Hunan, Guangxi, Guizhou, Sichuan, Yunnan, Xizang; INDIA, MYANMAR, THAILAND, BHUTAN, SIKKIM, NEPALHIMALAYA, VIETNAM] Prey on: *Aphis farinosa* Gmelin, *A. gossypii* Glover, *Macrosiphum avenae* Fabricius, *Myzus persicae* (Sulzer), *Schizaphis graminum* (Rondani), *Adelges*.
- Oenopia scalaris* (Timberlake, 1943) (*Protocaria*) [CHINA: Hebei, Fujian, Taiwan, Guangdong, Sichuan; INDOCHINA, JAPAN, KOREA, HAWAII, MICRONESIA] Prey on: aphids.
- Oenopia sexareata* (Mulsant, 1853) (*Coleophora*) [CHINA: Guangxi, Sichuan, Yunnan, Xizang; INDIA, MYANMAR, VIETNAM, INDONESIA, HIMALAYA, JAPAN] Prey on: aphids.
- Oenopia sexmaculata* Jing, 1986 [CHINA: Sichuan]
- Oenopia signatella* (Mulsant, 1866) [CHINA: Guangxi, Yunnan, Xizang; MYANMAR, INDIA, SRI LANKA]
- Oenopia takasago* (Sasaji, 1982) (*Chelonitis*) [CHINA: Taiwan]
- Ortalia bruneiana menglunensis* Pang et Mao, 1979 [CHINA: Guangxi, Yunnan; MALAYSIA] Prey on: *Aphis bambusae* Fullaway, *Aleurocanthus* on *Bambusa*.
- Ortalia jinghongensis* Pang et Mao, 1979 [CHINA: Yunnan] Prey on: aphids.
- Ortalia nigropectoralis* Pang et Mao, 1979 [CHINA: Yunnan]
- Ortalia pectoralis* Weise, 1901 [CHINA: Zhejiang, Fujian, Guangxi, Yunnan, Xizang; INDIA] Prey on: aphids.
- Ortalia yunnanensis* Pang et Mao, 1979 [CHINA: Guangdong, Hainan, Sichuan, Yunnan] Prey on: *Toxoptera citricida* (Kirkaldy), *Aphis craccivora* Koch.
- Oxynychua laexandras* Weise, 1890 [CHINA: Inner Mongolia]
- Palaeneda miniata* (Hope, 1831) (*Coccinella*) Syn. *Neda auriculata* Mulsant, 1866 [CHINA: Yunnan; INDIA]
- Pania insularis* (Sicard, 1912) (*Coleophora*) Syn. *Coleophora insularis* Weise, 1923; *Coleophora insularis* Miwa, 1931 [CHINA: Taiwan]
- Pania luteopustulata* (Mulsant, 1866) (*Oenopia*) Syn. *Oenopia praecae* Weise, 1891; *Coleophora korschefskyi* Mader, 1930 [CHINA: Henan, Shaanxi, Hubei, Anhui, Jiangxi, Fujian, Taiwan, Guangdong, Guangxi, Guizhou, Sichuan, Yunnan, Xizang; INDIA, MYANMAR, BHUTAN, THAILAND, NEPAL] Prey on: *Schizaphis graminum* (Rondani), aphids.
- Pania luteopustulata thibetana* Mulsant, 1853 [CHINA: Xizang]
- Pania victoriae* (Mulsant, 1866) (*Coccinella*) Syn. *Calvia thoraci-*

- ca Weise, 1902; *Coleophora insularis* Sicard, 1912 [CHINA: Taiwan]
- Parippodamia arctica (Schneider, 1792) [CHINA: Inner Mongolia, Hebei; N. EUROPE] Prey on: aphids.
- Phaenochilus metasternalis Miyatkae, 1970 [CHINA: Guangdong, Hainan, Hunan, Guangxi, Yunnan, Xizang] Prey on: aphids.
- Pharoscyrmus brunneosignatus Mader, 1949 [CHINA: Xinjiang; MONGOLIA]
- Pharoscyrmus taoi Sasaji, 1967 [CHINA: Fujian, Taiwan, Guangdong, Guangxi] Prey on: Aphis craccivora Koch, Rhopalosiphum pseudobrassicae (Davis).
- Phrynocaria approximans (Crotch, 1874) (*Coleophora*) Syn. *Chilomenes teretus* Ohta, 1929 [CHINA: Fujian, Taiwan, Guangdong, Hainan] Prey on: Toxoptera citricida (Kirkaldy).
- Phrynocaria congener (Billberg, 1808) (*Coccinella*) [CHINA: Jiangxi, Zhejiang, Fujian, Taiwan, Guangdong, Hongkong, Hainan, Guangxi, Sichuan, Yunnan; INDIA, JAPAN] Prey on: aphids, Macrosiphum avenae Fabricius, ruce planthoppers, .
- Phrynocaria nigrlimbata Jing, 1986 [CHINA: Yunnan] Prey on: aphids.
- Phrynocaria shirozui (Sasaji, 1982) (*Propylea*) [CHINA: Taiwan]
- Phymatosternus lewisii (Crotch, 1874) (*Platynaspis*) [CHINA: Jilin, Henan, Shaanxi, Hubei, Jiangsu, Jiangxi, Zhejiang, Fujian, Taiwan, Guangdong, Hainan, Guangxi, Guizhou, Yunnan; INDIA, MYANMAR, JAPAN, KOREA] Prey on: scale insects.
- Platynaspis angulimaculata Mader, 1938 [CHINA: Guangdong, Sichuan, Yunnan, Xizang; THAILAND] Prey on: Nippolachnus piri Matsumura, Pterochlorus tropicalis Van der Goot, aphids, scale insects.
- Platynaspis babai (Sasaji, 1988) (*Platynaspisius*) [CHINA: Taiwan]
- Platynaspis bimaculata Pang et Mao, 1979 [CHINA: Yunnan] Prey on: aphids.
- Platynaspis gressitti (Miyatake, 1961) (*Phymatosternus*) [CHINA: Fujian, Guangdong, Hainan, Yunnan] Prey on: aphids.
- Platynaspis hainanensis (Miyatake, 1961) (*Phymatosternus*) [CHINA: Hainan, Guangxi] Prey on: Cerataphis.
- Platynaspis hunagea Cao et Xiao, 1984 [CHINA: Yunnan] Prey on: aphids on Engelhardtia roxburghiana.
- Platynaspis maculosa Weise, 1910 [CHINA: Shaanxi, Gansu, Hubei, Jiangsu, Jiangxi, Zhejiang, Fujian, Guangdong, Hainan, Guangxi, Guizhou, Sichuan] Prey on: Aphis craccivora Koch, Longium sacchari (Zehntner), aphids, scale insects.
- Platynaspis ocellimaculata Pang et Mao, 1979 [CHINA: Yunnan] Prey on: aphids on Prunus persica, Cinnamomum camphora, Ficus.
- Platynaspis octoguttata (Miyatake, 1961) (*Phymatostethus*) [CHINA: Fujian, Guangdong, Yunnan] Prey on: Aphis craccivora Koch, Toxoptera aurantii Boyer et Fonscolombe.
- Platynaspis quinquepunctatus (Miyatake, 1961) (*Platynaspidius*) [CHINA: Taiwan]
- Platynaspis sexmaculata Cao et Xiao, 1984 [CHINA: Yunnan] Prey on: aphids on Ficus.
- Platynaspis tricolor (Hoang, 1983) [CHINA: Guangdong; VIETNAM]
- Plotina mulleri Mader, 1955 [CHINA: Fujian]
- Plotina versicolor Lewis, 1896 [CHINA: Fujian; JAPAN]
- Propylea japonica (Thunberg, 1781) (*Coccinella*) [CHINA: Heilongjiang, Jilin, Liaoning, Inner Mongolia, Hebei, Shanxi, Shaanxi, Qinghai, Gansu, Ningxia, Xinjiang, Henan, Shandong, Hubei, Anhui, Jiangsu, Jiangxi, Zhejiang, Fujian, China; CHINA: Taiwan, Guangdong, Hunan, Guangxi, Guizhou, Sichuan, Yunnan; INDIA, THAILAND, JAPAN, Former USSR] Prey on: Matsucoccus matsumurae Kuwana, aphids on Phaseolus, Aphis craccivora Koch, A. gossypii Glover, Lipaphis erysimi (Kaltenbach), Longium sacchari (Zehntner), Macrosiphum avenae Fabricius, Rhopalosiphum maidis Fitch, Schizaphis graminum (Rondani).
- Propylea luteopustulata (Mulsant, 1850) (*Oenopia*) Syn. *Coleophora korschefskeyi* Mader, 1926; *Oenopia hauseri* Mader, 1926; *Coleophora victoriae* Mulsant, 1866; *Coleophora insularis* Sicard, 1912; *Coleophora weisei* Miwa, 1931; *Coleophora octopunctata* Yablokoff - Kunzoryan, 1982; *Anatis tibetina* Mulsant, 1853; *Coleophora pedicata* Mulsant, 1853; *Coleophora mariae* Mulsant, 1853; *Caria thoracica* Weise, 1891; *Oenopia hauseri coalescens* Mader, 1930 [CHINA: Sichuan, Xizang; NEPAL, MYANMAR, SIKKIM]
- Propylea quatuordecimpunctata (Linnaeus, 1759) (*Coccinella*) Syn. *Propylea conglobata* Mulsant, 1866 [CHINA: Heilongjiang, Jilin, Liaoning, Inner Mongolia, Hebei, Henan, Shaanxi, Gansu, Xinjiang, Hubei, Jiangsu, Jiangxi, Fujian, Taiwan, Hongkong, Guizhou; JAPAN, KOREA, MONGOLIA, Former USSR, EUROPE, AFRICA] Prey on: aphids, scale insects, whiteflies.
- Pseudaspiderus lombatus Hoang, 1983 [CHINA: Hainan; VIETNAM]
- Pseudoscyrmus amplus Yang et Wu, 1972 [CHINA: Taiwan]
- Pseudoscyrmus ancistroides Pang et Huang, 1986 [CHINA: Fujian]
- Pseudoscyrmus anmashanus Yang, 1971 [CHINA: Taiwan]
- Pseudoscyrmus changi Yang, 1971 [CHINA: Taiwan]
- Pseudoscyrmus dapae Hoang, 1978 [CHINA: Henan, Fujian, Guangdong; VIETNAM]
- Pseudoscyrmus disselasmatus Pang et Haung, 1986 [CHINA: Henan, Fujian]
- Pseudoscyrmus fulvifemur Yang et Wu, 1972 [CHINA: Taiwan; JAPAN]
- Pseudoscyrmus fuscus Yang, 1971 [CHINA: Taiwan]
- Pseudoscyrmus hareja (Weise, 1879) (*Scymnus*) [CHINA: Taiwan, Guizhou; JAPAN, VIETNAM] Prey on: Pseudaulacaspis pentagona (Targioni), Unaspis yanonensis Kuwana.
- Pseudoscyrmus kurohime (Miyatake, 1969) (*Scymnus*) [CHINA: Fujian, Taiwan, Guangdong, Hainan, Guangxi, Guizhou, Yunnan; JAPAN, VIETNAM, MICRONESIA] Prey on: mealybugs, Ceratavavcuna lanigera Zehntner, Dysmicoccus brevipes (Cockerell).
- Pseudoscyrmus lamellatus Yang et Wu, 1972 [CHINA: Taiwan]
- Pseudoscyrmus lancetapicalis (Pang et Gordon, 1986) [CHINA: Guangdong]
- Pseudoscyrmus lewisi (Kamiya, 1961) (*Clitostethus*) Syn. *Clitostethus nakanei* Araki, 1964 [CHINA: Shaanxi, Hubei, Fujian; JAPAN, VIETNAM]
- Pseudoscyrmus montanus Yang, 1971 [CHINA: Taiwan]
- Pseudoscyrmus nagasakiensis (Kamiya, 1961) (*Clitostethus*) [CHINA: Taiwan; JAPAN]
- Pseudoscyrmus nepalius Miyatake, 1985 [CHINA: Guangdong; NEPAL]
- Pseudoscyrmus ohtai Yang et Wu, 1972 [CHINA: Hubei, Taiwan]
- Pseudoscyrmus orbiculatus Yang, 1971 [CHINA: Taiwan]
- Pseudoscyrmus paltatus Pang et Huang, 1986 [CHINA: Fujian]
- Pseudoscyrmus pronotus Pang et Huang, 1986 [CHINA: Fujian, Guangdong, Guizhou]
- Pseudoscyrmus quinquepunctatus (Weise, 1923) (*Nephus*) [CHINA: Taiwan, Hunan, Guizhou; JAPAN]
- Pseudoscyrmus seboskii (Ohta, 1929) [CHINA: Fujian, Guizhou; JAPAN]
- Pseudoscyrmus sylvaticus (Lewis, 1896) (*Scymnus*) [CHINA: Taiwan, Hainan; JAPAN, KOREA]
- Pseudoscyrmus tainanensis (Ohta,) (*Pullus*) [CHINA: Taiwan]
- Pssyllobora vigintiduopunctata (Linnaeus, 1758) (*Coccinella*) Syn. *Thea vigintiduopunctata* (Linnaeus, 1758) [CHINA: Henan,

- Henan, Shandong, Xinjiang, Sichuan; JAPAN, N. AMERICA] Prey on: powdery mildew on *Ailanthus*, fungi.
- Pullus niponicus* Lewis, 1896 [CHINA; Taiwan, JAPAN, KOREA]
- Pullus nubilus* Mulsant, 1850 [CHINA; Taiwan; INDIA]
- Pullus sodalis* Weise, 1923 [CHINA; Taiwan]
- Rhizobius formosanus* Weise, 1923 [CHINA; Taiwan]
- Rodolia breviscula* Weise, 1892 [CHINA: S. CHINA; INDIA, MYANMAR, SRI LANKA]
- Rodolia cardinalis* (Mulsant, 1850) (*Vedalia*) (introduced from AUSTRALIA) [CHINA: Shaanxi, Henan, Hubei, Jiangsu, Jiangxi, Zhejiang, Fujian, Taiwan, Guangdong, Hongkong, Guangxi, Guizhou, Sichuan, Yunnan; AUSTRALIA, JAPAN, INDONESIA, Former USSR, AMERICA, EUROPE, AFRICA] Prey on: *Icerya purchasi* Maskell.
- Rodolia concolor* (Lewis, 1879) (*Novius*) Syn. *Novius japonicus* (Crotch, 1873) [CHINA: Hebei, Henan, Gansu, Jiangsu; JAPAN] Prey on: *Tetraleurodes mori* (Quaintance), *Drosicha corpulenta* Kuwana.
- Rodolia formosana* Korschefsky, 1935 [CHINA; Taiwan]
- Rodolia fumida* Mulsant, 1850 Syn. *Rodolia roseipennis* Mulsant, 1850; *Rodolia testicolor* Mulsant, 1853 [CHINA: Fujian, Guangdong, Yunnan; INDIA, MYANMAR, SRI LANKA, BANGLADESH, MADAGASCAR]
- Rodolia guerini* (Crotch, 1874) (*Vedalia*) [CHINA: Zhejiang, Fujian, Guangdong, Guangxi, Sichuan, Yunnan; INDIA, SIKKIM]
- Rodolia hauseri* (Mader, 1930) (*Novius*) [CHINA: Yunnan]
- Rodolia limbata* (Motschulsky, 1866) (*Novius*) [CHINA: Heilongjiang, Jilin, Liaoning, Hebei, Shanxi, Shaanxi, Henan, Hubei, Jiangsu, Jiangxi, Zhejiang, Taiwan, Guangdong, Guangxi, Guizhou, Sichuan, Yunnan; JAPAN, KOREA, SIBERIA, MONGOLIA] Prey on: *Tetraleurodes mori* (Quaintance), *Icerya purchasi* Maskell, *Warajiococcus corpulentus* Kuwana, whiteflies.
- Rodolia marginata* Bielawski, Syn. *Rodolia arethusa* Mulsant, 1853; *Rodolia testicolor* Mulsant, 1853 [CHINA: Yunnan; INDIA] Prey on: scale insects.
- Rodolia octoguttata* Weise, 1910 [CHINA: Fujian, Guizhou, Sichuan, Yunnan; INDIA] Prey on: scale insects.
- Rodolia podagrica* Weise, 1908 [CHINA; PHILIPPINES, INDONESIA]
- Rodolia pumila* Weise, 1892 [CHINA: Xinjiang, Hubei, Zhejiang, Fujian, Guangdong, Hongkong, Hainan, Guangxi, Guizhou, Sichuan, Yunnan; JAPAN, MICRONESIA] Prey on: *Icerya purchasi* Maskell.
- Rodolia quadrimaculata* Mader, 1939 Syn. *Rodolia quadrimaculata* ab. *sexmaculata* Mader, 1939 [CHINA: Heilongjiang, Liaoning, Inner Mongolia, Henan, Hubei, Shaanxi, Taiwan, Guangdong, Hunan, Guangxi, Sichuan, Yunnan; JAPAN] Prey on: aphids, *Icerya purchasi* Maskell.
- Rodolia rubea* Mulsant, 1850 Syn. *Rodolia carneipellis* Mulsant, 1853 [CHINA: Hubei, Zhejiang, Fujian, Guangdong, Hainan; INDONESIA, CALEMANDAN, CELEBES] Prey on: scale insects.
- Rodolia rufocincta* Lewis, 1896 [CHINA: Beijing; JAPAN] Prey on: *Tetraleurodes mori* (Quaintance).
- Rodolia rufopilosa* Mulsant, 1850 [CHINA: Shaanxi, Gansu, Henan, Hubei, Anhui, Jiangsu, Jiangxi, Zhejiang, Fujian, Guangdong, Hongkong, Hainan, Hunan, Guangxi, Guizhou, Sichuan, Xizang; INDIA, MYANMAR, VIETNAM, INDONESIA, PHILIPPINES, JAPAN] Prey on: *Icerya purchasi* Maskell, *I. seychellerum* (Westwood), *Drosicha corpulenta* Kuwana, aphids, mites.
- Scotoscymnus bicolor* (Kamiya, 1965) (*Sukanakikona*) [CHINA: Taiwan; JAPAN]
- Scymnus accamptus* Pang et Pu, 1990 [CHINA: Guangxi]
- Scymnus acidotus* Pang et Huang, 1986 [CHINA: Fujian, Guangdong, Guizhou]
- Scymnus akonis* (Ohta, 1929) (*Pullus*) [CHINA; Taiwan]
- Scymnus ambonoidea* Pang, 1988 [CHINA: Guangdong, Guangxi]
- Scymnus andamanensis* Kapur, 1966 [CHINA: Shaanxi, Hunan, Guizhou; INDIA] Prey on: aphids.
- Scymnus apiciflavus* (Motschulsky, 1858) (*Diomus*) [CHINA: Shaanxi, Guangxi, Sichuan; INDIA, VIETNAM, SRI LANKA]
- Scymnus auritus* (Thunberg, 1795) (*Pullus*) Syn. *Pullus capitatus* Fabricius, 1798; *Pullus rufipes* Bris, 1863; *Pullus brisouti* Crotch, 1874; *Pullus fulvifrons* Stephen et Marsh, 1802; *Pullus minimus* Rossi, 1794 [CHINA: Anhui, Fujian, Guangdong, Sichuan; PALAEARCTIC]
- Scymnus babai* Sasaji, 1971 [CHINA: Heilongjiang, Jilin, Liaoning, Inner Mongolia, Hebei, Henan, Shandong, Shaanxi, Hubei, Jiangsu, Zhejiang, Fujian, Hunan, Sichuan, Yunnan; JAPAN, KOREA] Prey on: *Aphis gossypii* Glover, *A. pomi* De Geer, *Longiunguis scchari* (Zehntner), *Myzus persicae* (Silzer), *Rhopalosiphum maidis* Fitch, *Rh. padi* Linnaeus, *Rh. rufiabdominalis* (Sasaji).
- Scymnus brunnescens* Motschulsky, 1866 [CHINA: Fujian, Taiwan, Guangdong; INDIA, MYANMAR, SRI LANKA, INDONESIA, PHILIPPINES, JAPAN]
- Scymnus centralis* Kamiya, 1965 [CHINA: Taiwan, Guangdong, Guangxi, Hainan; JAPAN, VIETNAM]
- Scymnus chinensis* Jordan, 1894 [CHINA]
- Scymnus cnidatus* Pang et Pu, 1990 [CHINA: Guangxi]
- Scymnus comperei* Pang et Gordon, 1986 [CHINA: Guangdong] Prey on: *Panonychus citri* (McGregor).
- Scymnus compoceratus* Pang et Huang, 1986 [CHINA: Fujian]
- Scymnus contemptus* (Weise, 1923) (*Pullus*) [CHINA: Taiwan, Guangdong; JAPAN, VIETNAM]
- Scymnus crinitus* Fursch, 1966 [CHINA: Henan, Hebei, Hubei, Gansu, Sichuan]
- Scymnus curvus* Yang, 1978 [CHINA: Taiwan]
- Scymnus dactylicus* Pang, 1986 [CHINA: Guangdong]
- Scymnus dichorionicus* Pang et Huang, 1986 [CHINA: Hubei, Fujian]
- Scymnus dicorycus* Pang et Huang, 1986 [CHINA: Fujian, Guangdong, Hainan; VIETNAM]
- Scymnus dissolobus* Pang et Huang, 1985 [CHINA: Hubei, Fujian, Guangdong]
- Scymnus dorcatomoides* Weise, 1879 [CHINA: Hubei, Fujian, Taiwan; JAPAN, KURILE, VIETNAM]
- Scymnus dorytatus* Pang, 1988 [CHINA: Guangdong]
- Scymnus exocorycus* Pang et Huang, 1986 [CHINA: Fujian, Hainan]
- Scymnus ferrugatus* (Moll, 1885) (*Pullus*) [CHINA: Shaanxi, Fujian, Hunan; AISA, EUROPE, FRANCE, POLAND, PALAEARCTIC] Prey on: *Myzus persicae* (Sulzer).
- Scymnus folchini* Canepari, 1979 [CHINA: Hebei, Shandong, Zhejiang, Hainan, Sichuan]
- Scymnus formosanus* (Weise, 1923) (*Rhizobius*) [CHINA: Zhejiang, Taiwan, Guangdong, Guangxi, Guizhou, Sichuan; VIETNAM]
- Scymnus frontalis* Fabricius, 1878? [CHINA: Heilongjiang, Hebei, Henan, Shaanxi, Gansu, Xinjiang, Shandong, Hubei, Jiangxi, Zhejiang, Fujian; PALAEARCTIC, EUROPE, ASIA] Prey on: *Aphis gossypii* Glover, *A. robiniae* macchiati.
- Scymnus fujianensis* Pang et Gordon, 1986 [CHINA: Fujian]
- Scymnus fuscatus* Boheman, 1858 Syn. *Scymnus niponicus* Lewis, 1896; *Scymnus niponicus* ab. *nigripes* Ohta, 1929; *Scymnus niponicus* ab. *munaguronis* Ohta, 1929 [CHINA: Taiwan; INDIA, MYANMAR, THAILAND, CAMBODIA, SRI LANKA, INDONESIA, PHILIPPINES, SE ASIA, JAPAN, KOREA]

- Scymnus giganteus* Kamiya, 1961 [CHINA: Guangdong; JAPAN, VIETNAM]
- Scymnus hainanensis* Pang et Gordon, 1986? [CHINA: Hainan]
- Scymnus hatomensis* Kamiya, 1965 [CHINA: Taiwan; JAPAN]
- Scymnus hilaris* Motschulsky, 1858 Syn. *Pullus hilaris* ab. *avarus* Ohta, 1929; *Scymnus ishidae* Araki, 1963 [CHINA: Fujian, Taiwan; JAPAN, MYANMAR, INDIA, SRI LANKA] Prey on: scale insects.
- Scymnus hingstoni* Kapur, 1963 [CHINA: Fujian, Guangdong; INDIA, SIKKIM, VIETNAM]
- Scymnus hoffmanni* Weise, 1879 [CHINA: Heilongjiang, Jilin, Liaoning, Hebei, Henan, Shandong, Shaanxi, Gansu, Hubei, Anhui, Jiangsu, Jiangxi, Zhejiang, Fujian, Taiwan, Hunan, Guangxi, Sichuan, Yunnan; INDOCHINA, JAPAN, KOREA] Prey on: *Aphis gossypii* Glover, *Ceratovacuna lanigera* Zehntner, *Longiunguis sacchari* (Zehntner), *Schizaphis graminum* (Rondani), *Thrips oryzae* Williams, scale insects, spiders, mites, larvae of *Heliothis armigera* Huber.
- Scymnus impexus* Mulsant, 1850 [CHINA: Guangdong; EUROPE, N. AMERICA, N. AFRICA]
- Scymnus inderihensis* Mulsant, 1850 [CHINA: Shanxi, Shaanxi, Gansu, Xinjiang, Hebei, Hubei; MONGOLIA Former USSR]
- Scymnus ionochiatus* Pang et Huang, 1985 [CHINA: Fujian]
- Scymnus japonicus* Weise, 1879 [CHINA: Jilin, Liaoning, Hebei, Henan, Shandong, Hubei, Jiangsu, Jiangxi, Zhejiang, Fujian, Hainan, Guizhou, Sichuan, Yunnan, Xizang; JAPAN, KOREA] Prey on: aphids.
- Scymnus kabakovi* Hoang, 1982 [CHINA: Guangdong, Yunnan; VIETNAM]
- Scymnus kaguyahime* Kamiya, 1961 [CHINA: Zhejiang, Fujian; JAPAN]
- Scymnus kawamurai* (Ohta, 1929) (*Pullus*) [CHINA: Henan, Shaanxi, Shandong, Hubei, Anhui, Jiangsu, Zhejiang, Fujian, Guangdong, Hainan, Guizhou, Sichuan, Yunnan; JAPAN, INDIA] Prey on: *Aphis gossypii* Glover, *Macrosiphum avenae* Fabricius, *Myzus persicae* (Sulzer), *Schizaphis graminum* (Rondani), *Toxoptera citricida* (Kirkaldy).
- Scymnus klapperichi* Pang et Gordon, 1986 [CHINA: Fujian, Taiwan, Guangxi]
- Scymnus koebelei* (Pang et Gordon, 1986) [CHINA: Fujian, Guangdong, Hainan, Guangxi]
- Scymnus lenus* Yang, 1978 [CHINA: Fujian, Taiwan, Guangdong, Yunnan; VIETNAM]
- Scymnus leo* Yang, 1978 [CHINA: Taiwan]
- Scymnus lonchiatus* Pang et Huang, 1985 [CHINA: Fujian, Yunnan]
- Scymnus longisiphonulus* Cao et Xiao, 1984? [CHINA: Yunnan] Prey on: aphids on *Ficus*.
- Scymnus longmericus* Pang, 1986 [CHINA: Guangdong, Hainan]
- Scymnus manipulus* Fursch et Kreissl, 1967 [CHINA: Gansu, Jiangsu; Former UUSSR, CAUCASUS, MONGOLIA, C. ASIA]
- Scymnus minus* Capra et Fursch, 1967 [CHINA: Heilongjiang, Hebei, Shaanxi, Gansu, Xinjiang, Shandong, Hubei, Jiangxi, Zhejiang, Fujian, Guizhou, Yunnan; GERMANY, FRANCE] Prey on: *Aphis gossypii* Glover, *A. robiniae* Macchiatii.
- Scymnus mongolicus* Weise, 1890 [CHINA: Inner Mongolia, Gansu; MONGOLIA, IRAN, SAUDI ARABIA]
- Scymnus nankunicus* Pang, 1986 [CHINA: Fujian, Guangdong, Hainan]
- Scymnus nepentus* Pang et Huang, 1985 Syn. *Scymnus perdere* Yang, 1978 [CHINA: Zhejiang, Fujian, Taiwan, Guangdong]
- Scymnus nigrosuturalis* Kamiya, 1961 [CHINA: Taiwan; JAPAN, MICRONESIA]
- Scymnus notidamus* Pang et Huang, 1985 [CHINA: Fujian]
- Scymnus nubilus* Mulsant, 1850 [CHINA: Taiwan; INDIA, MYANMAR, NEPAL]
- Scymnus obsoletus* Weise, 1890 [CHINA: Xizang]
- Scymnus oestocraerus* Pang et Huang, 1985 [CHINA: Fujian, Taiwan, Hainan, Guangxi, Guizhou; VIETNAM]
- Scymnus oncosiphonus* Cao et Xiao, 1984 [CHINA: Yunnan] Prey on: aphids on *Ficus*.
- Scymnus orientalis* Mader, 1954 [CHINA: Fujian]
- Scymnus ovmaculatus* Sasaji, 1968 [CHINA: Guangdong; MALAYSIA]
- Scymnus paganus* (Lewis, 1896) (*Pullus*) [CHINA: Hubei, Taiwan, Sichuan; JAPAN] Prey on: aphids.
- Scymnus pangi* Fursch, 1989 Syn. *Scymnus notus* Pang et Pu, 1990 [CHINA: Taiwan, Guangdong, Hainan, Guangxi]
- Scymnus pelecoides* Pang et Huang, 1985 [CHINA: Zhejiang, Fujian]
- Scymnus postcalis* Sicard, 1912 [CHINA: Shaanxi, Henan, Hubei, Fujian, Taiwan, Guangdong, Guangxi, Guizhou, Yunnan; MYANMAR, INDIA, THAILAND, VIETNAM, SRI LANKA, JAPAN] Prey on: *Rhopalosiphum maidis* Fitch, aphids.
- Scymnus prosericatus* Pang, 1988 [CHINA: Guangdong]
- Scymnus prosphykontus* Pang et Huang, 1985 [CHINA: Fujian]
- Scymnus prostylotus* Pang et Huang, 1985 [CHINA: Fujian]
- Scymnus quadrillum* Motschulsky, 1858 [CHINA: Taiwan; INDIA, THAILAND, SRI LANKA, JAPAN]
- Scymnus quadrivulneratus* Mulsant, 1850 ? [CHINA: Hebei, Shanxi, Shaanxi, Gansu, Xinjiang, Henan; MONGOLIA, Former USSR] Prey on: *Aphis gossypii* Glover.
- Scymnus rhachiatus* Pang et Huang, 1985 [CHINA: Fujian]
- Scymnus rhamphiatus* Pang et Huang, 1985 [CHINA: Hubei, Zhejiang, Fujian, Hunan, Guizhou, Sichuan]
- Scymnus ruficeps* (Ohta, 1929) (*Pullus*) [CHINA: Taiwan; JAPAN]
- Scymnus scapanulus* Pang et Huang, 1986 [CHINA: Zhejiang, Fujian]
- Scymnus secula* Yang, 1971 [CHINA: Taiwan]
- Scymnus shirozui* Kamiya, 1965 [CHINA: Taiwan, Guangdong, Hainan]
- Scymnus singularis* Mader, 1954 [CHINA: Fujian]
- Scymnus sodalis* (Weise, 1923) (*Pullus*) [CHINA: Henan, Hubei, Jiangsu, Zhejiang, Fujian, Taiwan, Guangdong, Guangxi, Sichuan, Xizang; INDIA, VIETNAM, NEPAL, JAPAN] Prey on: mealybugs, aphids, red mites.
- Scymnus sternitus* Pang et Gordon, 1986 [CHINA: Guangxi]
- Scymnus syoitii* Sasaji, 1971 [CHINA: Fujian, JAPAN, VIETNAM]
- Scymnus tainanensis* (Ohta, 1929) (*Pullus*) [CHINA: Taiwan]
- Scymnus taiwanus* (Ohta, 1929) (*Pullus*) [CHINA: Zhejiang, Fujian, Taiwan, Guangdong, Hainan, Guangxi; VIETNAM, PHILIPPINES] Prey on: aphids.
- Scymnus takabayashii* (Ohta, 1929) (*Pullus*) [CHINA: Fujian, Hunan, Guizhou; JAPAN]
- Scymnus takasago* Kamiya, 1965 [CHINA: Taiwan]
- Scymnus tegminalis* Hoang, 1985 [CHINA: Fujian, Guangdong, VIETNAM]
- Scymnus tenuis* Yang, 1978 [CHINA: Fujian, Taiwan, Guangdong, Hainan, Yunnan; VIETNAM]
- Scymnus toxosiphonius* Pang et Huang, 1986 [CHINA: Fujian, Guizhou]
- Scymnus villis* (Weise, 1923) (*Pullus*) [CHINA: Taiwan]
- Scymnus xanthostethus* Pang et Pu, 1990 [CHINA: Guangxi]
- Scymnus yamato* Kamiya, 1961 [CHINA: Hebei, Henan, Jiangsu, Fujian, Guangdong, Sichuan; JAPAN, VIETNAM] Prey on: Aphid

- craccivora Koch, *A. gossypii* Glover, *Rhopalosiphum maidis* Fitch, Rh. nymphaeae Linnaeus.
- Scymnus yangi* Yu et Pang, 1992 Syn. *Scymnus bicolor* Yang, 1978; *Scymnus endocorycus* Pang et Huang, 1986; *Scymnus vishphuens* Hoang, 1982 [CHINA: Zhejiang, Fujian, Taiwan, Guangdong, Hainan, Hunan, Guizhou; VIETNAM]
- Scymnus yatomensis* Komiya, 1965 [CHINA: Hainan; JAPAN]
- Semiadalia decimguttata* Jing, 1986 [CHINA: Xizang]
- Senorycha grandis* (Thunberg, 1781) (*Coccinella*) [CHINA: NE CHINA, Fujian, Guangdong, Guangxi, Yunnan; INDIA, MYANMAR, INDONESIA, PHILIPPINES, JAPAN] Prey on: *Aphis bambusae* Fullaway.
- Serangium japonicum* Chapin, 1940 [CHINA: Shaanxi, Henan, Hubei, Jiangsu, Zhejiang, Fujian, Taiwan, Guangdong, Guangxi, Guizhou, Sichuan, Yunnan; JAPAN, KOREA] Prey on: *Aleurocanthus spiniferus* (Quaintance), *A. philippinensis* Quaintance et Baker, *A. aucubae* Kuwana, scale insects on Citrus.
- Shirozuella mirabilis* Sasaji, 1967 [CHINA: Taiwan]
- Singhikalia ornata* Kapur, 1963 [CHINA: Taiwan; VIETNAM, INDIA]
- Singhikalia subfasciata* Miyatake, 1972 [CHINA: Taiwan]
- Sospita gebleri* (Crotch, 1874) (*Mysia*) Syn. *Coccinella eramosa* Faldermann, 1833; *Noemysia kasai* Kurisaki, 1920 [CHINA: Inner Mongolia; JAPAN, SIBERIA]
- Sospita horni* (Crotch, 1872) [CHINA: Shaanxi; AMERICA] Prey on: aphids.
- Sospita oblongoguttata* (Linnaeus, 1758) (*Coccinella*) [CHINA: Heilongjiang, Jilin, Qinghai, Sichuan; JAPAN, KOREA, ASIA, EUROPE] Prey on: aphids.
- Sospita quadrivittata* Miyatake, 1965 [CHINA: Taiwan]
- Sphaerplotina hainanensis* Miyatake, 1969 [CHINA: Hainan]
- Spiladelphia longula* Barowsky, 1925? [CHINA: Xizang]
- Stethorus acuminatus* Motschulsky, [CHINA: Fujian]
- Stethorus aptus* Kapur, 1948 [CHINA: Zhejiang, Fujian, Taiwan, Guangdong, Guangxi; MALAYSIA, JAPAN] Prey on: *Panonychus citri* (McGregory).
- Stethorus binchuanensis* Pang et Mao, 1975 [CHINA: Sichuan, Yunnan] Prey on: *Panonychus citri* (McGregory).
- Stethorus cantonensis* Pang, 1966 [CHINA: Hubei, Jiangxi, Fujian, Guangdong, Hainan, Guangxi, Sichuan] Prey on: *Panonychus citri* (McGregory), *Oligonychus coffeae* (Nietner), red mites on *Carica papaya*.
- Stethorus chengi* Sasaji, 1968 [CHINA: Shaanxi, Henan, Hubei, Jiangsu, Zhejiang, Taiwan, Hunan, Guizhou, Sichuan] Prey on: red mites on beans, cotton, apple and Citrus.
- Stethorus dongchuanensis* Cao et Xiao, 1984 [CHINA: Yunnan] Prey on: *Panonychus citri* (McGregory).
- Stethorus gilvifrons* (Mulsant, 1850) [CHINA: Hongkong]
- Stethorus guangxiensis* Pang et Mao, 1975 [CHINA: Shaanxi, Hubei, Fujian, Guangxi, Sichuan] Prey on: *Panonychus citri* (McGregory).
- Stethorus hirashimai* Sasaji, 1968 [CHINA: Taiwan] Prey on: mites.
- Stethorus indica* Kapur, 1950 [CHINA: Guangxi, Yunnan; INDIA] Prey on: mites.
- Stethorus loi* Sasaji, 1968 [CHINA: Taiwan] Prey on: mites.
- Stethorus longisiphonulus* Pang, 1966 [CHINA: Fujian, Guangdong, Hainan; VIETNAM] Prey on: red mites on *Carica papaya*.
- Stethorus parapauperculus* Pang, 1966 [CHINA: Henan, Fujian, Guangdong, Hainan, Guangxi, Yunnan] Prey on: *Panonychus citri* (McGregory), red mites on *Saccharum sinensis*, *Carica papaya*.
- Stethorus punctillum* Weise, 1885? [CHINA: Heilongjiang, Liaoning, Hebei, Shandong, Henan, Shaanxi, Gansu, Xinjiang, Hubei, Jiangsu, Jiangxi, Zhejiang, Fujian, Guangxi, Sichuan; JAPAN, Former USSR, EUROPE, IRAN, ISRAEL, PALAEARCTIC] Prey on: *Panonychus citri* (McGregory), *Tetranychus urticae* (Koch), red mites on *Malus pumila*, *Morus alba*, *Pyrus*, *Crataegus pinnatifida*.
- Stethorus rani* Kapur, 1948 [CHINA: Fujian; MYANMAR, THAILAND, INDIA] Prey on: red mites on Citrus.
- Stethorus rotundatus* Motschulsky, 1859 [CHINA: Taiwan; INDIA, MYANMAR, PHILIPPINES, SRI LANKA]
- Stethorus shaanxiensis* Pang et Mao, 1975 [CHINA: Shaanxi, Gansu] Prey on: *Panonychus ulmi* (Koch), red mites on apple.
- Stethorus siphonulus* Kapur, 1948 [CHINA: Hubei, Fujian, Guangdong, Guangxi; MALAYSIA, THAILAND, HAWAII] Prey on: *Panonychus citri* (McGregory), *Oligonychus coffeae* (Nietner), *Polyphgotarsonemus latus* (Bank).
- Stethorus truncatus* Kapur, 1948 [CHINA: Guangdong; MALAYSIA]
- Stethorus vietnamicus* Hoang, 1979 [CHINA: Hainan, Guangxi; VIETNAM] Prey on: red mites on Citrus.
- Stethorus yingjiangensis* Cao et Xiao, 1984 [CHINA: Guizhou, Yunnan] Prey on: red mites on *Ficus*.
- Stethorus yunnanensis* Pang et Mao, 1979 [CHINA: Sichuan, Yunnan] Prey on: *Panonychus citri* (McGregory).
- Sticholotis confucii* Mulsant, 1850 [CHINA: Guangdong; MALAYSIA, SARAWAK]
- Sticholotis formosana* Weise, 1923 Syn. *Sticholotis substriatus* ab. *formosana*, Korschefsky, 1933 [CHINA: Taiwan, Guangdong, Hainan; CONGO]
- Sticholotis hirashimai* Sasaji, 1967 [CHINA: Taiwan]
- Sticholotis klapperichi* Mader, 1954 [CHINA: Fujian]
- Sticholotis maculata* Korschefsky, 1944 [CHINA: Taiwan]
- Sticholotis morimotoi* Kamiya, 1965 [CHINA: Taiwan; JAPAN]
- Sticholotis punctata* Crotch, 1874 [CHINA: Hubei, Fujian, Guizhou; JAPAN, PHILIPPINES, MALAYSIA] Prey on: *Parlatoria pergandi* Comstock, *Quadraspidiotus perniciosus* Comstock.
- Sticholotis ruficeps* Weise, 1902 [CHINA: Henan, Anhui, Guangdong; MALAYSIA, THAILAND, MICRONESIA] Prey on: aphids, giant coccids.
- Sticholotis taiwanensis* Miyatake, 1982 [CHINA: Taiwan]
- Sticholotis undecimpunctata* Miyatake, 1982 [CHINA: Taiwan]
- Subcoccinella vigintiquatuor punctata* (Linnaeus, 1758) (*Coccinella*) [CHINA: Liaoning; EUROPE, AFRICA, COAST of MEDITERRANEAN] Host: *Medicago*.
- Sukunakikona bicolor* Kamiya, 1965 [CHINA: Taiwan; JAPAN]
- Sukunakikona japonica* Kamiya, 1960 [CHINA: Taiwan; JAPAN]
- Sumnius brunneus* Jing, 1983 [CHINA: Henan, Sichuan, Yunnan] Prey on: *Warajiococcus corpulentus* Kuwana.
- Sumnius nigrofuscus* Jing, 1983 [CHINA: Yunnan]
- Sumnius petiolimaculatus* Jing, 1983 [CHINA: Yunnan]
- Sumnius vestitus* (Mulsant, 1850) Syn. *Sumnius cardoni* Weise, 1892 [CHINA: Fujian, Sichuan, Yunnan; VIETNAM, INDIA] Prey on: scale insects.
- Sumnius yunnanus* Mader, 1954 [CHINA: Shaanxi, Yunnan] Prey on: scale insects.
- Synonycha grandis* (Thunberg, 1781) (*Coccinella*) [CHINA: Jiangxi, Fujian, Taiwan, Guangdong, Hongkong, Hunan, Guangxi, Guizhou, Yunnan, Xizang; INDIA, INDONESIA, HIMALAYA, NEPAL, SIKKIM, MYANMAR, ASIA, JAPAN, NEW GUINEA, SRI LANKA, PHILIPPINES, AUSTRALIA] Prey on: aphids, aphids on *Camellia oleifera*, *Ceratovacuna lanigera* Zehntner, *Aphis saliceti* Kaltenbach.
- Telsimia chujoi* Miyatake, 1959 [CHINA: Taiwan; JAPAN]
- Telsimia emarginata* Chapin, 1926 [CHINA: Jilin, Henan, Hubei, Zhejiang, Fujian, Guangdong, Guangxi, Sichuan] Prey on: *Chrysophalus speermi*, *Ch. bifasciculatus* Ferris, *Ch. aonidium* (Linnaeus), *Aonidiella aurantii* (Maskell), *Parlatoria ziziphus*

- Lucas, *Pseudaulaspis pentagona* (Targioni), *Unaspis yanonensis* Kuwana.
- Telsimia huiliensis* Pang et Mao, 1979 [CHINA: Sichuan, Yunnan] Prey on: armored scale.
- Telsimia jingyangiensis* Pang et Mao, 1979 [CHINA: Guangxi, Sichuan, Yunnan] Prey on: armored scale.
- Telsimia nigra* (Weise, 1879) (*Penilia*) [CHINA: Fujian, Taiwan, Sichuan, Yunnan; JAPAN, PHILIPPINES] Prey on: *Parlatoria ziziphus* Lucas.
- Telsimia nigra centralis* Pang et Mao, 1979 [CHINA: Hubei, Fujian, Taiwan, Sichuan, Yunnan] Prey on: armored scale, *Aleurocanthus spiniferus* (Quaintance).
- Telsimia scymnoides* Miyatake, 1978 [CHINA: Taiwan]
- Telsimia shirozui* Miyatake, 1965 [CHINA: Taiwan]
- Telsimia sichuanensis* Pang et Mao, 1979 [CHINA: Shaanxi, Fujian, Guangxi, Sichuan] Prey on: *Parlatoria ziziphus* Lucas, *Pseudaulacaspis major* (Cockerell).
- Tythaspis sedecimpunctata* (Linnaeus, 1758) (*Coccinella*) [CHINA: Inner Mongolia; TURKEY, MONGOLIA, Former USSR, AFRICA]
- Vibidia duodecimguttata* (Poda, 1761) (*Thea*) [CHINA: Beijing, Henan, Shaanxi, Gansu, Hubei, Jiangsu, Fujian, Guangdong, Hunan, Guangxi, Sichuan, Yunnan; JAPAN, KOREA, SIBERIA, CAUCASUS, MINOR ASIA, C. ASIA, EUROPE] Feeding on: powdery mildew of *Ailanthus*.
- Vibidia huiliensis* Pang et Mao, 1979 [CHINA: Sichuan] Prey on: fungi.
- Vibidia korschevskyi* (Mader, 1930) (*Halysia*) [CHINA: Fujian, Sichuan, Shaanxi, Yunnan] Feeding on: powdery mildew.
- Vibidia luliangensis* Cao et Xiao, 1984 [CHINA: Sichuan, Yunnan] Feeding on: powdery mildew on *Malus pumila*.
- Vibidia xichanensis* Pang et Mao, 1979 [CHINA: Sichuan]
- Xanthadalia hieki* Yablokov - Kunzoryan, 1979 [CHINA: Sichuan, Yunnan, Xizang]
- Xanthadalia motuogensis* Jing, 1988 [CHINA: Xizang]
- (XIV) SUPERFAMILY MORDELLOIDEA
- Family **Cephaloidea**
- Cephaloon pallens* (Motschulsky, 1860) (*Sponidium*) [CHINA: NE CHINA; JAPAN, KOREA, SIBERIA]
- Family **Oedemeridae**
- Anacosessina tarsalis* Kono, 1937 [CHINA: Taiwan; JAPAN]
- Anocodes femorata* Ganglbauer, 1890 [CHINA: Gansu]
- Asclera binotata* Pic, 1927 [CHINA: Yunnan; or VIETNAM]
- Asclera brevicollis* Gressitt, 1939 [CHINA: Jiangxi, Fujian, Guangdong]
- Asclera diluta* Gressitt, 1939 [CHINA: Guangdong]
- Asclera formosana* (Pic, 1910) (*Ishnomera*) [CHINA: Taiwan]
- Asclera incostata* Pic, 1926 Syn. *Asclera incostata* var. *obscurimembra* Pic, *Asclera incostata* var. *ruficornis* Pic, 1926 [CHINA: Gansu, Xizang]
- Asclera licenti* Pic, [CHINA]
- Asclera loi* Gressitt, 1939 [CHINA: Hainan]
- Asclera maculicollis* Ganglbauer, 1890 [CHINA: Gansu]
- Asclera obsoleta* Ganglbauer, 1890 [CHINA: Shanxi]
- Asclera rugosipennis* (Pic, 1910) (*Ishnomera*) [CHINA: Taiwan]
- Asclera subrugosa* Kono, 1937 [CHINA: Taiwan; JAPAN]
- Asclera sulcicollis* Pic, 1929 [CHINA: NE CHINA]
- Chrysanthera intergricollis* Heyden, 1886 [CHINA: AMUR, SIBERIA]
- Copidita testaceopyga* Pic, 1923 [CHINA]
- Ditylus laevis* (Fabricius, 1887) [CHINA: NE CHINA; JAPAN, KOREA, SIBERIA, KURILE, SACHALIN, EUROPE]
- Eobia ambusta* Lewis, 1895 [CHINA: Taiwan; JAPAN]
- Eobia chinensis* (Hope, 1842) (*Nacerdes*) Syn. *Eobia chinensis* var. *kotoensis* Kono, 1937; *Sessinia chinensis* (Hope, 1842) [CHINA: Zhejiang, Taiwan; JAPAN]
- Eobia curticornis* Pic, 1911 [CHINA: Taiwan]
- Eobia florilega* Lewis, 1895 [CHINA: Mainland of CHINA, Taiwan; JAPAN]
- Eobia formosana* Pic, 1923 [CHINA: Taiwan]
- Eobia pallidipennis* Gressitt, 1939 [CHINA: Hainan]
- Eobia piceopyga* Pic, 1923 [CHINA]
- Eobia testaceiceps* Pic, 1923 [CHINA]
- Ganglbaueria collaris* Semenov, 1891 [CHINA: Xinjiang]
- Homomorpha crucifera* Semenov, 1898 [CHINA: Xinjiang]
- Nacerda coarctata manchurica* Magistretti, [CHINA: NE CHINA]
- Nacerda davidis* Fairmaire, 1886 [CHINA: Gansu, Sichuan; SIBERIA]
- Nacerda melanura* (Linnaeus, 1758) [CHINA: Taiwan; JAPAN]
- Nacerda stangulata* (Fairmaire, 1889) (*Anocodes*) [CHINA: Sichuan]
- Oedemera analis* Fairmaire, 1888 [CHINA: Beijing]
- Oedemera angusticollis* Fairmaire, 1888 [CHINA: Beijing]
- Oedemera angustipennis* Gressitt, 1939 [CHINA: Guangdong]
- Oedemera flaviventris* Fairmaire, 1891 [CHINA: Hubei, Sichuan]
- Oedemera inapicalis* Pic, 1910 [CHINA]
- Oedemera lurida* Marshall, 1802 [CHINA; SIBERIA, EUROPE, ALGERIA]
- Oedemera nasalis* Reitter, 1896 Syn. *Oedemera nasalis* var. *ros-trifera* Reitter, 1896 [CHINA: Xinjiang]
- Oedemera nigripes* Ganglbauer, 1889 [CHINA: Xinjiang]
- Oedemerus pallidipes* Pic, 1907 [CHINA]
- Oedemerus sauteri* Pic, 1927 [CHINA: Taiwan]
- Oedemerus testaceithorax* (Pic, 1936) [CHINA: Taiwan, Guangdong; JAPAN]
- Patalia testaceipennis* Pic, 1923 [CHINA: Taiwan]
- Peronocnemis davidis* Fairmaire, 1886 [CHINA: Sichuan]
- Probosca haemorrhoidalis* Pic, 1907 [CHINA]
- Probosca maculicollis* Pic, 1907 [CHINA]
- Pseudolycus uniformis nigriceps* Pic, 1909 [CHINA: Yunnan]
- Schistopselaphus sonani* Kono, 1937 [CHINA: Taiwan]
- Sessinia fuscotrigosa* Fairmaire, 1879 [CHINA]
- Sessinia sinensis* (Gemm. 1870) (*Ananca*) Syn. *Sessinia chinensis* Boheman, 1855 [CHINA: Hongkong]
- Sparedropsis davidis* Fairmaire, 1888 [CHINA: Beijing]
- Sparedropsis subserratus* Gressitt, 1939 [CHINA: Hainan]
- Xanthochroa apicalis* Kono, [CHINA: Taiwan]
- Xanthochroa atripennis* Pic, 1934 [CHINA]
- Xanthochroa baiharana* Kono, [CHINA: Taiwan]
- Xanthochroa davidis* (Fairmaire, 1886) (*Peronocnemis*) [CHINA: Gansu, Sichuan; SIBERIA]
- Xanthochroa fulvicrus* Fairmaire, 1889 [CHINA: Sichuan]
- Xanthochroa katoi* Kono, 1932 [CHINA: Hubei; JAPAN]
- Xanthochroa metallipennis* Fairmaire, 1889 [CHINA: Sichuan]
- Xanthochroa potanini* Ganglbauer, 1889 [CHINA: Sichuan]
- Xanthochroa semenovi* Ganglbauer, 1889 [CHINA: Gansu]
- Xanthochroa strangulata* (Fairmaire, 1889) (*Anocodes*) [CHINA: Sichuan]
- Xanthochroa taiwana* Kono, 1932 [CHINA: Taiwan]
- Xanthochroa waterhousei* Harold, 1875 Syn. *Xanthochroa waterhousei* var. *subniolacea* Pic, 1911 [CHINA: Hubei, Taiwan; KOREA, SACHALIN, JAPAN]
- Xanthochroa yunnana* Pic, 1923 [CHINA: Yunnan]