

Fig. 33. Male genitalia of *M. picta*. (A and B) Ventral and lateral views of phallobase without and with trabes; (C and D) entire sipho and enlarged view of apex. Scale bar = 0.2 mm.




Fig. 34. Male genitalia of *M. hudsonica*. (A and B) Ventral and lateral views of phallobase without and with trabes; (C and D) entire sipho and enlarged view of apex. Scale bar = 0.2 mm.

lawed before widdle cometimes normanion into




Fig. 35. Male genitalia of *M. quinquelineata*. (A and B) Ventral and lateral views of phallobase without and with trabes; (C and D) entire sipho and enlarged view of apex. Scale bar = 0.2 mm.




Fig. 36. Male genitalia of *M. luteodorsa*. (A and B) Ventral and lateral views of phallobase without and with trabes; (C and D) entire sipho and enlarged view of apex. Scale bar = 0.2 mm.

nating in Engel. spruce; 2 mi. W of Boulder; Durango, Junction Cr. Rd., 10,000 ft.; El Paso Co.; Estes Park, 9,000 ft.; Mt. Evans, Doolittle Ranch, 9,800 ft.; Mt. Manitou. *New Mexico:* Cloudcroft.




Fig. 37. Male genitalia of *M. cyathigera*. (A and B) Ventral and lateral views of phallobase without and with trabes; (C and D) entire sipho and enlarged view of apex. Scale bar = 0.2 mm.




Fig. 38. Male genitalia of *M. lynx*. (A and B) Ventral and lateral views of phallobase without and with trabes; (C and D) entire sipho and enlarged view of apex. Scale bar = 0.2 mm.




Fig. 39. Male genitalia of *M. labyrinthica*. (A and B) Ventral and lateral views of phallobase without and with trabes; (C and D) entire sipho and enlarged view of apex. Scale bar = 0.2 mm.






Fig. 40. Male genitalia of *M. mexicana* sp. nov. (A and B) Ventral and lateral views of phallobase without and with trabes; (C and D) entire sipho and enlarged view of apex. Scale bar = 0.2 mm.



**Fig. 41.** Male genitalia of *M. curva* sp. nov. (A and B) Ventral and lateral views of phallobase without and with trabes; (C and D) entire sipho and enlarged view of apex. Scale bar = 0.2 mm.

it and lateral spot; spots forming "M" enlarged or variously connected; lateral spots reduced or absent. In *lynx* form, elongate medial elytral spot



 $\textbf{Fig. 42-44.} \quad \textbf{Female genitalia.} \ \, \textbf{(42)} \ \, \textbf{\textit{M. picta}}; \ \, \textbf{(43)} \ \, \textbf{\textit{M. hudsonica}}; \ \, \textbf{(44)} \ \, \textbf{\textit{M. luteodorsa}}. \ \, \textbf{Scale bar} = 0.1 \ \, \text{mm.}$ 




Fig. 45-47. Female genitalia. (45) M. cyathigera; (46) M. lynx; (47) M. labyrinthica. Scale bar = 0.1 mm.




Fig. 48–50. Female genitalia. (48) M. curva sp. nov.; (49) M. mexicana sp. nov.; (50) M. quinquelineata. Scale bar = 0.1 mm.