

Aphids of the family Eriosomatidae (Insecta: Homoptera) in Belarus

Тли семейства Eriosomatidae (Insecta: Homoptera) Беларуси

S.V. BUGA, D.G. ZHOROV, N.V. LESHCHINSKAYA & A.V. STEKOLSHCHIKOV*

С.В. Буга, Д.Г. ЖОРОВ, Н.В. ЛЕЩИНСКАЯ, А.В. СТЕКОЛЬЩИКОВ

S.V. Buga, Department of Zoology, Belarusian State University, 4 Nezavisimosti av., 220030 Minsk, Republic of Belarus. E-mail: sergey.buga@gmail.com

D.G. Zhorov, Department of Zoology, Belarusian State University, 4 Nezavisimosti av., 220030 Minsk, Republic of Belarus. E-mail: zhorovdg@mail.ru

N.V. Lyashchynskaya, Scientific and Practical Centre for Bioresources, 27 Akademicheskaja st., 220072 Minsk, Republic of Belarus. E-mail: nadzeya.liashchynskaya@gmail.com

A.V. Stekolshchikov, Zoological Institute, Russian Academy of Sciences, 1 Universitetskaya nab., 199034 St Petersburg, Russia. E-mail: aphids@zin.ru

Investigation of Eriosomatidae in Belarus started in the 1900s. However, a regional checklist of Eriosomatidae aphids was not published till now. Based on material preserved in the collections of Belarusian State University and the Zoological Institute of the Russian Academy of Sciences, and data scattered across publications we have compiled the first comprehensive list of Eriosomatidae recorded from Belarus and briefly analysed it from the geographical and ecological points of views. The list includes 12 genera and 27 species. *Pemphigus passeki* Börner, 1952 has been recorded from Belarus for the first time. The registered diversity is less abundant than in the adjacent Poland. In the accordance with the global pattern of aphids host plant connections, a single species permanently inhabit woody plants (monoecious species), the other ones are dioecious and migrate from woody plants to woody (eight species) or herbaceous (12 species) plants. Four species is anholocyclic.

Исследования фауны тлей семейства Eriosomatidae в Беларуси были начаты еще в первые десятилетия XX века, однако до настоящего времени ни одного регионального списка эриосоматид не было опубликовано. Целью данной работы было обобщение сведений о Eriosomatidae Беларуси, имеющихся в разрозненных публикациях, которые были дополнены данными, полученными в ходе собственных исследований. Это позволило впервые составить аннотированный список эриосоматид фауны Беларуси и кратко проанализировать особенности их образа жизни. Список включает 27 видов из 12 родов. *Pemphigus passeki* Börner, 1952 впервые указан для фауны Беларуси. Выявленное на территории Беларуси число видов семейства оказалось меньше, чем в соседней Польше. Только один вид однодомен и постоянно обитает на одном растении-хозяине. Большинство тлей-эриосоматид фауны Беларуси являются двудомными и мигрируют с древесных растений на другие древесные (восемь видов) или травянистые (12 видов) растения; четыре вида неполноциклические, то есть развиваются без смены растений-хозяев.

Key words: gall-forming insects, monoecious, dioecious, Rhynchota, Homoptera, Eriosomatidae, Aphidinae, invasive species

Ключевые слова: галлообразующие насекомые, однодомные, двудомные, Rhynchota, Homoptera, Eriosomatidae, Aphidinae, инвазивные виды

* Corresponding author.

INTRODUCTION

Belarus, a small (207 600 km²) Central European country, is located in the transitional territory between two ecological zones: the European deciduous forests and the North Eurasian evergreen coniferous forests. The country is divided into six administrative provinces (Fig. 1).

Study on Eriosomatidae was started in the 1900s, when prof. A.K. Mordvilko collected aphids in different parts of Belarus. Two species of Eriosomatidae, *Eriosoma lanuginosum* (Hartig, 1839) and *Kaltenbachiella* (as *Tetraneura*) *pallida* (Haliday, 1838), are recorded for Grodno Province by A.K. Mordvilko in his monography (Mordvilko, 1924) devoted to the wooly apple aphid (*Eriosoma lanigerum* (Hausmann, 1802)) and other Eriosomea. In 1930, Mordvilko listed for Eastern (Soviet part) Belarus 28 aphid species, including 13 species of Eriosomatidae: *Eriosoma patchiae* (Börner et Blunck, 1916), *E. lanuginosum*, *Forda formicaria* Heyden, 1837, *F. marginata* Koch, 1857 (as *Forda trivialis* Pass.), *Geoica urticaria* (Passerini, 1856), *Pachypappa vesicalis* Koch, 1856, *P.* (as *Asiphum*) *tremulae* (Linnaeus, 1761), *Pachypappella* (as *Pachypappa*) *lactea* (Tullgren, 1909), *Paracletus cimiciformis* von Heyden, 1837, *Pemphigus bursarius* (Linnaeus, 1758), *P. populinigrae* (Schrank, 1801) (as *P. filaginis* B.d.F.), *P. spyrothecae* Passerini, 1856, and *Thecabius affinis* (Kaltenbach, 1843). At the same time several cecidological papers reported cecidia-forming aphis of Western Belarus (territory of Poland at that time). Seven species of Eriosomatidae (*Eriosoma ulmi* (Linnaeus, 1758), *Pachypappa marsupialis* Koch, 1856 (as *Pemphigus marsupialis* Courceh.), *Pemphigus bursarius*, *P. populinigrae* (as *P. filaginis* B. d. F.), *P. spyrothecae*, *Prociphilus xylostei* (de Geer, 1773), and *Thecabius* (as *Pemphigus*) *affinis*) were recorded by Z. Fiedorowiczowna (1930), and four species (*Pemphigus bursarius*, *P. spyrothecae*, *P. xylostei*, *Tetraneura* (as *Schizoneura*) *ulmi*) by J. Perepeczko-Baumanowa (1934).

In 1955, G.Ch. Shaposhnikov recorded six species of Eriosomatidae, *Eriosoma lanuginosum*, *E. patchiae*, *Pachypappa warschavensis* (Nasonov, 1894) (as *Asiphum varsoviensis* Mordv.), *P. tremulae* (as *Asiphum tremulae* Deg.), *Pemphigus borealis* Tullgren, 1909, and *Pemphigus immunitis* Buckton, 1896 (as *Pemphigus lichensteini* Tullgr.), as pests of forest woody plants in Belarus (Shaposhnikov, 1955). After the 1950s a study of aphids, pests of ornamental plants, was started in Belarus by N.L. Losinskaya (1960) who recorded two species of Eriosomatidae in parks. Information on Eriosomatidae species was also given in the monographies of S.V. Gorlenko & N.A. Pan'ko (1967, 1972), S.V. Gorlenko, A.I. Blintsov & N.A. Pan'ko (1988).

Comprehensive studies of the aphid fauna in Belarus were started in the 1980s. Till now several papers on the local fauna of Belarusian regions have been published (Buga, 1989, 1995; Stekolshchikov & Buga, 2006; Buga et al., 2008; Rakauskas & Buga, 2010; Stekolshchikov et al., 2008, 2010, 2012), and a special attention was paid to species damaging woody plants (Buga & Gorlenka, 1987; Buga, 1997, 2001; Buga & Stekolshchikov, 2009). However, a general regional checklist of Eriosomatidae has not been published yet.

MATERIAL AND METHODS

The paper is based on the material collected by authors since 1983. Although this information had been partly published in Russian or Belarusian, we prefer to repeat the collection data in English as electronic supplementary material to this paper. We also listed the material collected by T.V. Tikhomirov, F.V. Sautkin, D.L. Petrov, A.S. Roginsky & O.V. Sinchuk, and microscope slides of Eriosomatids collected by A.K. Mordvilko and storaged in the collection of the Zoological Institute of RAS (St Petersburg).

Microscope slides were prepared using Faure-Berlese mounting fluid or Canada balsam (Blackman & Eastop, 2000). The




Fig. 1. Administrative provinces of Belarus and geographic points of aphid sampling and registration.

material is deposited in the Department of Zoology of Belarusian State University (Minsk, Belarus) and the Zoological Institute of RAS (St Petersburg). Data extracted from publications are given in a uniform manner, but the completeness of the data depends on the original publications. When different variants of transliteration of geographical names are more or less widely used in the literature, we provide the additional variants in parentheses after the main variant. The special keys proposed by B. Osiadacz & R. Hałaj (Hałaj & Osiadacz, 2013; Osiadacz & Hałaj, 2014) have been used for identification of *Pemphigus* Hartig, 1839 species on the basis of galls formed on poplars. Aphid synonymy follows Remaudière and Remaudière (1997) with subsequent additions (Eastop & Blackman, 2005; Holman, 2009; Blackman & Eastop, 2011; Favret, 2016). Aphid taxonomic classification follows Shaposhnikov (1964). In this classification, Eriosomatidae (as Pemphigidae) is considered as a separate family

of Aphidoidea. This opinion was also supported by Heie & Wegierek (2009).

Abbreviations

Collectors: AM – A.K. Mordvilko, AR – A.S. Roginsky, DZh. – D.G. Zhorov, DP – D.L. Petrov, OS – O.V. Sinchuk, SB – S.V. Buga, VKh – V.I. Khvir, VT – V.N. Tikhomirov.

Localities: BBR – Berezina Biosphere Reserve, BPF – Belovezha Primeval Forest, CBG – Central Botany Gardens of the National Academy of Sciences of Belarus, hort. cv. – cultivars, NBR – Naliboki Biological Reserve, PLR – Priluki landscape reserve, RWS – railway station.

LIST OF SPECIES

The list of species is published as an electronic supplementary material which is available from:

<http://www.zin.ru/journals/zsr/content.asp?year=2016>.

DISCUSSION

Presently, the fauna of Belarus includes 27 species from 12 genera of Eriosomatidae. *Pemphigus paskevi* Börner, 1952 has been recorded from Belarus for the first time on the basis of material collected in 1900 by A.K. Mordvilko. *Pachypappa populi* (Linnaeus, 1758) was first mentioned by S.V. Buga (2001) with reference to A.K. Mordvilko (1930); however, reexamination of the cited publication revealed that Mordvilko (1930) did not record this species directly for Belarus. Three species of *Pachypappa* Koch, 1856 are considered for Belarus according to literature data only (Mordvilko, 1930; Fiedorowiczówna, 1930 and Shaposhnikov, 1955).

The species richness in Belarus is not so high compared with the neighbouring countries like Poland (49 species, 19 genera) (Hałaj, Osiadacz & Poljaković-Pajnik, 2016; Osiadacz & Hałaj, 2009, 2010, 2012; Wojciechowski et al., 2015). One of the reasons of this is the absence of such thermophilic species as *Aploneura lentisci* (Passerini, 1856) or *Mimeura ulmiphila* (del Guercio, 1917), etc. The Eriosomatidae were more extensively studied in Minsk Province (Fig. 1). Data from Belarus Lakeland (Poozer'e) were received mainly from the Gorodock Highland (Rakauskas & Buga, 2010).

The majority of registered species are common for Europe or the Palaearctic. At least one species, *Pemphigus borealis* Tullgren, 1909, is invasive and obviously has been brought from Far East (Shaposhnikov, 1955) with saplings of introduced host plants such as poplar *Populus laurifolia*. *Pemphigus spyrothecae* is common on *P. italicica* and very rare on other poplars. The species have never been registered on "wild" trees of *P. nigra* growing in floodplain of Pripyat' and other rivers in Belarusian Polesie.

As opposed to the majority of Eriosomatidae species, *P. spyrothecae* is monoecious. Dioecious aphids migrate from woody plants to woody (eight species) or herbaceous (12

species) plants. At least four Fordini species, *Forda formicaria* Heyden, 1837, *F. marginata* Koch, 1857, *Geoica urticularia* (Passerini, 1856), and *Prociphilus cimiciformis* von Heyden, 1837, inhabit plant roots and are anholocyclic under the condition of Belarus, because their primary host plants are exotic *Pistacia* spp. and absent in the local flora. Secondary host plants of *Pachypappa marsupialis* Koch, 1856 and *P. warshavensis* (Nasonov, 1894) are unknown. Poplars (*Populus* spp.) are the primary hosts of 14 species of Pemphigini, elms (*Ulmus* spp.) are the primary hosts of six species of Eriosomatini. Common spruce (*Picea abies*) is the secondary host plant for five Erisomatidae species. Host plant species are represented by eight families. One anholocyclic species, *P. cimiciformis*, is able to feed on the roots of a great variety of plants.

Some Erisomatidae species, *Pemphigus spyrothecae* Passerini, 1856, *P. bursarius* (Linnaeus, 1758), and *Eriosoma ulmi* (Linnaeus, 1758), are common and harmful in ornamental green stands in Belarus.

ACKNOWLEDGEMENTS

We are grateful to F.V. Sautkin, D.L. Petrov, A.S. Roginsky, O.V. Sinchuk, V.N. Tikhomirov for providing aphid and cecidological materials and to V.N. Tikhomirov (Belarusian State University, Minsk) for his help in the identification of host plants. This study was supported partly by the Belarusian Republic Foundation for Basic Research (grant no B15–063).

REFERENCES

- Blackman R.L. & Eastop V.F.** 2000. *Aphids on the World's Crops, an Identification and Information Guide*, 2nd Edition. Chichester: John Wiley & Sons. 476 p.
- Blackman R.L. & Eastop V.F.** 2011. Additions and amendments to "Aphids on the World's Plants". *Zootaxa*, **2774**: 57–68.
- Blackman R.L. & Eastop V.F.** 2016. *Aphids on the World's Plants. An online identification and information guide* [online]. Available from: www.aphidsontheworldsplants.info [updated 18 September 2016; visited 1 De-

- cember 2016].
- Buga S.V. & Gorlenka S.U.** 1987. Character of dendrocolous aphid complex organization in urban green stands. *Vesti Akademii nauk BSSR, Seriya biologicheskikh Nauk*, 2: 13–17. (In Belarusian).
- Buga S.V.** 1987. Plant introduction and dendrocolous aphid fauna of Byelorussia. In: *Migratsiya patogennykh organizmov pri introduksii rasteniy* [Migration of pathogenic organisms during plant introduction]: 87–89. Apatity: Kolskiy nauchnyy centr AN SSSR. (In Russian).
- Buga S.V.** 1988. *Aphids (Homoptera, Aphidinea) of the woody plants of the Central botanical Gardens of the Academy of Sciences of Belarusian SSR. 1. Families Phylloxeridae and Pemphigidae*. Moscow: VINITI. 10 p. (In Russian).
- Buga S.V.** 1989a. Order Homoptera. In: **Sokolov V.E.** (Ed.). *Flora i fauna zapovednikov SSSR: Nasekomye Berezinskogo zapovednika* [Flora and Fauna of the USSR reserves: Insects of Berezina reserve]: 8–14. Moscow: VINITI. (In Russian).
- Buga S.V.** 1989b. Dendrocolous aphids of green stands in Central Botanical Gardens of the Academy of Sciences of Byelorussian SSR. In: *Zashchita rasteniy v botanicheskikh sadakh i sohranenie ekologicheskogo ravnovesiya* [Plant protection in botanical gardens and ecological balance maintaining]: 32–33. Riga. (In Russian).
- Buga S.V.** 1989c. *Dendrofil'nye tli – vrediteli zelenykh nasazhdeniy Byelorussii (vidovoy sostav i biologicheskoe obosnovanie zashchitnykh meropriyatiy)* [Dendrocolous aphids as the pests of green stands in Belarus (species composition and biological basis of protection activities)]. Abstract of PhD thesis. 18 p. (In Russian).
- Buga S.V.** 1995. Dendrocolous aphids (Homoptera: Aphidoidea) of Naliboki reserve. *Newsletters of Belarusian University. Ser. 2*, 1: 40–43. (In Russian).
- Buga S.V.** 1997. Dendrocolous aphids (Homoptera: Aphidoidea) of non-park green stands in Minsk. *Newsletters of Belarusian University. Ser. 2*, 3: 36–39. (In Russian).
- Buga S.V. & Jarigo V.E.** 1999a. Current situation and prospects of aphid investigations at Bialowieza Primeval Forest. In: *Belavezhskaya pushcha na rubezhe tret'ego tysyacheletiya* [Bialowieza Primeval Forest in the Third Millennium]: 360. Minsk: Belarusian University Publishers. (In Russian).
- Buga S.V. & Jarigo V.E.** 1999b. Aphids on the trees of the Central Botanical Gardens of Belarusian National Academy of Sciences. *Aphids and other homopterous Insects*, 7: 61–65.
- Buga S.V., Kaminski V.A. & Jarigo V.E.** 1999. Dendrocolous aphids (Insecta: Homoptera: Aphidoidea) of Priluki reserve. In: *Belavezhskaya pushcha na rubezhe tret'ego tysyacheletiya* [Bialowieza Primeval Forest in the Third Millennium]: 358–359. Minsk: Belarusian University Publishers. (In Russian).
- Buga S.V.** 2000. Expansion of new species of aphids on the trees with respect to plant introduction activity in Belarus. *Aphids and Other Homopterous Insects*, 7: 57–60.
- Buga S.V.** 2001. *Dendrofil'nye tli Belorusi* [Dendrocolous aphids of Belarus]. Minsk: Belarusian University Publishers. 98 p. (In Russian).
- Buga S.V.** 2002. *Struktura i ekologicheskie osnovy formirovaniya fauny dendrofil'nykh tley Belarusi* [Taxonomical structure of dendrocolous aphids fauna in Belarus and ecological basis of its formation]. Abstract of Sc.D. thesis. 33 p. (In Russian).
- Buga S.V., Leshchynskaya N.V. & Stekolshchikov A.V.** 2008. Addition to the aphid (Homoptera: Aphidinea) fauna of Belavezha Primeval Forest. *Newsletters of Belarusian University. Ser. 2*, 2: 57–61. (In Russian).
- Buga S.V. & Stekolshchikov A.V.** 2009. Aphids as pests of fruit- and berry-producing plants in Byelorussia. *Redia*, 92: 239–242.
- Buga S.V., Petrov D.L., Rhyzhaya A.V. & Sautkin F.V.** 2010. *Phytophagous pests of woody plants in Minsk and Grodno*. Minsk: Belarusian University Publishers. 40 p. (In Russian).
- Buga S.V. & Stekolshchikov A.V.** 2012. Aphids of the tribe Macrosiphini (Insecta: Homoptera: Aphididae) in Belarus. *Zoosystematica Rossica*, 21(1): 63–96.
- Buga S.V.** 2016. Order Sternorrhyncha. In: **Lukashuk A.O. & Tsinkevich V.A.** (Eds.). *Biologicheskoe raznoobrazie Berezinskogo biosfernogo zapovednika: nogokhvostki (Collembola) i nasekomye (Insecta)* [Biological diversity of Berezinski biosphere reserve: springtails (Collembola) and insects (Insecta)]: 29. Minsk: Belaruski dom pechat. (In Russian).
- Eastop V.F. & Blackman R.L.** 2005. Some new

- synonyms in Aphididae (Hemiptera: Sternorrhyncha). *Zootaxa*, **1089**: 1–36.
- Favret C.** 2016. *Aphid Species File (Version 5.0/5.0)* [online]. Available from: <http://aphid.speciesfile.org/> [updated 21 November 2016; visited 3 December 2016].
- Fiedorowiczówna Z.** 1930. Zooecidia of plants collected in Disna and Braslav districts of Vilno province. *Prace Towarzystwa Przyjaciół Nauk w Wilnie. Wydział nauk matematycznych i przyrodniczych*, **6**: 25–59. (In Polish).
- Gorlenko S.V. & Pan'ko N.A.** 1967. *Vrediteli i bolezni introdutsirovannykh rasteniy* [Pests and diseases of introduced plants]. Minsk: Nauka i tekhnika. 136 p. (In Russian).
- Gorlenko S.V. & Pan'ko N.A.** 1972. *Formirovaniye mikroflory i entomofauny gorodskikh zelenykh nasazdeniy* [Genesis of microflora and entomofauna in urban green stands]. Minsk: Nauka i tekhnika. 168 p. (In Russian).
- Gorlenko S.V., Blintsov A.I & Pan'ko N.A.** 1988. *Ustoychivost' drevesnykh rasteniy k bioticheskim faktoram* [Resistance of woody plants to biotic factors]. Minsk: Nauka i tekhnika. 189 p. (In Russian).
- Hałaj R. & Osiadacz B.** 2013. European gall-forming *Pemphigus* (Aphidoidea: Eriosomatidae). *Zoologischer Anzeiger*, **252**: 417–423.
- Hałaj R., Osiadacz B. & Poljaković-Pajnik L.** 2016. Upper Silesia – a Polish bridgehead gained by *Prociphilus (Meliahizophagus) fraxinifoliae* (Riley, 1879), an alien aphid. *Acta entomologica silesiana*, **24**: 1–9.
- Heie O.E.** 1980. *Aphidoidea (Hemiptera) of Fennoscandia and Denmark. I. The families Mindaridae, Hormaphididae, Thelaxidae, Anoeciidae and Pemphigidae*. Klapenborg: Scandinavian Science Press. 236 p.
- Heie O.E. & Wiegerek P.** 2009. Diagnoses of the higher taxa of Aphidomorpha (Hemiptera: Sternorrhyncha). *Redia*, **92**: 261–269.
- Holman J.** 2009. *Host Plant Catalog of Aphids: Palearctic Region*. Berlin: Springer. 1140 p.
- Jarigo V.E. & Buga S.V.** 1999. Taxonomical structure of coniferous aphids (Insecta: Homoptera: Aphidoidea) in Belarus. In: *Strukturno-funktional'noe sostoyanie biologicheskogo raznoobraziya zhivotnogo mira Belarusi* [Structural and functional state of the animal biological diversity in Belarus]: 357–359. Minsk: Institut zoologii NAN Belarusi. (In Russian).
- Jarigo V.E.** 1999. Pestful coniferous aphids (Insecta: Homoptera: Aphidoidea) in Belarus. In: *Les. Nauka. Molodezh* [Forest. Science. Youth]: 209–211. Gomel. (In Russian).
- Losinskaya N.L.** 1960. The main representatives of tree and shrub pestful entomofauna at the Central Botany Gardens of the BSSR Academy of Sciences. In: *Central'nyy Botanicheskiy Sad Akademii Nauk BSSR: Sbornik nauchnykh rabot* [The Central Botany Gardens of the Belarusian SSR Academy of Science: Miscellaneous of scientific works], **1**: 114–121. (In Russian).
- Mordvilko A.K.** 1924. *The Woolly Apple Aphid (Eriosoma lanigerum Hausmann) and other Eriosomea. Biology and distribution*. Lenigrad: Novaya Derevnya. 109 p. (in Russian).
- Mordvilko A.K.** 1930. The preliminary report on Belarusian aphid (Aphidae) fauna investigations in Belarus in summer 1929. *Matař'yal' da vyvuchen'ya flery i fauny Belarusi*, **5**: 161–164. (In Belarusian).
- Moskvina I.P., Petrov D.L. & Buga S.V.** 2001. The structure of entomophagous communities for the gall-forming aphids common species (family Pemphigidae). In: *Raznoobrazie zhivotnogo mira Belarusi: itogi izucheniya i perspektivy sokhraneniya* [The diversity of fauna of Belarus: study results and conservation perspectives]: 117–119. Minsk. (In Russian).
- Mormyleva V.F.** 1981. Aphids of grain cereals in Belarus and measures of its control. In: *Aukshtikal'nene A.M., Zayanchkauš as A.P., Semyanov V.P. & Yakimavichus A.B.* (Eds). *Noveyshie dostizheniya sel'skokhozyaystvennoy entomologii (Trudy vos'mogo s'ezda Vsevoyuznogo entomologicheskogo obshchestva, Vilnius, 9–13 oktyabrya 1979)* [Newest achievements of agricultural Entomology (Proceedings of the 8th congress of the All-union entomological society, Vilnius, October, 9–13, 1979)]: 119–122. Vilnius. (In Russian).
- Osiadacz B. & Hałaj R.** 2009. The aphids (Hemiptera: Sternorrhyncha: Aphidinea) of Poland. A Distributional Checklist. *Polish Entomological Monographs*, **6**: 1–96.
- Osiadacz B. & Hałaj R.** 2010. Systematic Review of Aphids (Hemiptera: Sternorrhyncha: Aphidomorpha) of Poland with Host Plant Index. A Monographic Survey. *Silesian Natural History Monographs*, **1**: 1–192.
- Osiadacz B. & Hałaj R.** 2012. The Update of Systematic Review of Aphids (Hemiptera: Sternorrhyncha: Aphidomorpha) of Poland with Host Plant Index. *Wiadomości Entomologiczne*, **31**(4): 230–241.

- Osiadacz B. & Halaj R.** 2014. First records of gall-inducing aphid *Pemphigus populi* (Hemiptera: Aphidoidea, Eriosomatidae) in Poland with gall-based key to Central and North European species of the genus. *Entomologica Fennica*, **25**: 16–25.
- Pan'ko N.A.** 1969. On pest fauna of urban green stands in the South of Belarus. In: **Smolsky N.V.** (Ed.). *Introduktsiya rasteniy i okhrana prirody* [Plant introduction and environmental protection]: 204–214. Minsk: Nauka i tekhnika. (In Russian).
- Perepeczko-Baumanowa J.** 1934. Zooecidia collected in Oshmyany and the nearest vicinity. *Prace Towarzystwa Przyjaciół Nauk w Wilnie. Wydział nauk matematycznych i przyrodniczych*, **9**: 239–261. (In Polish).
- Petrov D.L.** 2004. Common gall-forming species in Belarus. *Newsletters of Belarusian University. Ser. 2*, **2**: 63–68. (In Russian).
- Petrov D.L.** 2006. *Gall-forming arthropods of trees in Belarus*. Minsk: BelISA. 140 p. (In Russian).
- Petrov D.L. & Buga S.V.** 2008a. *Gall-forming arthropods as the pests of green stands in Belarus*. Minsk: Belarusian University Publishers. 38 p. (In Russian).
- Petrov D.L. & Buga S.V.** 2008b. *Gall-forming arthropods are the pests of green stands in Belarus*. Minsk: Belarusian University Publishers. 42 p. (In Russian).
- Petrov D.L.** 2013. Entomophagous species of common gall-forming aphids (family Pemphigidae). In: *Zoologicheskie chteniya* [Zoological lectures]: 239–241. Grodno: Grodno University Publishers. (In Russian).
- Petrov D.L.** 2014. The structure of entomophagous community of *Eriosoma ulmi* (L.) on *Ulmus glabra* (Huds.) in green stands of Belarus. *Trudy Belorusskogo gosudarstvennogo Universiteta*, **9**(2): 104–109. (In Russian).
- Rakauskas R. & Buga S.** 2010. Contribution to the knowledge of the aphid (Hemiptera, Sternorrhyncha: Aphidoidea) fauna of the Gorodok Highland, Belarus. *Acta Zoologica Lituanica*, **20** (4): 205–224.
- Remaudière G. & Remaudière M.** 1997. *Catalogue des Aphididae du monde – Catalogue of the world's Aphididae (Homoptera, Aphidoidea)*. Paris: INRA editions. 473 p.
- Rupais A.A.** 1989. *Tli (Aphidodea) Latvii* [The aphids (Aphidodea) of Latvia]. Riga: Zinatne. 331 p. (In Russian).
- Rhyzhaya A.V. & Buga S.V.** 2012. Common coleopteran and homopterous insects which are phytophagous on woody plants in Minsk and Grodno urbocenosis. *Newsletters of Grodno University. Ser. 5*, **2**(131): 120–130. (In Russian).
- Shaposhnikov G.Ch.** 1964. Suborder Aphidinea – Aphids. In: **Bei-Bienko G.Ya.** (Ed.). *Opredelitel' nasekomykh Evropeyskoy chasti SSSR* [Key to the insects of the European part of the USSR], **1**: 489–616. Moscow-Leningrad: Nauka. (In Russian).
- Shaposhnikov G.Ch.** 1985. The main features of the evolution of aphids. In: *Evolution and biosystematics of aphids. Proceedings of the International Aphidological Symposium at Jabłonna, Poland, April 1981*: 19–99. Wrocław: Ossolineum.
- Stekolshchikov A.V. & Buga S.V.** 2006. Notes on *Trichosiphon aphis* (*Xenomysus*) *cortices* Aizenberg, 1935 and *T. (X.) foliotus* Shaposhnikov in Juchnevitch, 1968, nomen nudum (Homoptera: Aphididae). *Zoosystematica Rossica*, **15**(1): 63–68.
- Stekolshchikov A.V., Buga S.V. & Leshchinskaya N.V.** 2008. Addition to the aphid fauna of Byelorussia (Homoptera: Aphidoidea) with detailed description of sexuales of *Semiaphis anthrisci* (Kaltenbach, 1843). *Zoosystematica Rossica*, **17**(1): 61–66.
- Stekolshchikov A.V., Leshchinskaya N.V., Buga S.V. & Voronova N.V.** 2010. Recent additions to the aphid (Homoptera: Aphidoidea) fauna of Byelorussia. *Zoosystematica Rossica*, **19** (1): 35–38.
- Wojciechowski W., Depa Ł., Kanturski M., Wegierek P. & Wieczorek K.** 2015. An annotated checklist of the Aphids (Hemiptera: Aphidomorpha) of Poland. *Polish Journal of Entomology*, **84**(4): 383–420.

ADDENDA

Electronic supplementary material. Text file. File format: PDF. Available from: <http://www.zin.ru/journals/zsr/content.asp?year=2016>

Explanation note. List of species of the family Eriosomatidae in Belarus.

Received 5 Dec. 2016 / Accepted 14 Dec. 2016

Editorial responsibility: I.A. Gavrilov-Zimin