



Agriculture
Canada

Checklist of

BEEETLES


of Canada and Alaska



Checklist of

BEEETLES

of Canada and Alaska



Edited by

Yves Bousquet
Biosystematics Research Centre
Ottawa, Ontario

Research Branch
Agriculture Canada
Publication 1861/E

1991

©Minister of Supply and Services Canada 1991

Available in Canada through

Associated Bookstores
and other booksellers

or by mail from

Canada Communication Group — Publishing
Ottawa, Canada K1A 0S9

Cat. no. A43-1861/1991E

ISBN 0-660-13767-4

Canadian Cataloguing in Publication Data

Checklist of beetles of Canada and Alaska

(Publication ; 1861/E)

Includes bibliographical references.

Cat. no. A43-1861/1991E

ISBN 0-660-13767-4

1. Beetles-Canada-Nomenclature. 2. Beetles-Alaska-Nomenclature. I. Bousquet, Yves. II. Canada. Agriculture Canada. Research Branch. III. Series: Publication (Canada. Agriculture Canada). English ; 1861/E.

QL581.C4 1991 595.76 C91-099101-4

Staff Editor
Frances Smith

CONTENTS

List of contributors	v	Superfamily Scarabaeoidea	143
Acknowledgments	v	Family Lucanidae (J. McNamara)	143
Summary	vi	Family Passalidae (J. McNamara)	145
Résumé	vi	Family Scarabaeidae (J. McNamara)	145
Synoptic table	1	Superfamily Byrrhoidea	158
Introduction	4	Family Byrrhidae (P.J. Johnson)	158
Format	4	Superfamily Buprestoidea	160
Subsequent revisions of the checklist	5	Family Buprestidae (A. Davies)	160
Abbreviations and symbols	5	Superfamily Dryopoidea	169
ORDER COLEOPTERA	6	Family Ptilodactylidae (L. LeSage)	169
Suborder Archostemata	6	Family Heteroceridae (L. LeSage)	169
Family Cupedidae (J.M. Campbell)	6	Family Limnichidae (L. LeSage)	171
Family Micromalthidae (J.M. Campbell)	6	Family Dryopidae (L. LeSage)	171
Suborder Adephaga	7	Family Elmidae (L. LeSage)	172
Family Rhysodidae (Y. Bousquet)	7	Family Psephenidae (L. LeSage)	174
Family Trachypachidae (Y. Bousquet)	7	Superfamily Elateroidea	174
Family Carabidae (Y. Bousquet)	8	Family Artematopidae (Y. Bousquet)	174
Family Haliplidae (R.E. Roughley)	60	Family Elateridae (Y. Bousquet)	175
Family Amphizoidae (Y. Bousquet)	61	Family Throscidae (Y. Bousquet)	186
Family Noteridae (R.E. Roughley)	61	Family Eucnemidae (Y. Bousquet)	186
Family Dytiscidae (D.J. Larson and R.E. Roughley)	62	Superfamily Cantharoidea	188
Family Gyrinidae (R.E. Roughley)	72	Family Lycidae (J. McNamara)	188
Suborder Polyphaga	74	Family Phengodidae (J. McNamara)	190
Superfamily Staphylinoidea	74	Family Lampyridae (J. McNamara)	190
Family Hydraenidae (R.E. Roughley)	74	Family Cantharidae (J. McNamara)	192
Family Ptiliidae (J.M. Campbell)	75	Superfamily Dermestoidea	195
Family Limulodidae (J.M. Campbell)	76	Family Derodontidae (D.E. Bright)	195
Family Agyrtidae (J.M. Campbell)	77	Family Nosodendridae (J.M. Campbell)	196
Family Leiodidae (S.B. Peck)	77	Family Dermestidae (Y. Bousquet)	196
Family Scydmaenidae (A. Davies)	82	Superfamily Bostrichoidea	199
Family Micropeplidae (J.M. Campbell)	84	Family Bostrichidae (J. McNamara)	199
Family Silphidae (J.M. Campbell)	84	Family Lyctidae (J. McNamara)	201
Family Staphylinidae (J.M. Campbell and A. Davies)	86	Family Anobiidae (J. McNamara)	201
Family Scaphidiidae (J.M. Campbell)	124	Family Ptinidae (Y. Bousquet)	205
Family Pselaphidae (A. Davies)	125	Superfamily Lymexyloidea	206
Superfamily Hydrophiloidea	130	Family Lymexylidae (J.M. Campbell)	206
Family Hydrophilidae (R.E. Roughley)	130	Superfamily Cleroidea	206
Family Georyssidae (R.E. Roughley)	135	Family Trogossitidae (Y. Bousquet)	206
Family Sphaeritidae (R.E. Roughley)	135	Family Cleridae (J. McNamara)	208
Family Histeridae (A. Davies)	135	Family Melyridae (D.E. Bright)	211
Superfamily Eucinetidae	141	Superfamily Cucujoidea	213
Family Eucinetidae (J.M. Campbell)	141	Family Sphindidae (J.M. Campbell)	213
Family Clambidae (J.M. Campbell)	141	Family Nitidulidae (J. McNamara)	214
Family Scirtidae (J.M. Campbell)	142	Family Rhizophagidae (Y. Bousquet)	218
Superfamily Dascilloidea	143	Family Cucujidae (Y. Bousquet)	219
Family Dascillidae (J.M. Campbell)	143	Family Cryptophagidae (Y. Bousquet)	221
Family Rhipiceridae (J.M. Campbell)	143	Family Languriidae (J.M. Campbell)	223
		Family Erotylidae (J.M. Campbell)	225
		Family Phalacridae (J.M. Campbell)	226

Family Cerylonidae (J.M. Campbell)	227	Family Anthicidae (Y. Bousquet)	271
Family Corylophidae (J.M. Campbell)	228	Family Pedilidae (Y. Bousquet)	274
Family Coccinellidae (J. McNamara)	229	Family Aderidae (Y. Bousquet)	275
Family Endomychidae (J.M. Campbell)	237	Family Scraptiidae (L. LeSage)	275
Family Lathridiidae (Y. Bousquet)	239	Superfamily Chrysomeloidea	277
Family Biphyllidae (J.M. Campbell)	241	Family Cerambycidae (J. McNamara)	277
Family Byturidae (J.M. Campbell)	241	Family Bruchidae (J. McNamara)	300
Superfamily Tenebrionoidea	242	Family Chrysomelidae (L. LeSage)	301
Family Mycetophagidae (Y. Bousquet)	242	Superfamily Curculionoidea	323
Family Tetratomidae (L. LeSage)	243	Family Nemonychidae (J. McNamara)	323
Family Ciidae (J. McNamara)	244	Family Anthribidae (J. McNamara)	324
Family Melandryidae (L. LeSage)	245	Family Ithyceridae (J. McNamara)	325
Family Mordellidae (J. McNamara)	248	Family Attelabidae (J. McNamara)	326
Family Rhipiphoridae (J.M. Campbell)	250	Family Rhynchitidae (J. McNamara)	326
Family Colydiidae (Y. Bousquet)	251	Family Brentidae (J. McNamara)	327
Family Zopheridae (J.M. Campbell)	252	Family Apionidae (J. McNamara)	328
Family Tenebrionidae (Y. Bousquet and J.M. Campbell)	253	Family Curculionidae (J. McNamara)	329
Family Lagriidae (J.M. Campbell)	261	Family Scolytidae (J. McNamara)	356
Family Cephaloidae (J.M. Campbell)	262	Family Platypodidae (J. McNamara)	365
Family Meloidae (J.M. Campbell)	263	ORDER STREPSIPTERA	366
Family Oedemeridae (J.M. Campbell)	266	Suborder Stylopida	366
Family Prostomidae (J.M. Campbell)	267	Family Corioxenidae (S.B. Peck)	366
Family Synchronidae (L. LeSage)	267	Family Halictophagidae (S.B. Peck)	366
Family Mycteridae (J.M. Campbell)	267	Family Stylopidae (S.B. Peck)	367
Family Boridae (J.M. Campbell)	268	References	368
Family Pythidae (J.M. Campbell)	268	Index	397
Family Pyrochroidae (J.M. Campbell)	269		
Family Salpingidae (J.M. Campbell)	270		

LIST OF CONTRIBUTORS

- Y. Bousquet, Biosystematics Research Centre,
Agriculture Canada, Ottawa, Ontario K1A 0C6
- D.E. Bright, Biosystematics Research Centre,
Agriculture Canada, Ottawa, Ontario K1A 0C6
- J.M. Campbell, Biosystematics Research Centre,
Agriculture Canada, Ottawa, Ontario K1A 0C6
- A. Davies, Biosystematics Research Centre,
Agriculture Canada, Ottawa, Ontario K1A 0C6
- P.J. Johnson, Department of Entomology,
University of Wisconsin, Madison, Wisconsin
53706
- D.J. Larson, Department of Biology, Memorial
University of Newfoundland, St. John's,
Newfoundland A1B 3X9
- L. LeSage, Biosystematics Research Centre,
Agriculture Canada, Ottawa, Ontario K1A 0C6
- J. McNamara, Biosystematics Research Centre,
Agriculture Canada, Ottawa, Ontario K1A 0C6
- S.B. Peck, Department of Biology, Carleton
University, Ottawa, Ontario K1S 5B6
- R.E. Roughley, Department of Entomology,
University of Manitoba, Winnipeg, Manitoba
R3T 2N2

ACKNOWLEDGMENTS

We wish to thank the following persons for reviewing sections of the checklist: R.L. Aalbu (Tenebrionidae); R.S. Anderson (Silphidae, Curculionidae); F.G. Andrews (Lathridiidae); I.S. Askevold (Chrysomelidae); F.M. Atton (Gyrinidae); E.U. Balsbaugh (Chrysomelidae); R. Baranowski (Leiodidae); W.F. Barr (Cleridae); E.C. Becker (Elateroidea); H.P. Brown (Dryopoidea); D. Chandler (Anthicidae, Pselaphidae, Melandryidae, Tetratomidae, Synchronidae, Scraptiidae); J. Cicero (Lampyridae); W. Connell (Nitidulidae); M. Daccordi (Chrysomelinae); J.G. Edwards (Synetinae); D.G. Furth (Chrysomelidae); R.E. Gordon (Coccinellidae); L.H. Herman (Staphylinidae); A.T. Howden (Curculionidae); H. Howden (Scarabaeidae); C. Johnson (Ptiliidae); J.B. Karren (Chlamisinae); J.M. Kingsolver (Bruchidae); J.L. Krysan (Chrysomelidae); S. Laplante (Cleridae, Cerambycidae); J.F. Lawrence (Melandryidae, Tetratomidae, Synchronidae, Scraptiidae); I. Lobl (Scaphiidae); W.V. Miller (Heteroceridae); A.R.

Moldenke (Clytrinae); H.G. Nelson (Dryopoidea); A.F. Newton Jr. (Leiodidae, Staphylinidae); S. Oygur (Gyrinidae); J. Pakaluk (Corylophidae); J.D. Pinto (Meloidae); D. Pollock (Mycteridae, Boridae, Pythidae, Salpingidae); V. Puthz (Steninae); E.G. Riley (Cassidinae); W.T. Schultz (Eumolpinae); T.N. Seeno (Chrysomelidae); R.B. Selander (Meloidae); S.A. Slipinski (Cerylonidae); C.L. Staines (Hispidinae); W.E. Steiner Jr. (Phalacridae); J.B. Stribling (Ptilodactylidae); M. Thayer (Staphylinidae); M.C. Thomas (Cucujidae, Prostomidae); C.A. Triplehorn (Tenebrionidae); R.E. White (Anobiidae, Chrysomelidae); D.P. Wooldridge (Limnichidae); D.K. Young (Scirtidae, Pyrochroidae); P. Zwick (Leiodidae).

Thanks are also extended to B.F. Carr, J.L. Carr, and D.W. Langor, for providing several provincial records; to D. Beauchamp, B. Walter, L. Robillard, and R. Hutchinson, for typing the checklist; and to A. Smetana, for checking the termination of all species-group names.

SUMMARY

All species and subspecies of beetles known to occur in Canada and Alaska are listed along with their synonyms; these names are arranged alphabetically under the respective genus-group names. Valid genus-group and family-group names are listed in phylogenetic order. No new taxa, combinations, or synonyms are proposed. Species distributions are presented in tabular form, indicating the

presence or absence of the taxa in each province and territory, in Labrador, and in Alaska. Introduced and holarctic species are highlighted. References to pertinent identification keys are indicated under the corresponding supraspecific taxon. The work also includes a brief introduction for every family treated and an index to all supraspecific names.

RÉSUMÉ

L'ouvrage répertorie toutes les espèces et sous-espèces de Coléoptères connues du Canada et de l'Alaska ainsi que leurs synonymes; ces noms sont énumérés par ordre alphabétique à la suite du nom du groupe-genre correspondant. Les noms valides des groupe-genre et groupe-famille sont présentés par ordre phylogénétique. Cette liste ne propose aucun nouveau taxon, combinaison ou synonyme. La répartition des espèces est présentée sous forme

tabulaire, indiquant la présence ou l'absence des taxa dans chaque province et territoire, le Labrador et l'Alaska. On met en relief les espèces introduites ou holarctiques et on mentionne, sous le taxon supraspécifique correspondant, les références contenant des tableaux de détermination utiles. L'ouvrage comprend aussi une courte introduction sur chaque famille considérée ainsi qu'un index des noms supraspécifiques.

Synoptic table

Family	AK	YK	NT	BC	AB	SK	MB	ON	PQ	NB	NS	PE	LB	NF	T	H	I
Cupedidae	-	-	-	1	-	-	-	2	1	-	-	-	-	-	3	-	-
Micromalthidae	-	-	-	1	-	-	-	-	-	-	-	-	-	-	1	-	-
Rhysodidae	-	-	-	1	-	-	-	1	-	-	-	-	-	-	2	-	-
Trachypachidae	1	1	1	2	1	1	-	-	-	-	-	-	-	-	2	-	-
Carabidae	237	164	207	483	386	314	350	503	451	243	244	137	94	175	946	84	46
Halipidae	3	4	9	23	15	10	13	17	18	3	3	-	2	4	37	1	-
Amphizoidae	1	2	-	3	2	-	-	-	-	-	-	-	-	-	3	-	-
Noteridae	-	-	-	-	-	-	-	2	1	-	-	-	-	-	2	-	-
Dytiscidae	92	111	118	167	153	124	145	155	147	82	77	31	74	78	262	45	-
Gyrinidae	7	10	14	18	21	23	22	27	30	14	10	5	8	10	36	4	-
Hydraenidae	-	-	4	21	10	4	5	5	7	3	1	-	-	1	27	-	-
Ptilidae	5	-	-	16	-	-	1	2	10	5	1	-	-	-	32	1	11
Limulodidae	-	-	-	-	-	-	-	1	-	-	-	-	-	-	1	-	-
Agyrtidae	4	-	1	6	3	-	-	1	-	-	-	-	-	-	8	-	-
Leioidae	36	18	16	84	39	18	26	57	53	19	15	-	7	10	128	6	1
Scydmaenidae	5	-	-	12	1	1	3	29	14	8	6	-	1	-	48	-	-
Micropeplidae	4	1	1	8	4	2	1	5	4	-	-	-	-	-	10	1	-
Silphidae	8	6	9	13	17	18	18	18	15	10	11	7	1	6	25	4	-
Staphylinidae	308	172	155	581	265	126	278	441	460	167	195	19	63	171	1129	88	99
Scaphidiidae	1	1	1	4	1	2	3	13	10	-	-	-	-	1	17	-	-
Pselaphidae	10	1	3	23	16	9	16	56	38	15	11	2	1	3	85	-	2
Hydrophilidae	24	20	36	81	65	52	64	92	86	38	38	2	5	24	149	11	18
Georyssidae	-	-	-	1	1	-	-	-	1	-	-	-	-	-	1	-	-
Sphaeritidae	1	-	-	1	1	-	-	-	-	-	-	-	-	-	1	-	-
Histeridae	1	2	2	57	42	25	43	65	58	13	12	3	-	7	118	1	10
Eucinetidae	-	1	2	5	3	3	4	4	4	2	2	-	-	-	7	-	1
Clambidae	1	-	-	3	3	-	-	1	6	-	-	-	-	-	7	1	1
Scirtidae	1	1	2	10	4	3	3	15	15	7	3	-	-	5	26	-	-
Dascillidae	-	-	-	1	-	-	-	-	-	-	-	-	-	-	1	-	-
Rhipiceridae	-	-	-	-	-	-	-	1	-	-	-	-	-	-	1	-	-
Lucanidae	-	1	2	7	2	2	3	8	5	2	3	-	-	1	14	-	-
Passalidae	-	-	-	-	-	-	1	1	-	-	-	-	-	-	1	-	-
Scarabaeidae	13	9	17	88	96	58	85	136	109	41	47	11	7	20	248	1	21
Byrrhidae	14	9	13	15	12	5	7	10	12	2	4	-	5	4	26	3	2
Buprestidae	11	13	18	75	57	41	63	99	80	36	22	7	4	5	152	1	2
Ptilodactylidae	-	-	-	1	-	-	1	2	3	-	-	1	-	-	3	-	-
Heteroceridae	2	4	5	9	16	14	19	16	16	5	-	1	-	-	25	-	-
Limnichidae	-	-	-	1	-	-	1	4	2	-	-	-	-	-	4	-	-
Dryopidae	-	-	-	2	1	1	1	2	5	2	2	1	-	1	6	-	1
Elmidae	-	1	1	14	11	4	7	16	21	10	11	5	2	5	32	-	-
Psephenidae	-	-	-	-	-	-	-	3	2	1	2	-	-	-	3	-	-
Artematopidae	-	-	1	3	1	-	1	2	2	1	-	-	-	-	4	-	-
Elateridae	53	32	49	194	131	88	99	175	165	81	87	21	25	46	369	6	5

Synopsis table (continued)

Family	AK	YK	NT	BC	AB	SK	MB	ON	PQ	NB	NS	PE	LB	NF	T	H	I
Throscidae	-	-	2	6	1	-	1	5	5	2	2	-	-	1	9	-	-
Eucnemidae	1	-	-	13	2	1	3	18	20	5	3	-	-	1	32	-	-
Lycidae	3	1	1	8	5	4	8	25	19	5	5	-	1	1	29	-	-
Phengodidae	-	-	-	-	-	-	-	1	-	-	-	-	-	-	1	-	-
Lampyridae	-	-	3	9	3	5	12	19	14	8	10	1	-	3	29	-	1
Cantharidae	15	8	13	57	20	9	22	48	48	22	25	1	5	16	126	-	2
Derodontidae	2	-	-	5	3	-	-	2	1	2	3	-	-	1	8	-	1
Nosodendridae	-	-	-	1	-	-	-	1	-	-	-	-	-	-	2	-	-
Dermestidae	6	3	6	31	16	10	14	24	19	4	5	4	-	4	48	-	17
Bostrichidae	1	1	3	8	3	2	6	13	6	1	1	-	-	-	18	1	1
Lyctidae	-	-	-	4	-	-	1	4	3	1	2	-	-	-	6	-	2
Anobiidae	4	6	2	35	18	9	21	42	40	8	7	-	-	7	79	-	5
Ptinidae	1	-	-	13	5	6	7	11	11	9	10	1	-	5	16	-	13
Lymexylidae	-	-	-	-	1	-	-	1	1	1	1	-	-	1	1	-	-
Trogossitidae	4	4	5	17	6	6	6	12	9	2	2	1	1	3	22	2	-
Cleridae	5	4	5	22	12	10	14	32	24	11	4	-	-	4	50	-	5
Melyridae	-	2	3	35	16	5	6	10	10	1	1	-	-	1	49	-	2
Sphindidae	-	-	-	2	1	-	-	3	4	-	-	-	-	-	5	-	-
Nitidulidae	17	10	8	63	44	16	45	56	63	13	15	3	1	13	101	3	7
Rhizophagidae	3	1	-	15	7	3	4	15	11	4	3	-	-	5	24	-	3
Cucujidae	3	1	2	15	10	9	17	26	24	5	4	1	1	5	31	1	10
Cryptophagidae	26	1	3	47	15	7	12	19	17	2	3	-	2	3	66	11	8
Languriidae	-	-	-	1	-	1	2	6	5	-	-	-	-	-	7	-	1
Erotylidae	-	-	-	5	6	4	5	14	14	2	1	-	-	-	20	-	-
Phalacridae	-	-	-	3	-	-	1	3	2	-	-	-	-	-	5	-	-
Cerylonidae	1	-	-	3	2	-	1	6	3	1	2	-	-	-	7	-	1
Corylophidae	-	1	2	7	4	2	1	4	3	1	-	-	-	-	11	2	2
Coccinellidae	26	31	35	94	83	70	56	83	77	37	25	6	9	13	162	2	8
Endomychidae	-	-	1	7	2	2	5	10	8	2	6	1	-	2	15	-	2
Lathridiidae	19	5	6	34	9	8	17	22	22	6	8	1	1	4	55	6	15
Biphylidae	-	-	-	-	-	-	-	1	1	1	-	-	-	-	1	-	-
Byturidae	1	1	1	1	1	1	1	1	1	1	1	1	-	1	1	-	1
Mycetophagidae	-	-	-	6	4	2	7	10	12	2	1	-	-	-	15	-	-
Tetratomidae	1	1	-	2	1	1	-	5	7	1	-	-	-	1	9	-	1
Ciidae	6	-	6	19	10	5	14	16	17	5	6	-	-	6	29	1	1
Melandryidae	4	2	3	27	10	5	9	27	42	11	10	-	3	13	55	3	2
Mordellidae	-	1	6	18	10	12	37	49	29	14	7	1	-	-	70	-	-
Rhipiphoridae	-	-	-	3	2	2	-	5	2	-	-	-	-	-	10	-	-
Colydiidae	-	-	2	10	3	1	1	10	5	1	-	-	-	1	21	-	1
Zopheridae	-	-	-	1	1	-	-	1	1	1	-	-	-	1	1	-	-
Tenebrionidae	3	3	8	69	32	22	32	70	63	16	14	5	-	7	135	1	11
Lagriidae	-	-	1	1	1	-	3	5	5	1	1	-	-	-	6	-	-
Cephaloidea	1	1	2	5	3	-	1	4	5	1	-	1	2	2	9	-	-

Synoptic table (concluded)

Family	AK	YK	NT	BC	AB	SK	MB	ON	PQ	NB	NS	PE	LB	NF	T	H	I
Meloidae	3	1	3	25	26	23	20	17	12	3	3	-	-	-	49	-	-
Oedemeridae	2	-	-	10	4	-	3	5	5	1	2	-	-	1	13	-	1
Prostomidae	-	-	-	1	-	-	-	-	-	-	-	-	-	-	1	-	-
Synchroidea	-	-	-	-	-	-	1	1	1	-	-	-	-	-	1	-	-
Mycteridae	-	-	-	2	-	-	-	2	2	-	-	-	-	-	4	-	-
Boridae	-	-	1	1	2	2	2	2	2	1	-	-	-	-	2	-	-
Pythidae	3	3	3	5	4	3	4	4	4	3	4	-	1	2	6	-	-
Pyrochroidae	2	-	1	5	4	1	3	7	7	3	4	-	-	1	12	-	-
Salpingidae	3	1	1	7	1	1	1	3	2	1	1	-	1	2	10	-	1
Anthicidae	2	2	5	29	25	24	19	25	21	9	5	1	-	3	55	-	3
Pedilidae	-	-	-	8	4	1	2	6	9	1	1	-	-	-	19	-	-
Aderidae	-	-	-	1	-	-	-	8	8	1	1	-	-	-	9	-	1
Scaptiidae	2	1	-	16	5	1	1	8	10	3	2	-	-	3	21	-	-
Cerambycidae	31	27	32	145	88	56	100	209	178	68	51	11	13	24	354	4	1
Bruchidae	1	1	2	10	8	7	9	11	5	2	2	1	-	1	19	-	6
Chrysomelidae	42	41	72	181	217	174	254	363	274	116	103	19	14	34	569	9	38
Nemonychidae	-	-	-	6	3	-	1	2	3	1	-	-	-	-	8	-	-
Anthribidae	2	2	2	4	8	6	7	12	12	1	1	-	-	1	18	1	1
Ithyceridae	-	-	-	-	-	-	-	1	1	-	-	-	-	-	1	-	-
Attelabidae	-	-	-	-	-	1	2	3	4	1	2	-	-	-	4	-	-
Rhynchitidae	-	-	3	5	6	5	6	7	5	3	3	2	-	2	13	-	-
Brentidae	-	-	-	-	-	-	-	1	1	-	-	-	-	-	1	-	-
Apionidae	3	3	3	12	20	13	15	17	13	2	2	-	-	1	35	-	1
Curculionidae	61	51	55	261	206	124	179	312	280	72	91	17	11	68	609	14	69
Scolytidae	52	33	31	134	80	40	47	89	81	53	46	8	6	27	204	-	5
Platypodidae	-	-	-	1	-	-	-	-	-	-	-	-	-	-	1	-	-
STREPSIPTERA	-	-	-	2	1	3	-	2	2	-	1	-	-	-	11	-	-
TOTAL	1205	838	1031	3628	2464	1673	2351	3843	3466	1365	1320	340	373	872	7447	316	469

INTRODUCTION

Beetles represent the most diverse group of living organisms on earth. More than 250 000 species have been described, and the actual number may easily exceed a million. Given such staggering figures, it is not surprising that Coleoptera can be found nearly everywhere, from the tropics to the Arctic and from caves to the summit of high mountains. Coleoptera species are particularly abundant on vegetation, inside rotten wood, under the bark of dead trees, and in soil, ground litter, fungi, dung, carrion, rodent burrows, and nests. Several groups are adapted to life in fresh or brackish water. Although some dytiscids can venture into salt water and some species do complete all stages of their life cycle in the intertidal zone where they are periodically immersed, no beetles have become completely adapted to marine life. Beetles are important to humans because many species are beneficial in the biological control of pest arthropods and weeds or are major pests of stored products, fabrics, forests, wood products, and crops.

No comprehensive checklist of Coleoptera is available for Canada and Alaska. Leng's catalogue (1920) and its supplements (Leng and Mutchler 1927, 1933; Blackwelder 1939a; Blackwelder and Blackwelder 1948) as well as the world-based *Coleopterorum Catalogus* list most of the species that occur in Canada and Alaska. However, these publications are out of date for many groups, and their distribution data are incomplete. Beaulne (1913-1929) started a checklist of the beetles of Canada but never completed it, and as pointed out by Lindrøth (1966) for the ground beetles, many of Beaulne's distributional records are inaccurate.

This checklist is intended to provide the correct names of the beetles that occur in Canada and Alaska, give a brief survey of the distribution of the taxa in the area covered, and lead the users to the proper publications for the identification of the taxa. The checklist of Strepsiptera has been added at the end simply because the group has often been included in the Coleoptera.

FORMAT

With some exceptions reflecting the author's personal perspective, the concept of the families used in this checklist follows that of Lawrence (1982a). Major changes concern the trachypachids, limulodids, scaphidiids, lyctids, lagriids, pedilids, rhynchitids, scolytids, and platypodids, which are treated as distinct families, and the leptinids, trogids, and geotrupids, which are included respectively within the Leiodidae and the Scarabaeidae for the last two groups. The superfamily concepts and the listing of the families follow Lawrence (1982a).

The classification and information for each family are the responsibility of the respective authors. No new taxa, combinations, or synonyms are proposed. Variety and form names published before 1961 are given subspecific rank unless they were deemed to be synonyms or their status has been changed (article 45 (g) of the International Code of Zoological Nomenclature, 3rd ed.). Varieties and forms published after 1960 are omitted. New distributional records are listed but not identified as such.

The authors of species-group names are listed in parentheses when the original combination differs from the one used in the checklist. To avoid confusion, initials are included for several authors, such as C.R. Sahlberg, R.F. Sahlberg, J.R. Sahlberg, J. Müller, O.F. Müller, W. Horn, J.E. LeConte, and O. Fabricius. LeConte, Horn, and Fabricius stand for

J.L. LeConte, G. Horn, and J.C. Fabricius respectively. An asterisk [*] placed after a species-group name indicates that the taxon is Holarctic, a dagger [†] that it is introduced in North America, and a double dagger [‡] that it is *probably* introduced in North America.

All synonyms based on material collected in North America north of Mexico should be included and are listed in italics. Synonyms of Holarctic and introduced species based on material from outside North America have been included occasionally, particularly when the name has been in use in North American literature. Misidentifications, spelling errors, and invalid emendations are not listed. The ending of all species-group names has been checked to conform with the gender of their respective genera.

Valid genus-group names are in capital letters, their synonyms in italics. All synonyms of genus-group names whose type species are indigenous to North America north of Mexico should be included. Some of the synonyms whose type species are not indigenous to the continent are included also when the name has been used in North American literature.

References to keys are indicated under the corresponding supraspecific taxon. It should be noted that only references including keys useful for the determination of all or most Canadian taxa of the group treated are considered. Keys published

before the 1930s are usually out of date, and most of them are therefore omitted.

An abbreviation of two letters is used to indicate Canadian provinces and Alaskan records (see section entitled "Abbreviations and symbols"). Labrador is considered to be a distinct geographic entity. The distributional citation in the checklist is from west to east.

The distributional records are based on published records and on specimens in the Canadian National Collection. For some families, records

from other collections have been included. Doubtful records from the literature that apparently are based on misidentifications or mislabeled specimens are rejected.

A question mark (?) inserted in a geographic column indicates that the taxon was recorded in the area, but the record is questionable. The absence of a geographic entry after a species-group name indicates that the taxon was recorded in Canada, but the province was not specified.

SUBSEQUENT REVISIONS OF THE CHECKLIST

This publication is set as a data base that can be easily updated to maintain an accurate summary of our current knowledge of the beetles of Canada and Alaska. The authors intend to maintain a corrected version of the checklist and to republish it whenever we have accumulated enough changes. The authors would appreciate the assistance of all users of this checklist in calling errors and omis-

sions to their attention. We would also welcome receiving specimens that represent new provincial or Alaskan records. Specialists and other users may send material to the Biosystematics Research Centre for confirmation, and with the sender's permission, for incorporation as voucher specimens in the Canadian National Collection.

ABBREVIATIONS AND SYMBOLS

AB Alberta
AK Alaska
BC British Columbia
LB Labrador
MB Manitoba
NB New Brunswick
NF Newfoundland
NS Nova Scotia

NT Northwest Territories
ON Ontario
PE Prince Edward Island
PQ Quebec
SK Saskatchewan
YK Yukon Territory
H Holarctic
I Introduced in North America

* Holarctic
† Introduced in North America

‡ Probably introduced in North America
§ No subgeneric name available

Family CORYLOPHIDAE minute fungus beetles

J.M. Campbell

Members of Corylophidae, a small family of about 435 species, are found in all major faunal regions, but only 11 species are known to occur in Canada. Adults are usually collected in moldy leaf litter and piles of grass cuttings. Adults and larvae are thought to feed on fungal spores.

The taxonomy of the Canadian corylophid species is poorly known; reliable species identification is not possible at this time, and a number of

species have yet to be discovered. Casey (1900) prepared the last revision covering the North American species of Corylophidae. The classification adopted here is based on Paulian's (1950) reclassification of the Old World taxa.

YK (1); NT (2); BC (7); AB (4); SK (2); MB (1); ON (4); PQ (3); NB (1) • I (2)

Subfamily CORYLOPHINAE Tribe Orthoperini

Genus ORTHOPERUS Stephens

<i>O. atomus</i> (Gyllenhal)†	-	-	-	BC	-	-	-	-	-	-	-	-	-	-
<i>O. scutellaris</i> LeConte	-	-	NT	BC	AB	SK	-	-	-	-	-	-	-	-
<i>lucidus</i> Casey														
<i>O. suturalis</i> LeConte	-	-	-	-	-	-	-	ON	-	-	-	-	-	-

Tribe Rypobiini

Genus GRONEVUS Casey

<i>G. fuscicornis</i> Casey	-	-	-	-	-	-	-	ON	PQ	-	-	-	-	-
-----------------------------	---	---	---	---	---	---	---	----	----	---	---	---	---	---

Subfamily SERICODERINAE Tribe Sericoderini

Genus SERICODERUS Stephens

<i>S. lateralis</i> (Gyllenhal)†	-	-	-	BC	-	-	-	-	PQ	-	-	-	-	-
----------------------------------	---	---	---	----	---	---	---	---	----	---	---	---	---	---

Subfamily SACIINAE Tribe Saciini

Genus MOLAMBA Casey

<i>M. biguttata</i> (LeConte)	-	-	-	BC	AB	-	-	-	-	-	-	-	-	-
<i>M. fasciata</i> (Say)	-	-	-	-	-	-	-	ON	PQ	NB	-	-	-	-
<i>subfasciata</i> (Guérin-Méneville)														
<i>M. obesa</i> Casey	-	-	-	BC	-	-	-	-	-	-	-	-	-	-

Genus SACIUM LeConte

Parmulus Harold

<i>S. lugubre</i> LeConte	-	YK	NT	BC	AB	SK	MB	ON	-	-	-	-	-	-
<i>S. montanum</i> Casey	-	-	-	BC	-	-	-	-	-	-	-	-	-	-

Genus ARTHROLIPS Wollaston

<i>A. decolor</i> (LeConte)														
<i>californicus</i> (Matthews)	-	-	-	-	AB	-	-	-	-	-	-	-	-	-

Family COCCINELLIDAE ladybird beetles

J. McNamara

Coccinellidae comprises 162 species and subspecies in Canada and Alaska. Many species are introduced in North America, some of these purposely for biological control. Most coccinellid adults and larvae are beneficial, because they are predators of aphids, scale insects, and mites. However, in the absence of preferred food, many species feed on the immature stages of other insects. The species of *Psyllobora* feed on fungi, and the only species of *Epilachna* that occurs in Canada feeds on plants of the family Leguminaceae.

The taxonomy of the North American Coccinellidae is well known, thanks to the recent

monograph by Gordon (1985). Only some of the most recent references on coccinellids have been listed in the checklist. Good regional works on Coccinellidae include those of Chapin (1974) for Louisiana, Belicek (1976) for western Canada and Alaska, and Laroche (1979) for Quebec.

AK (26); YK (31); NT (35); BC (94); AB (83); SK (70); MB (56); ON (83); PQ (77); NB (37); NS (25); PE (6); LB (9); NF (13) • H (2); I (8)

Subfamily STICHOLOTIDINAE Tribe Microweisini

Genus MICROWEISEA Cockerell

Epismilia Cockerell

Pseudoweisea Schwarz

Smilia Weise

Key to species: Gordon (1970a)

M. misella (LeConte)	-	-	-	BC	AB	SK	MB	ON	PQ	NB	-	-	-	NF
----------------------	---	---	---	----	----	----	----	----	----	----	---	---	---	----

Genus COCCIDOPHILUS Brethes

Cryptoweisea Gordon

Key to North American species: Gordon (1970a)

C. marginatus (LeConte)	-	-	-	BC	AB	SK	MB	ON	PQ	NB	-	-	-	-
-------------------------	---	---	---	----	----	----	----	----	----	----	---	---	---	---

Tribe Serangiini

Genus DELPHASTUS Casey

Key to species: Gordon (1970b)

D. pusillus (LeConte)	-	-	-	-	-	-	-	ON	PQ	-	-	-	-	-
<i>puncticollis</i> (LeConte)														
<i>sonoricus</i> Casey														

Subfamily SCYMNINAE Tribe Stethorini

Genus STETHORUS Weise

Key to New World species: Gordon and Chapin (1983)

S. punctillum Weise†	-	-	-	BC	-	-	-	ON	-	-	-	-	-	-
S. punctum punctum (LeConte)	-	-	-	-	AB	SK	-	ON	PQ	-	NS	-	-	-
S. punctum picipes Casey	-	-	-	BC	-	-	-	-	-	-	-	-	-	-
<i>brevis</i> Casey														

Tribe Scymnini

Key to North American species: Gordon (1976)

Genus DIDION Casey

D. longulum Casey	-	YK	NT	BC	AB	SK	-	-	PQ	-	-	-	-	-
<i>occiduum</i> (Casey)														
<i>parviceps</i> Casey														
D. nanum (LeConte)	-	-	-	-	-	-	-	ON	PQ	-	-	-	-	-
D. punctatum (Melsheimer)	-	-	NT	BC	AB	SK	MB	ON	PQ	NB	-	PE	-	NF

Coccinellidae

Genus SCYMNUS Kugelann

(Subgenus SCYMNUS s. str.)

<i>S. americanus</i> Mulsant	-	-	-	-	-	-	-	ON	PQ	-	-	-	-	-
<i>S. apicanus pseudapicanus</i> Gordon	-	-	-	-	AB	SK	MB	-	-	-	-	-	-	-
<i>borealis</i> Gordon														
<i>S. caurinus</i> Horn	-	-	-	BC	AB	-	-	-	-	-	-	-	-	-
<i>aluticollis</i> Casey														
<i>S. fenderi</i> Malkin	-	-	-	BC	-	-	-	-	-	-	-	-	-	-
<i>S. indianensis</i> Weise	-	-	-	-	-	-	-	ON	-	-	-	-	-	-
<i>rusticus</i> Casey														
<i>S. nebulosus</i> LeConte	-	-	-	BC	-	-	-	-	-	-	-	-	-	-
<i>harneyi</i> Hatch														
<i>infuscatus</i> Boheman														
<i>phelpsii</i> Crotch														
<i>S. opaculus</i> Horn	-	-	-	BC	AB	SK	-	-	-	-	-	-	-	-
<i>S. paracanus linearis</i> Gordon	-	-	-	-	AB	SK	MB	-	-	-	-	-	-	-
(Subgenus PULLUS Mulsant)														
<i>S. abbreviatus</i> LeConte	-	-	-	-	-	-	-	ON	PQ	-	-	-	-	-
<i>S. aquilonarius</i> Gordon	-	-	-	-	AB	SK	-	-	-	-	-	-	-	-
<i>S. ardelio</i> Horn	-	-	-	BC	AB	-	-	-	-	-	-	-	-	-
<i>apacheanus</i> Casey														
<i>decipiens</i> Casey														
<i>sanctus</i> Casey														
<i>S. aridus</i> Casey	-	-	-	BC	-	-	-	-	-	-	-	-	-	-
<i>S. brullei</i> Mulsant	-	-	-	-	-	SK	MB	ON	PQ	-	NS	-	-	-
<i>divisus</i> Casey														
<i>laurenticus</i> Casey														
<i>lodi</i> Stehr														
<i>subaeneus</i> Casey														
<i>S. calaveras</i> Casey	-	-	-	BC	-	-	-	-	-	-	-	-	-	-
<i>saginitus</i> Casey														
<i>strenuus</i> Casey														
<i>stygius</i> Casey														
<i>tenuivestis</i> Casey														
<i>S. carri</i> Gordon	-	-	-	-	AB	SK	-	-	-	-	-	-	-	-
<i>S. caudalis</i> LeConte	-	-	-	-	-	SK	MB	ON	PQ	-	-	-	-	-
<i>agricola</i> Casey														
<i>indutus</i> Casey														
<i>natchezianus</i> Casey														
<i>S. cervicalis</i> Mulsant	-	-	-	-	-	-	-	ON	-	-	-	-	-	-
<i>S. compar</i> Casey	-	-	-	-	-	-	-	ON	-	-	-	-	-	-
<i>impunctus</i> Wingo														
<i>vicksburgicus</i> Casey														
<i>S. coniferarum</i> Crotch	-	-	-	BC	-	-	-	-	-	-	-	-	-	-
<i>S. fraternus</i> LeConte	-	-	-	-	-	-	-	ON	PQ	-	-	-	-	-
<i>dentipes</i> Fall														
<i>haemorrhous</i> LeConte														
<i>S. humboldti</i> Casey	-	-	-	BC	-	-	-	-	-	-	-	-	-	-
<i>S. impexus</i> Mulsant†	-	-	-	BC	-	-	-	-	-	NB	NS	-	-	-
<i>S. impletus</i> Gordon	-	-	-	BC	-	-	-	-	-	-	-	-	-	-
<i>S. iowensis</i> Casey	-	-	-	BC	-	SK	MB	ON	PQ	NB	-	-	-	-
<i>collaris</i> Melsheimer														
<i>melsheimeri</i> Weise														
<i>S. kansanus</i> Casey	-	-	-	-	-	-	-	ON	PQ	-	-	-	-	-
<i>S. lacustris</i> LeConte	-	YK	NT	BC	AB	SK	MB	ON	PQ	NB	NS	-	-	-
<i>cultratus</i> Wingo														
<i>S. marginicollis</i> Mannerheim	-	-	-	BC	AB	-	-	ON	PQ	-	-	-	-	-
<i>borealis</i> Hatch														
<i>californicus</i> Boheman														
<i>desertorum</i> Casey														
<i>S. postpictus</i> Casey	-	-	-	BC	AB	SK	-	-	-	-	-	-	-	-
<i>S. puncticollis</i> LeConte	-	-	-	-	-	-	-	ON	PQ	-	-	-	-	-
<i>S. rubricaudus</i> Casey	-	-	-	-	-	-	-	ON	PQ	-	-	-	-	-
<i>canterius</i> Casey														
<i>chromopygus</i> Casey														
<i>texanus</i> Casey														
<i>S. securus</i> J. Chapin	-	-	-	-	-	-	-	ON	-	-	-	-	-	-
<i>S. solidus</i> Casey	-	-	-	BC	-	-	-	-	-	-	-	-	-	-
<i>blaisdelli</i> Casey														

<i>S. suturalis</i> Thunberg†	-	-	-	-	-	-	-	-	PQ	-	-	-	-	-
<i>S. tahoensis</i> Casey	-	-	-	BC	-	-	-	-	-	-	-	-	-	-
<i>S. tenebrosus</i> Mulsant	-	-	NT	-	AB	-	-	ON	PQ	-	-	-	-	-
<i>S. uncus</i> Wingo	-	-	-	-	AB	SK	MB	ON	-	-	-	-	-	-
<i>S. utahensis</i> Gordon	-	-	-	-	AB	-	-	-	-	-	-	-	-	-
Genus NEPHUS Mulsant														
(Subgenus NEPHUS s. str.)														
<i>N. ornatus naviculatus</i> (Casey)	AK	-	-	BC	AB	SK	MB	ON	PQ	-	-	-	-	NF
<i>kincaidi</i> (Hatch)	-	-	-	-	-	-	-	ON	PQ	NB	NS	-	-	-
<i>N. ornatus ornatus</i> (LeConte)	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>frosti</i> (Casey)	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>sanguinifer</i> (Casey)	-	-	-	-	-	-	-	-	-	-	-	-	-	-
(Subgenus TURBOSCYMNUS Gordon)														
<i>N. georgei</i> (Weise)	AK	YK	NT	BC	AB	-	-	ON	PQ	-	-	-	-	NF
<i>bisignatus</i> (Horn)	-	-	-	-	-	-	-	-	-	-	-	-	-	-
(Subgenus SCYMNOBIUS Casey)														
<i>N. flavifrons</i> (Melsheimer)	-	-	-	-	-	-	-	ON	PQ	-	-	-	-	-
<i>bioculatus</i> (Mulsant)	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>guttiger</i> (Mulsant)	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>ludovicianus</i> (Casey)	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>marginellus</i> (Mulsant)	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>N. intrusus</i> (Horn)	-	-	-	-	-	-	-	ON	-	-	-	-	-	-
<i>inops</i> (Casey)	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>N. sordidus</i> (Horn)	-	-	-	-	AB	SK	MB	ON	-	-	-	-	-	-
<i>intrusoides</i> (Hatch)	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Genus DIOMUS Mulsant														
<i>D. amabilis</i> (LeConte)	-	-	-	-	-	-	-	-	PQ	-	-	-	-	-
<i>dulcis</i> (Casey)	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>emertoni</i> (Casey)	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>D. debilis</i> (LeConte)	-	-	-	-	AB	SK	-	ON	PQ	-	-	-	-	-
<i>aeger</i> (Casey)	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>minutissimus</i> (Casey)	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>molliculus</i> (Casey)	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>pusio</i> (Casey)	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>D. liebecki</i> (Horn)	-	-	-	-	-	-	MB	ON	-	-	-	-	-	-
<i>adulans</i> (Casey)	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>ohioensis</i> (Stehr)	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>D. terminatus</i> (Say)	-	-	-	-	-	-	-	ON	-	-	-	-	-	-
<i>femoralis</i> (LeConte)	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>partitus</i> (Casey)	-	-	-	-	-	-	-	-	-	-	-	-	-	-

Tribe Hyperaspini

Genus HYPERASPIDIUS Crotch														
<i>H. comparatus</i> Casey	-	-	-	BC	-	-	-	-	-	-	-	-	-	-
<i>juniperus</i> Nunenmacher	-	-	-	-	AB	-	-	-	-	-	-	-	-	-
<i>H. hercules</i> Belicek	-	-	-	-	AB	-	-	-	-	-	-	-	-	-
<i>H. insignis</i> Casey	-	-	-	-	AB	-	-	-	-	-	-	-	-	-
<i>H. mimus</i> Casey	-	-	-	-	AB	SK	-	-	-	-	-	-	-	-
<i>carri</i> Nunenmacher	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>coloradensis</i> Nunenmacher	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>H. vittigerus</i> (LeConte)	-	YK	-	BC	AB	SK	-	-	-	-	-	-	-	-
Genus HYPERASPIS Redtenbacher														
<i>Oxynychus</i> LeConte	-	-	-	-	-	-	-	ON	PQ	NB	-	-	-	-
<i>H. bigeminata</i> (Randall)	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>guexi</i> Mulsant	-	-	-	-	-	SK	MB	ON	PQ	NB	-	-	-	-
<i>H. binotata</i> (Say)	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>affinis</i> (Randall)	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>leucopsis</i> Melsheimer	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>normata</i> (Say)	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>paspalis</i> Watson	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>H. borealis</i> Dobzhansky	-	YK	-	BC	AB	-	-	-	-	-	-	-	-	-
<i>elali</i> Nutting	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>obscura</i> Malkin	-	-	-	-	-	-	-	-	-	-	-	-	-	-

<i>schuhi</i> Hatch															
<i>simuloides</i> Hatch															
H. consimilis LeConte	-	-	NT	BC	AB	SK	MB	-	PQ	NB	-	-	-	-	-
<i>canadensis</i> Dobzhansky															
H. conviva Casey	-	-	-	-	-	SK	MB	ON	PQ	-	-	-	-	-	-
<i>congressis</i> Watson															
<i>insolens</i> Casey															
H. disconotata Mulsant	-	-	-	-	AB	-	MB	ON	PQ	NB	-	-	-	-	-
H. dissoluta dissoluta Crotch	-	-	-	BC	-	-	-	-	-	-	-	-	-	-	-
<i>coloradana</i> Casey															
<i>nupta</i> Casey															
H. fastidiosa Casey	-	-	-	BC	AB	-	-	-	-	-	-	-	-	-	-
<i>septentrionis</i> Dobzhansky															
H. inflexa Casey	-	-	-	BC	AB	SK	MB	-	PQ	-	-	-	-	-	-
<i>serena</i> Casey															
H. jasperensis Belicek	-	-	NT	-	AB	-	-	-	-	-	-	-	-	-	-
H. lateralis Mulsant	-	-	-	BC	AB	SK	-	-	-	-	-	-	-	-	-
<i>flammula</i> Nunenmacher															
<i>idae</i> Nunenmacher															
<i>laevipennis</i> Casey															
<i>montanica</i> Casey															
<i>nigrocauda</i> Dobzhansky															
<i>omissa</i> Casey															
<i>wellmani</i> Nunenmacher															
H. moerens (LeConte)	-	-	-	-	-	-	-	ON	PQ	-	-	-	-	-	-
H. octavia Casey	-	-	-	-	-	SK	-	ON	PQ	NB	-	-	-	-	-
H. oregona Dobzhansky	-	-	-	BC	AB	-	-	-	-	-	-	-	-	-	-
<i>lanei</i> Hatch															
H. postica LeConte	-	-	-	BC	AB	SK	-	-	-	-	-	-	-	-	-
<i>angustula</i> Casey															
<i>elliptica</i> Casey															
<i>essigi</i> Malkin															
H. pratensis LeConte	-	-	-	-	-	-	-	-	PQ	-	-	-	-	-	-
H. proba (Say)	-	-	-	-	-	SK	MB	ON	PQ	-	-	-	-	-	-
<i>trinifer</i> Casey															
H. quadrivittata LeConte	-	-	-	BC	AB	SK	-	-	-	-	-	-	-	-	-
<i>tetraneura</i> Casey															
H. signata signata (Olivier)	-	-	-	-	-	SK	-	-	-	-	-	-	-	-	-
<i>taedata</i> LeConte															
H. simulatrix Dobzhansky	-	-	-	BC	AB	SK	-	-	-	-	-	-	-	-	-
H. troglodytes Mulsant	-	-	-	-	-	-	MB	-	PQ	-	NS	-	-	-	-
<i>discreta</i> LeConte															
<i>novascotiae</i> E.A. Chapin															
H. undulata (Say)	-	-	-	BC	AB	SK	MB	ON	PQ	NB	-	-	-	-	-
<i>elegans</i> Mulsant															
<i>guttifera</i> Weise															
<i>maculifera</i> Melsheimer															
Genus BRACHIACANTHA Dejean															
B. albifrons (Say)	-	-	NT	-	AB	SK	MB	-	-	-	-	-	-	-	-
<i>pacifica</i> Casey															
B. decempustulata (Melsheimer)	-	-	-	-	-	-	-	ON	PQ	NB	NS	-	-	-	-
<i>stellata</i> Casey															
<i>troglydytes</i> Mulsant															
B. dentipes (Fabricius)	-	-	-	-	-	-	-	ON	-	-	-	-	-	-	-
<i>separata</i> Leng															
<i>socialis</i> Casey															
B. quadripunctata quadripunctata Melsheimer	-	-	-	-	-	-	-	ON	PQ	-	-	-	-	-	-
<i>basalis</i> Melsheimer															
<i>confusa</i> Mulsant															
<i>diversa</i> Mulsant															
B. rotunda Gordon	-	-	-	-	-	-	-	ON	PQ	-	-	-	-	-	-
B. ursina (Fabricius)	-	-	-	-	AB	SK	MB	ON	PQ	NB	NS	-	-	-	-
<i>congruens</i> Casey															
B. uteella Casey	-	-	-	BC	-	-	-	-	-	-	-	-	-	-	-
<i>fenyessi</i> Leng															
<i>lengi</i> Nunenmacher															
<i>sonorana</i> Casey															

Subfamily CHILOCORINAE**Tribe Chilacorini**

Key to genera: Chapin (1965)

Genus BRUMOIDES E.A. Chapin

B. histrio (Fall)	-	-	-	BC	-	-	-	-	-	-	-	-	-	-
<i>parvicollis</i> (Casey)	-	-	-	BC	-	-	-	-	-	-	-	-	-	-
B. septentrionis septentrionis (Weise)	AK	YK	NT	BC	AB	SK	MB	ON	-	-	-	-	-	-
<i>desertorum</i> (Casey)	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>nevadensis</i> (Leng)	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>ovoideus</i> (Casey)	-	-	-	-	-	-	-	-	-	-	-	-	-	-
B. septentrionis davisi (Leng)	-	-	-	-	-	-	-	ON	PQ	-	-	-	-	-

Genus EXOCHOMUS Redtenbacher

E. aethiops (Bland)	-	-	-	-	AB	SK	-	-	-	-	-	-	-	-
<i>mormonicus</i> Casey	-	-	-	-	AB	SK	-	-	-	-	-	-	-	-
E. californicus Casey	-	-	-	BC	-	-	-	-	-	-	-	-	-	-

Genus CHILOCORUS Leach

C. hexacyclus Smith	-	-	-	-	AB	SK	-	-	-	-	-	-	-	-
C. stigma (Say)	-	-	-	-	AB	SK	-	-	-	-	-	-	-	-
<i>bivulnerus</i> Mulsant	-	-	-	-	AB	SK	MB	ON	PQ	NB	NS	-	-	-
C. tricyclus Smith	-	-	-	BC	-	-	-	-	-	-	-	-	-	-

Subfamily COCCIDULINAE**Tribe Coccidulini**

Genus COCCIDULA Kugelann

C. lepida LeConte	AK	YK	NT	BC	AB	SK	MB	ON	PQ	NB	-	-	-	-
<i>occidentalis</i> Horn	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>reitteri</i> Dodge	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>suturalis</i> Reitter	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>suturalis</i> Weise	-	-	-	-	-	-	-	-	-	-	-	-	-	-

Genus RHYZOBIOUS Stephens

Lindorus Casey

R. lophanthae (Blaisdell)†	-	-	-	BC	-	-	-	-	-	-	-	-	-	-
----------------------------	---	---	---	----	---	---	---	---	---	---	---	---	---	---

Subfamily COCCINELLINAE**Tribe Coccinellini**

Genus ANISOSTICTA Dejean

A. bitriangularis (Say)	AK	YK	NT	BC	AB	SK	MB	ON	PQ	NB	NS	-	LB	NF
<i>irregularis</i> Weise	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>multiguttata</i> (Randall)	-	-	-	-	-	-	-	-	-	-	-	-	-	-
A. borealis Timberlake	AK	YK	NT	-	-	-	MB	-	PQ	-	-	-	-	-

Genus MACRONAEMIA Casey

M. episcopalis (Kirby)	AK	YK	NT	BC	AB	SK	MB	ON	PQ	-	-	-	-	-
------------------------	----	----	----	----	----	----	----	----	----	---	---	---	---	---

Genus PARANAEMIA Casey

P. vittigera (Mannerheim)	-	-	-	?	?	-	-	-	-	-	-	-	-	-
<i>similis</i> Casey	-	-	-	-	-	-	-	-	-	-	-	-	-	-

Genus COLEOMEGILLA Timberlake

Megilla Mulsant

C. maculata lengi Timberlake	-	-	-	-	-	-	-	ON	PQ	-	-	-	-	-
------------------------------	---	---	---	---	---	---	---	----	----	---	---	---	---	---

Genus CERATOMEGILLA Crotch

C. ulkei Crotch	AK	YK	NT	BC	AB	-	-	ON	-	-	-	-	-	-
<i>parva</i> (Watson)	-	-	-	-	-	-	-	-	-	-	-	-	-	-

Genus HIPPODAMIA Dejean

H. americana Crotch	-	-	NT	BC	AB	SK	MB	ON	PQ	-	-	-	-	-
H. apicalis Casey	-	-	-	BC	-	-	-	-	-	-	-	-	-	-
<i>lengi</i> Johnson														
<i>nigromaculata</i> (Nunenmacher)														
<i>tricolor</i> Nunenmacher														
H. arctica (Schneider)*	AK	YK	NT	BC	-	-	-	-	PQ	-	-	-	LB	-
H. caseyi Johnson	-	-	-	BC	AB	-	-	-	-	-	-	-	-	-
H. convergens Guérin-Ménéville	-	-	-	BC	AB	SK	MB	ON	PQ	NB	-	-	-	-
<i>juncta</i> Casey														
<i>modesta</i> Melsheimer														
<i>obsoleta</i> Crotch														
<i>praticola</i> Mulsant														
H. expurgata Casey	-	YK	-	BC	AB	SK	-	-	-	-	-	-	-	-
H. falcigera Crotch	AK	YK	NT	BC	AB	SK	MB	-	PQ	-	-	-	-	-
<i>albertana</i> Casey														
<i>cottlei</i> (Nunenmacher)														
H. glacialis glacialis (Fabricius)	-	-	NT	BC	AB	SK	MB	ON	PQ	-	-	-	-	-
H. glacialis lecontei Mulsant	-	-	NT	-	AB	SK	MB	-	-	-	-	-	-	-
<i>abducens</i> Casey														
<i>defecta</i> Johnson														
<i>hoppingi</i> Nunenmacher														
<i>mackenzie</i> E.A. Chapin														
<i>pseudoglacialis</i> Johnson														
H. lunatomaculata dobzhanskyi	-	YK	-	BC	-	-	-	-	-	-	-	-	-	-
E.A. Chapin														
H. moesta moesta LeConte	-	-	-	BC	-	-	-	-	-	-	-	-	-	-
H. moesta bowditchi Johnson	-	-	-	BC	AB	SK	MB	-	-	-	-	-	-	-
H. oregonensis Crotch	-	-	-	BC	AB	-	-	-	-	-	-	-	-	-
<i>cockerelli</i> Johnson														
<i>dispar</i> Casey														
<i>liliputana</i> Casey														
<i>puncticollis</i> Casey														
H. parenthesis (Say)	AK	YK	NT	BC	AB	SK	MB	ON	PQ	NB	NS	-	-	-
<i>albomacula</i> Fitch														
<i>approximata</i> Fitch														
<i>confluenta</i> Fitch														
<i>connata</i> Fitch														
<i>discopunctata</i> Fitch														
<i>insulata</i> Fitch														
<i>linearis</i> Fitch														
<i>lituricollis</i> Fitch														
<i>nimia</i> Fitch														
<i>permacrifrons</i> Fitch														
<i>triangularis</i> Fitch														
<i>tridens</i> (Kirby)														
<i>tridentifrons</i> Fitch														
H. quindecimmaculata Mulsant	-	-	-	-	-	SK	-	ON	-	-	-	-	-	-
H. quinquesignata quinquesignata	AK	YK	NT	BC	AB	SK	MB	ON	PQ	NB	-	PE	-	-
(Kirby)														
<i>coccinea</i> Casey														
<i>leporina</i> Mulsant														
<i>mulsanti</i> LeConte														
<i>pugetana</i> Casey														
<i>quadraria</i> Casey														
<i>subsimilis</i> Casey														
<i>uteana</i> Casey														
<i>vernix</i> Casey														
H. quinquesignata ambigua LeConte	-	-	-	BC	-	-	-	-	-	-	-	-	-	-
<i>obliqua</i> Casey														
<i>punctulata</i> LeConte														
H. sinuata crotchi Casey	-	-	NT	BC	AB	SK	MB	-	-	-	-	-	-	-
<i>disjuncta</i> Timberlake														
<i>fontinalis</i> Casey														
<i>straminea</i> E.A. Chapin														
H. sinuata spuria LeConte	AK	YK	-	BC	-	-	-	-	-	-	-	-	-	-
<i>complex</i> Casey														
H. tredecimpunctata tibialis (Say)	AK	YK	NT	BC	AB	SK	MB	ON	PQ	NB	NS	PE	LB	NF

<i>H. variegata</i> (Goeze)†	-	-	-	-	-	-	-	ON	PQ	-	-	-	-	-
<i>H. washingtoni</i> Timberlake	-	-	-	BC	-	-	-	-	-	-	-	-	-	-
Genus ANATIS Mulsant														
Key to species: Watson (1976)														
<i>A. labiculata</i> (Say)	-	-	-	-	-	-	MB	ON	PQ	NB	-	-	-	-
<i>canadensis</i> Provancher														
<i>caseyi</i> Westcott														
<i>quindecimpunctata</i> (Olivier)														
<i>A. lecontei</i> Casey	-	-	-	BC	AB	-	-	-	-	-	-	-	-	-
<i>A. mali</i> (Say)	AK	YK	NT	BC	AB	SK	MB	ON	PQ	NB	NS	PE	-	NF
<i>borealis</i> Belicek														
<i>signaticollis</i> Mulsant														
<i>A. rathvoni</i> (LeConte)	-	-	-	BC	AB	SK	-	-	-	-	-	-	-	-
Genus MYZIA Mulsant														
<i>Neomysia</i> Casey														
<i>Paramysia</i> Reitter														
<i>M. interrupta</i> (Casey)	-	-	-	BC	AB	-	-	-	-	-	-	-	-	-
<i>caseyi</i> (Timberlake)														
<i>M. pullata</i> (Say)	-	YK	-	-	AB	SK	MB	ON	PQ	NB	NS	-	LB	-
<i>montana</i> (Casey)														
<i>notans</i> (Randall)														
<i>randalli</i> (Casey)														
<i>M. subvittata</i> (Mulsant)	-	-	-	BC	AB	-	-	-	-	-	-	-	-	-
<i>hornii</i> Crotch														
<i>oregona</i> (Casey)														
Genus CALVIA Mulsant														
<i>Anisocalvia</i> Crotch														
<i>C. quatuordecimguttata</i> (Linné)*	AK	YK	NT	BC	AB	SK	MB	ON	PQ	NB	NS	-	LB	NF
<i>bicordifera</i> (Casey)														
<i>cardisce</i> (Randall)														
<i>complexa</i> Nunenmacher														
<i>elliptica</i> (Casey)														
<i>hesperica</i> (Crotch)														
<i>incarnata</i> (Kirby)														
<i>lacustris</i> (Casey)														
<i>obliqua</i> (Randall)														
<i>postplagiata</i> (Casey)														
<i>quadrisignata</i> (Casey)														
<i>sicardi</i> (Nunenmacher)														
<i>similis</i> (Randall)														
<i>uniformis</i> (Casey)														
<i>vancouveri</i> (Casey)														
<i>victoriana</i> (Casey)														
Genus ADALIA Mulsant														
<i>A. bipunctata</i> (Linné)‡	AK	YK	NT	BC	AB	SK	MB	ON	PQ	NB	NS	-	LB	NF
<i>annectans</i> (Crotch)														
<i>bioculata</i> (Say)														
<i>coloradensis</i> Casey														
<i>disjuncta</i> (Randall)														
<i>duplicata</i> Leng														
<i>frigida</i> (Schneider)														
<i>humeralis</i> (Say)														
<i>ludovicae</i> Mulsant														
<i>melanopleura</i> (LeConte)														
<i>ocellata</i> Johnson														
<i>ophthalmica</i> Mulsant														
<i>ornatella</i> Casey														
<i>ovipennis</i> Casey														
<i>postica</i> Johnson														
<i>sempustulata</i> Johnson														
<i>transversalis</i> Casey														
Genus COCCINELLA Linné														
Key to North American species: Brown (1962b)														

Coccinellidae

<i>C. alta</i> W.J. Brown	-	-	-	BC	AB	-	-	-	-	-	-	-	-	-
<i>suturalis</i> Casey														
<i>C. californica</i> Mannerheim	AK	-	-	BC	AB	-	-	-	-	-	-	-	-	-
<i>melanocollis</i> Johnson														
<i>C. fulgida</i> Watson	AK	YK	NT	BC	-	-	-	ON	PQ	-	-	-	-	-
<i>C. hieroglyphica humboldtiensis</i>	-	-	-	BC	-	-	-	-	-	-	-	-	-	-
Nunenmacher														
<i>C. hieroglyphica kirbyi</i> Crotch	-	YK	NT	BC	AB	SK	MB	ON	PQ	NB	NS	-	-	-
<i>tricuspidis</i> Kirby														
<i>C. hieroglyphica mannerheimi</i>	AK	YK	NT	-	-	-	MB	-	-	-	-	-	-	-
Mulsant														
<i>C. johnsoni</i> Casey	AK	-	-	BC	-	-	-	-	-	-	-	-	-	-
<i>C. monticola</i> Mulsant	AK	-	NT	BC	AB	SK	MB	ON	PQ	NB	NS	-	-	-
<i>alutacea</i> Casey														
<i>biguttata</i> Johnson														
<i>confluenta</i> Johnson														
<i>impressa</i> Casey														
<i>lacustris</i> LeConte														
<i>nevadica</i> Casey														
<i>postica</i> Johnson														
<i>sellica</i> Johnson														
<i>C. novemnotata</i> Herbst	-	-	NT	BC	AB	SK	MB	ON	PQ	-	-	-	-	-
<i>confluenta</i> Fitch														
<i>conjuncta</i> Fitch														
<i>degener</i> Casey														
<i>divisicollis</i> Fitch														
<i>franciscana</i> Mulsant														
<i>inaequalis</i> Fitch														
<i>oregona</i> Casey														
<i>parvamaculata</i> Fitch														
<i>C. prolongata</i> prolongata Crotch	-	-	-	BC	AB	-	-	-	-	-	-	-	-	-
<i>C. septempunctata</i> Linné†	-	-	-	BC	AB	SK	MB	ON	PQ	NB	NS	-	-	-
<i>C. transversoguttata richardsoni</i>	AK	YK	NT	BC	AB	SK	MB	ON	PQ	NB	NS	-	LB	NF
W.J. Brown														
<i>interrupta</i> Fitch														
<i>quinenotata</i> Kirby														
<i>zetterstedti</i> Mader														
<i>C. trifasciata perplexa</i> Mulsant	AK	YK	NT	BC	AB	SK	MB	ON	PQ	NB	NS	-	LB	NF
<i>C. trifasciata subversa</i> LeConte	-	-	-	BC	-	-	-	-	-	-	-	-	-	-
<i>barda</i> LeConte														
<i>euginii</i> Mulsant														
<i>juliana</i> Mulsant														
<i>praedicta</i> Mader														
<i>C. undecimpunctata undecimpunctata</i>	-	-	-	BC	-	-	-	ON	PQ	NB	NS	PE	-	NF
Linné†														
Genus PROPYLAEA Mulsant														
<i>P. quatuordecimpunctata</i> (Linné)†	-	-	-	-	-	-	-	ON	PQ	NB	-	-	-	-
Genus CYCLONEDA Crotch														
<i>C. munda</i> (Say)	-	-	-	-	-	-	MB	ON	PQ	-	-	-	-	-
<i>C. polita</i> Casey	-	-	-	BC	AB	SK	-	-	-	-	-	-	-	-
<i>flava</i> Timberlake														
Genus OLLA Casey														
<i>O. v-nigrum</i> (Mulsant)	-	-	-	BC	AB	SK	MB	ON	-	-	-	-	-	-
<i>abdominalis</i> (Say)														
<i>arizonae</i> Casey														
<i>fenestralis</i> Casey														
<i>minuta</i> Casey														
<i>plagiata</i> Casey														
<i>sayi</i> (Crotch)														
<i>semilunaris</i> Johnson														
<i>sobrina</i> Casey														
Genus NEOHARMONIA Crotch														
<i>Agrabia</i> Casey														
<i>Harmoniaspis</i> Casey														
<i>Neoharmonia</i> Casey														

N. venusta venusta (Melsheimer)	-	-	-	-	-	-	-	ON	-	-	-	-	-	-
<i>centralis</i> Casey														
<i>dissimila</i> Blatchley														
<i>fattigi</i> Blatchley														
<i>notulata</i> (Mulsant)														

Genus MULSANTINA Weise
Pseudocleis Casey
 Key to species: Chapin (1985)

M. hudsonica (Casey)	AK	YK	NT	BC	AB	SK	MB	ON	PQ	NB	NS	PE	LB	-
M. picta (Randall)	-	YK	-	BC	AB	SK	MB	ON	PQ	NB	NS	-	-	-
<i>blanchardi</i> (Johnson)														
<i>concinata</i> (Melsheimer)														
<i>contexta</i> (Mulsant)														
<i>impictipennis</i> (Weise)														
<i>minor</i> (Casey)														
<i>nubilata</i> (Casey)														

Tribe Psylloborini

Genus PSYLLOBORA Dejean

P. borealis Casey	-	-	-	BC	-	-	-	-	-	-	-	-	-	-
<i>deficiens</i> Casey														
P. vigintimaculata (Say)	AK	YK	NT	BC	AB	SK	MB	ON	PQ	NB	NS	-	-	NF
<i>obsoleta</i> Casey														
<i>separata</i> Casey														
<i>taedata</i> LeConte														

Subfamily EPILACHNINAE

Key to New World species: Gordon (1975)

Tribe Epilachnini

Genus EPILACHNA Dejean

E. varivestis Mulsant	-	-	-	-	-	-	-	ON	PQ	-	-	-	-	-
<i>cervina</i> Mulsant														
<i>corrupta</i> Mulsant														
<i>cuprea</i> Cockerell														
<i>genuina</i> Mulsant														
<i>juncta</i> Johnson														
<i>maculiventris</i> Bland														
<i>toweri</i> Johnson														

Family ENDOMYCHIDAE handsome fungus beetles

J.M. Campbell

The family Endomychidae is found in most regions of the world, but the majority of species occur in tropical areas. The range of 15 species extends northward into Canada. Adults and larvae may be collected on soft fungi, in leaf litter, or under bark with fungal growth. One species, *Mycetaea subterranea* (*M. hirta* in most publications), is a pest of stored products. Most species feed on fungi.

Because of their moderately large size and often bright coloration, many North American species are

taxonomically well known. However, the family has never been revised for North America. Strohecker (1953) reviewed the genera of the world and published a catalog of the North American species (1986). Hatch (1962) treated the genera and species that occur in the Pacific Northwest. A modern revision of the North American species of this family is needed.

NT (1); BC (7); AB (2); SK (2); MB (5); ON (10); PQ (8); NB (2); NS (6); PE (1); NF (2) • I (2)

Subfamily MYCETAEINAE
Tribe Mycetaeini

Genus SYMBIOTES Redtenbacher

S. duryi Blatchley	-	-	-	-	-	-	-	ON	-	-	-	-	-	-
<i>lacustris</i> Casey														
<i>oblongus</i> Casey														
<i>pilosus</i> Casey														
<i>waltoni</i> Dury														
S. gibberosus (Lucas)†	-	-	-	-	-	-	-	ON	-	-	-	-	-	-
<i>duryi</i> Walton														
<i>montanus</i> Casey														

Genus MYCETAEA Stephens

M. subterranea (Fabricius)†	-	-	-	BC	-	-	-	ON	PQ	-	NS	PE	-	NF
<i>fumata</i> Stephens														
<i>hirta</i> (Marsham)														

Tribe Leiestini

Key to North American species: Blaisdell (1931)

Genus RHANIDEA Strohecker

Rhanis LeConte

R. unicolor (Ziegler)	-	-	-	-	-	-	-	ON	PQ	-	-	-	-	-
<i>apicalis</i> (Melsheimer)														
<i>haemorrhoidalis</i> (Guérin-Méneville)														

Genus STETHORHANIS Blaisdell

S. borealis Blaisdell	-	-	-	BC	-	-	-	-	-	-	-	-	-	-
-----------------------	---	---	---	----	---	---	---	---	---	---	---	---	---	---

Genus PHYMAPHORA Newman

P. californica Horn	-	-	-	BC	-	-	-	-	-	-	-	-	-	-
P. pulchella Newman	-	-	-	-	-	-	MB	ON	PQ	-	NS	-	-	-
<i>crassicornis</i> (Melsheimer)														
<i>puncticollis</i> (Ziegler)														

Subfamily STENOTARSINAE
Tribe Stenotarsini

Genus DANAE Reiche

D. testacea (Ziegler)	-	-	-	-	-	-	-	ON	PQ	-	NS	-	-	-
<i>ferruginea</i> (Guérin-Méneville)														
<i>lutea</i> (Melsheimer)														
<i>pilosa</i> (Melsheimer)														

Subfamily EUMORPHINAE
Tribe Lycoperdinini

Genus LYCOPERDINA Latreille

(Subgenus GOLGIA Mulsant)

L. ferruginea LeConte	-	-	-	BC	AB	SK	MB	ON	PQ	-	NS	-	-	-
<i>angulata</i> (Say)														
<i>glabrata</i> Guérin-Méneville														

Tribe Mycetinini

Genus APHORISTA Gorham

A. laeta (LeConte)	-	-	-	BC	-	-	-	-	-	-	-	-	-	-
A. vittata (Fabricius)	-	-	-	-	-	-	MB	ON	PQ	-	-	-	-	-
<i>distincta</i> (Say)														
<i>lineata</i> (Olivier)														
<i>ovipennis</i> Casey														

Genus MYCETINA Mulsant

M. idahoensis Fall

M. perpulchra (Newman)

quadriguttata (Melsheimer)

-	-	-	BC	-	-	-	-	-	-	-	-	-	-	-
-	-	-	-	-	-	MB	ON	PQ	NB	NS	-	-	-	-

Subfamily ENDOMYCHINAE
Tribe Endomychini

Genus ENDOMYCHUS Panzer

E. biguttatus Say

E. limbatus (Horn)

endomychoides (Fall)

-	-	NT	-	-	SK	MB	ON	PQ	NB	NS	-	-	-	NF
-	-	-	BC	AB	-	-	-	-	-	-	-	-	-	-

Family LATHRIDIIDAE minute brown scavenger beetles

Y. Bousquet

In Canada and Alaska 55 species of lathridiids have been recorded. These species are usually found on foliage, in rotten wood, under logs, and under the bark of dead trees, as well as in fungi and nests of social Hymenoptera, birds, and mammals. A number of species, including *Cartodere constricta* and *Aridius nodifer*, are regularly found in stored products, especially those that are damp and moldy. Adults and larvae feed on fungi, particularly molds.

Lathridiidae is another family of Cucujoidea that has been poorly studied in North America.

Most of the genera, particularly those of the subfamily Corticariinae, are in need of modern revision. The generic classification of the Lathridiinae used here follows that of Walkley (1952).

AK (19); YK (5); NT (6); BC (34); AB (9); SK (8); MB (17); ON (22); PQ (22); NB (6); NS (8); PE (1); LB (1); NF (4) • H (6); I (15)

Subfamily LATHRIDIINAE

Genus DIENERELLA Reitter

Microgramme Walkley

D. arga (Reitter)†

D. cordicollis (Mannerheim)

D. costulata (Reitter)†

D. filiformis (Gyllenhal)†

D. filum (Aubé)†

D. ruficollis (Marsham)†

pulicaria (Melsheimer)

-	-	-	-	-	-	-	ON	PQ	-	-	-	-	-	-
AK	-	-	-	-	-	-	-	-	-	-	-	-	-	-
-	-	-	-	-	-	MB	ON	PQ	-	-	-	-	-	-
-	-	-	-	-	SK	MB	-	-	NB	-	PE	-	-	-
-	-	-	BC	-	-	MB	ON	PQ	-	-	-	-	-	-
-	-	-	BC	-	-	-	ON	PQ	-	NS	-	-	-	NF

Genus ENICMUS C.G. Thomson

E. aterrimus Motschulsky

ferrugineus Belon*opaculus* LeConte

E. cordatus Belon

E. fictus Fall

E. mimus Fall

E. nitens Fall

E. tenuicornis LeConte

laticollis LeConte

-	-	-	-	-	-	-	ON	PQ	-	-	-	-	-	-
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
-	-	-	BC	-	-	-	-	-	-	-	-	-	-	-
AK	-	NT	BC	AB	SK	MB	ON	PQ	-	-	-	-	-	-
AK	-	NT	BC	AB	SK	MB	-	-	-	-	-	-	-	-
-	-	-	-	AB	-	MB	ON	-	-	-	-	-	-	-
-	-	-	BC	-	-	-	-	PQ	-	-	-	-	-	-

Genus THES Semenov

T. bergrothi (Reitter)†

tricariniatus (W.J. Brown)

-	-	-	BC	-	SK	MB	-	-	NB	NS	-	-	-	-
---	---	---	----	---	----	----	---	---	----	----	---	---	---	---

Genus LATHRIDIUS Herbst

Conithassa C.G. Thomson

L. consimilis Mannerheim*

parallellocollis Mannerheim

L. crenatus LeConte

AK	-	-	-	-	-	-	-	-	-	-	-	-	-	-
-	-	-	BC	-	-	-	-	-	-	-	-	-	-	-

- Casey, T.L. 1897. Coleopterological notices, VII. *Annals of the New York Academy of Sciences* 9:285-684.
- Casey, T.L. 1898. Studies in the Ptinidae, Cioidae and Sphindidae of America. *Journal of the New York Entomological Society* 6:61-93.
- Casey, T.L. 1900. Review of the American Corylophidae, Cryptophagidae, Tritomidae, and Dermestidae, with other studies. *Journal of the New York Entomological Society* 8:51-172.
- Casey, T.L. 1916. Some random studies among the Clavicornia. Pages 35-292 in *Memoirs on the Coleoptera*, Vol. 7. New Era Printing, Lancaster, Pa. 300 pp.
- Cawthra, E.M. 1957. Description of a new species of *Grypoidius* Schönherr (Col.: Curculionidae) with a key to the genus. *Proceedings of the Royal Entomological Society of London, Series B, Taxonomy* 26:127-130.
- Chagnon, G.; Robert, A. 1962. Principaux Coléoptères de la province de Québec. Les Presses de l'Université de Montréal, Montreal, Que. 440 pp.
- Chamberlin, W.J. 1926. Catalogue of the Buprestidae of North America north of Mexico. Privately published, Corvallis, Ore. 289 pp.
- Chandler, D.S. 1975. A revision of *Tanarthrus* LeConte with a presentation of its mid-Cenozoic speciation (Coleoptera: Anthicidae). *Transactions of the American Entomological Society* 101:319-354.
- Chandler, D.S. 1977. New *Mecynotarsus* with a key to the New World species (Coleoptera: Anthicidae). *The Coleopterists Bulletin* 31:363-370.
- Chandler, D.S. 1982. A revision of North American *Notoxus* with a cladistic analysis of the New World species (Coleoptera: Anthicidae). *Entomography* 1:333-438.
- Chandler, D.S. 1990a. Insecta: Coleoptera Pselaphidae. Pages 1175-1190 in Dindal, D.L., ed. *Soil biology guide*. John Wiley & Sons, New York, N.Y. 1349 pp.
- Chandler, D.S. 1990b. The Pselaphidae (Coleoptera) of Latimer County, Oklahoma, with revisions of four genera from eastern North America Part 1. Faroninae and Euplectinae. *Transactions of the American Entomological Society* 115(1989):503-529.
- Chantal, C. 1985. Les Tetratomidae (Coleoptera) du Québec. *Faberies* 11:43-76.
- Chapin, E.A. 1965. The genera of the Chilacorini (Coleoptera, Coccinellidae). *Bulletin of the Museum of Comparative Zoology at Harvard College* 133:227-271.
- Chapin, J.B. 1974. The Coccinellidae of Louisiana (Insecta: Coleoptera). Louisiana State University, Agricultural Experiment Station, Bulletin No. 682. 87 pp.
- Chapin, J.B. 1985. Revision of the genus *Mulsantina* Weise (Coleoptera: Coccinellidae). *Annals of the Entomological Society of America* 78:348-368.
- Chemsak, J.A. 1963. Taxonomy and bionomics of the genus *Tetraopes* (Cerambycidae: Coleoptera). University of California Publications in Entomology No. 30. 90 pp.
- Chemsak, J.A.; Linsley, E.G. 1982. Checklist of the Cerambycidae and Disteniidae of North America, Central America, and the West Indies (Coleoptera). Plexus Publishing, Medford, N.J. 138 pp.
- Clark, S.M. 1983. A revision of the genus *Microrhopala* (Coleoptera: Chrysomelidae) in America north of Mexico. *The Great Basin Naturalist* 43:597-618.
- Clark, W.E. 1971. A taxonomic revision of the weevil genus *Tychius* Germar in America north of Mexico (Coleoptera: Curculionidae). *Brigham Young University Science Bulletin (Biological Series)* 13(3):1-39.
- Clark, W.E. 1978. The weevil genus *Sibinia* Germar: Natural history, taxonomy, phylogeny, and zoogeography, with revision of the New World species (Coleoptera: Curculionidae). *Quaestiones entomologicae* 14:91-387.
- Clark, W.E. 1980. Revision of Nearctic weevils of the genus *Lignyodes* Dejean (Coleoptera: Curculionidae). *Transactions of the American Entomological Society* 106:273-326.
- Clark, W.E. 1982. Classification of the weevil tribe Lignyodini (Coleoptera, Curculionidae, Tychiinae), with revision of the genus *Plocetes*. *Transactions of the American Entomological Society* 108:11-151.
- Clark, W.E. 1987. Revision of the Nearctic species of *Pseudanthonomus* Dietz (Coleoptera: Curculionidae). *The Coleopterists Bulletin* 41:263-285.
- Cobos, A. 1964. Materiales para el estudio de la familia Eucnemidae. Primera parte (Coleoptera). *Eos* 40:289-435.



GO SATO 1990

Canada