

REVISION OF THE AUSTRALIAN COCCINELLIDAE (COLEOPTERA). PART 4. TRIBE TELSIMINI

ADAM ŚLIPIŃSKI¹, HONG PANG² and ROBERT D. POPE³

¹CSIRO Entomology, GPO Box 1700, Canberra, ACT 2601, Australia;
e-mail: Adam.Slipinski@ento.csiro.au

²State Key Laboratory of Biocontrol, Institute of Entomology, Zhongshan University Guangzhou
510275, China

³Brackley Burn, Park Street; Slinfold, Sussex RH13 7RU, England

Abstract.— The Australian members of the Coccinellid tribe Telsimini are revised which now includes 2 genera and 16 species. Nomenclatural history, diagnoses and distribution are provided for each genus and species. Keys to the genera and species are also presented. The lectotypes are designated for: *Lipernes creber* Blackburn, 1895, *L. gibbosus* Blackburn, 1895, *L. subviridis* Blackburn, 1892 and *Serangium obscuripes* Lea, 1902. *Telsimia abdita*, *T. acaciae*, *T. cassinula*, *T. glorious*, *T. leucoceps*, *T. occidua*, *T. palitans*, *T. rossi* and *T. rotunda* spp. nov. are described from Australia. *Serangium obscuripes* Lea, 1902 is transferred to *Telsimia* and *T. elainae* Chazeau is newly recorded from Australia.



Key words.— Entomology, taxonomy, review, Cucujoidea, Coccinellidae, Telsimini, New Guinea, Australia.