
# **PROCEEDINGS**

of the

# ENTOMOLOGICAL SOCIETY



of WASHINGTON

# PUBLISHED QUARTERLY

# **CONTENTS**

BAKER, G. T.—Morphology of the puparium of <i>Lipoptena mazamae</i> (Diptera: Hippoboscidae) BOTTORFF, R. L., S. W. SZCZYTKO and A. W. KNIGHT.—Descriptions of a new species and three incompletely known species of western nearctic <i>Isoperla</i> (Plecoptera: Per-	224
lodidae)	286
BROWN, B. V.— <i>Triphleba vitrinervis</i> (Malloch), an unrecognized species of <i>Crinophleba</i> Borgmeier (Diptera: Phoridae)	27 I
DAVIDSON, R. L. and H. J. LEE, JR.—Distribution and habitat of <i>Dyschirius campicola</i> Lindroth (Coleoptera: Carabidac) with new state records for Ohio and Illinois, first records east of the Mississippi River	205
DAVIS, D. R.—First record of a bagworm moth from Hawaii: description and introduction of <i>Brachycyttarus griseus</i> De Joannis (Lepidoptera: Psychidae)	259
DIETRICH, C. H. and S. H. McKAMEY.—Three new idiocerine leafhoppers (Homoptera: Cicadellidae) from Guyana with notes on ant-mutualism and subsociality	214
FREIDBERG, A. and W. N. MATHIS.—A new species of Craspedoxantha and a revised phylogeny for the genus (Diptera: Tephritidae)	325
GRISSELL, E. E. and M. E. SCHAUFF.—A synopsis of the seed-feeding genus Bephratelloides (Chalcidoidea: Eurytomidae)	I 7 7
GROGAN, W. L., JR. and W. W. WIRTH.—A new species of the minute predaceous midge genus Nannohelea from Sri Lanka (Diptera: Ceratopogonidae)	347
HERATY, J. M. and K. N. BARBER.—Biology of Obeza floridana (Ashmead) and Pseudo- chalcura gibbosa (Provancher) (Hymenoptera: Eucharitidae)	248
HOFFMAN, K. N. and J. C. MORSE.—Descriptions of the females of three <i>Polycentropus</i> species (Trichoptera: Polycentropodidae)	274
JOHNSON, N. F.— <i>Telenomus</i> (Hymenoptera: Scelionidae) egg parasites of <i>Erinnyis ello</i> (Lepidoptera: Sphingidae)	306
LaSALLE, J. and M. E. SCHAUFF.—New subfamily placement for some North American Eulophidae (Hymenoptera, Chalcidoidea)	282
(Continued on back cover)	

## THE

# **ENTOMOLOGICAL SOCIETY**

## OF WASHINGTON

ORGANIZED MARCH 12, 1884

#### **OFFICERS FOR 1990**

JEFFREY R. ALDRICH, President
DAVID R. SMITH, President-Elect
RICHARD G. ROBBINS, Recording Secretary
HOLLIS B. WILLIAMS, Corresponding Secretary
JAMES B. STRIBLING, Custodian

NORMAN E. WOODLEY, Treasurer GARY STECK, Program Chairman GEOFFREY B. WHITE, Membership Chairman F. Christian Thompson, Past President

ROBERT D. GORDON, Editor
THOMAS J. HENRY, Associate Editor

DONALD R. DAVIS

Publications Committee
Donald R. Whitehead
F. Christian Thompson

GEORGE C. STEYSKAL

Honorary President Curtis W. Sabrosky

LOUISE M. RUSSELL

Honorary Members
ALAN STONE

THEODORE L. BISSELL

All correspondence concerning Society business should be mailed to the appropriate officer at the following address: Entomological Society of Washington, % Department of Entomology, NHB 168, Smithsonian Institution, Washington, D.C. 20560.

MEETINGS.—Regular meetings of the Society are held in the Natural History Building, Smithsonian Institution, on the first Thursday of each month from October to June, inclusive, at 8 P.M. Minutes of meetings are published regularly in the *Proceedings*.

MEMBERSHIP.—Members shall be persons who have demonstrated interest in the science of entomology. Annual dues for members are \$20.00 (U.S. currency) of which \$18.00 is for a subscription to the Proceedings of the Entomological Society of Washington for one year.

PROCEEDINGS.—The *Proceedings* are published quarterly beginning in January by The Entomological Society of Washington, % Department of Entomology, NHB-168, Smithsonian Institution, Washington, D.C. Members in good standing receive the Proceedings of the Entomological Society of Washington. Nonmember subscriptions are \$50.00 per year, domestic, and \$60.00 per year, foreign (U.S. currency), payable in advance. Foreign delivery cannot be guaranteed. All remittances should be made payable to *The Entomological Society of Washington*.

The Society does not exchange its publications for those of other societies.

# PLEASE SEE P. 318 OF THE APRIL, 1989 ISSUE FOR INFORMATION REGARDING PREPARATION OF MANUSCRIPTS.

#### STATEMENT OF OWNERSHIP

Title of Publication: Proceedings of the Entomological Society of Washington.

Frequency of Issue: Quarterly (January, April, July, October).

Location of Office of Publication, Business Office of Publisher and Owner: The Entomological Society of Washington, % Department of Entomology, Smithsonian Institution, 10th and Constitution NW, Washington, D.C. 20560.

Editor: Robert D. Gordon, Systematic Entomology Laboratory, ARS, % Department of Entomology, Smithsonian Institution, 10th and Constitution NW, Washington, D.C. 20560.

Books for Review: T. Henry, Entomology, Smithsonian Institution, 10th and Constitution NW, Washington, D.C. 20560.

Managing Editor and Known Bondholders or other Security Holders: none.

#### This issue was mailed 1 May 1990

Second Class Postage Paid at Washington, D.C. and additional mailing office.

PRINTED BY ALLEN PRESS, INC., LAWRENCE, KANSAS 66044, USA

THIS DUDING ATION IS DOINTED ON ACID EDEE DADED

# A SURVEY OF THE COCCINELLIDAE (COLEOPTERA) ASSOCIATED WITH NURSERY STOCK IN MARYLAND

C. L. Staines, Jr., M. J. Rothschild, and R. B. Trumbule

Maryland Department of Agriculture, Office of Plant Industries and Resource Conservation, Plant Protection Section, 50 Harry S Truman Parkway, Annapolis, Maryland 21401.

Abstract. — Twenty-eight species of adult Coecinellidae were collected in Maryland nurseries from 1986 to 1988. Host plants, presence of prey, and active predation (if any) were recorded for each species. The most commonly detected species were Coccinella septempunctata L., Coleomegilla maculata lengi Timberlake, and Hippodamia convergens Guérin. Coccinellidae populations were seldom high and the number of times beetles were collected was low compared to the number of nurseries visited.

Key Words: host plants, prey

Recently there has been a shift in the pest control practices of commercial nurseries away from cover sprays to spot spraying. Under this new regime, pest control by natural enemies is both desirable and feasible. The Coceinellidae have long been known as major predators of various Homoptera and Acari. Gordon's (1985) treatise on North American Coccinellidae has facilitated adult identification to species. A survey of the Coccinellidae found in Maryland nurseries was undertaken to determine the variety of species present, their frequency of occurrence, and abundances.

### MATERIALS AND METHODS

Adult Coccinellidae were collected during routine nursery inspections. In addition to the typical collection data, host plants, presence and type of prey, and feeding activity by the coccinellids were also noted. Specimens were placed in 70% ethyl alcohol and taken to the laboratory for mounting. Adults were identified by the senior author using Gordon (1985). No efforts were made to identify larvae. Voucher specimens are de-

posited in the Maryland Department of Agriculture collection.

#### RESULTS AND DISCUSSION

Twenty-eight species representing eighteen genera were collected. Coccinellidae were found in 67 locations in 1986, 82 locations in 1987, and 37 locations in 1988. One reason for the low detection rate in 1988 is that several field personnel only reported unusual eoccinellids. Possibly another reason was that the drought conditions and high temperatures during the summer made most adult eoccinellids seek protected locations. Prey numbers were also low, possibly due to the same conditions.

Four species were found in high numbers, but most were observed as occasional adults scattered over large areas. The species found are listed below with the counties, host plants (names as listed in Hortus III, 1976), prey associations or feeding activities, and the months in which specimens were collected. The relative frequency of each species is indicated.

#### SCYMNINAE

# Scymnini

Scymnus cervicalis Mulsant: Anne Arundel, Prince George's. Catalpa sp., Quercus sp. Associated with aphids (Homoptera: Aphididae). May. Two locations.

S. tenebrosus Mulsant: Anne Arundel, Montgomery. Catalpa sp., Malus sp. Associated with aphids. May. Two locations.

Stethorus punctum punctum (LeConte): Prince George's. Gleditsia tricanthos L., Platanus sp., Quercus sp. April, July. Two locations.

# Hyperaspini

Brachicantha felina (Fab.): Montgomery, Prince George's. Betula sp., Malus sp. Associated with aphids. May, June, August. Four locations.

B. rotunda Gordon: Harford, Montgomery. Betula sp., Malus sp., Robinia pseudoacacia L. Associated with aphids. Four locations.

B. quadripunctata quadripunctata (Melsheimer): Prince George's. May. One specimen.

B. ursina (Fab.): Baltimore. June. One specimen.

Hyperaspis binotata (Say): Baltimore, Prince George's. Prunus sp.. Quercus sp. April, May. Associated with Melanaspis obscura (Comstock) (Homoptera: Diaspididae). Two locations.

H. signata signata (Olivier): Anne Arundel, Carroll. May, June. Juniperus scorulorum Sarg. Feeding on Carulaspis sp. (Homoptera: Diaspididae). Three specimens.

*H. proba* (Say): Baltimore, Kent, Prince George's. *Quercus* sp. July, November. Associated with *M. obscura*. Three locations.

## CHILOCORINAE

#### Chilocorini

Chilocorus kuwanae Silvestri: Specimens of this species were released against various Diaspididae. Recoveries were made in Prince George's and Worcester. Ilex cor-

nuta Lindl. & Paxt., I. crenata Thunb., Quercus spp. Feeding on Hemiberlesia lataniae (Signoret), Lopholeucaspis japonica (Cockerell), M. obscura. Two locations.

C. stigma (Say): Baltimore, Carroll, Howard, Kent, Montgomery, Prince George's, St. Mary's, Somerset. Acer sp., A. palmatum Thunb., Euonymus alata (Thunb.) Sieb., Fraxinus sp., Prunus sp., Quercus sp., Q. palustris Muenchh., Q. robur L. Associated with Lepidosaphes yangagicola Kuwana, M. obscura, Pseudaulacaspis sp. (Diasipididae), Eulecanium cerasorum (Cockerell) (Coccidae). March, April, May, June, July, September, November. Nineteen locations. This species was common in only one location. All other collections represent only a few individuals.

Exochomus marginipennis (LeConte): Kent, St. Mary's. Robinia pseudoacacia. May, November. Two specimens.

#### Coccidulinae

#### Coccidulini

Rhyzobius lophanthae (Blasidell): Montgomery. Pinus sp. October. One specimen.

#### Coccinellinae

### Coccinellini

Adalia bipunctata (L.): Anne Arundel. Baltimore, Caroline, Carroll, Harford, Kent. Montgomery, Prince George's, St. Mary's. Acer rubrum L., Betula sp., Carpinus betulus L., Elageanus umbellata Thunb., Euonymus alata, Forsythia intermedia Zab., G. triacanthos, Lonicera sempervirens L., Malus sp., Picea abies (L.) Karst., Pinus mugo Turra, P. strobus L., Prunus sp., Pyracantha sp., Quercus bicolor Willd., Rosa rugosa Thunb., Salix sp., Sorbus acuparia L., Spirea nipponica Maxim., Viburnum sp., V. rhytidophyllum Hemsl. Feeding on aphids. April, May, June, Adalia bipunctata was the fourth most common species found in the survey (47 locations, five commonly).

Anatis labiculata (Say): Anne Arundel, Baltimore, Carroll, Howard, Montgomery,

Prince George's. Acer rubrum, A. saccharum Marsh., Betula sp., Cornus florida L., Ligustrum sp., Pyracantha sp., Quercus sp., Q. acutissima Carruth., Tsuga canadensis (L.) Carriere, Ulmus sp., Viburnum sp. April, May, June. Thirteen locations.

A. mali (Say): Harford, Prince George's. Pinus sylvestris L. May. Two specimens.

Axion tripustulatum (DeGeer): Prince George's. Quercus sp. Associated with M. obscura. April. One specimen.

Coccinella novemnotata Herbst: Allegany, Carroll. June, July. Two locations. This species used to be very common in Maryland, judging by the number of specimens in student collections at the University of Maryland and observations by the senior author. Since the introduction of *C. septempunctata* L., the species has only been collected twice in 1986. Whether this is due to a natural decline or competitive displacement is not known at present.

C. septempunctata L.: Allegany, Anne Arundel, Baltimore, Carroll, Charles, Frederick, Harford, Howard, Kent, Montgomery, Prince George's, Queen Anne's, Somerset, St. Mary's, Talbot, Wicomico, Worcester. Acer saccharum, Betula sp., Crategus sp., Elageanus umbellata, Euonymus sp., Fraxinus sp., Hemerocallis sp., Ilex cornuta Lindl. & Paxt., Juniperus sp., Malus sp., Pieris japonica (Thunb.) D. Don, Pinus sp., P. sylvestris L., P. thunbergiana Franco, Prunus sp., Pyracantha sp., Pyrus calleryana Decne., Rhododendron sp., Rosa sp., Quercus sp., Q. robur, Spirea sp., Taxus sp., Tilia sp., Tsuga canadensis, Viburnum sp., Wisteria sp., Zelkova serrata (Thunb.) Makino. March, April, May, June, July, August, September, October, November, This was the most commonly collected species in the survey (87 locations). Feeding on aphids. Coccinella septempunctata was taken in large numbers on several occasions. On three separate occasions adult C. septempunctata were collected on the bark of cut Christmas trees of Picea abies. P. pungens Engelm., and Pinus sylvestris. The origin of these trees was eastern Pennsylvania. The movement of cut Christmas trees provides an interesting method of distribution for this species. This species has also been collected in a greenhouse complex.

Coleomegilla maculata lengi Timberlake: Anne Arundel, Baltimore, Caroline, Carroll, Kent, Prince George's, Queen Anne's, St. Mary's, Somerset, Talbot, Wicomico, Worcester. Acer rubrum, Coreopsis sp., Euonymus alata, E. japonica Thunb., Forsythia intermedia. Gleditsia tricanthos. Hedera helix L., Hibiscus sp., Ilex crenata, I. opaca Ait., Juglanus sp., Magnolia sp., Phlox sp., Pinus sp., P. sylvestris, Prunus sp., Pyrancantha sp., Rosa sp., R. rugosa, Quercus sp., Salix sp., Spirea sp., Taxus sp., Tsuga canadensis, Vitis sp., Z. serrata. Feeding on aphids; associated with Pineus strobi (Hartig) (Homoptera: Phylloxeridae) and Dymicoccus wisteriae (Green) (Pseudococcidae). April, May, June, July, August. This was the second most common species in this survey (45 locations).

Cycloneda munda (Say): Anne Arundel, Baltimore, Charles, Harford, Montgomery, Prince George's, Queen Anne's, Wicomico. Acer sp., Betula pendula Roth, Malus sp., Pyracantha sp., Quercus palustris Muenchh., Taxus sp. Feeding on aphids. May, June, July, August. Fifteen locations.

Olla v-nigrium (Mulsant): Anne Arundel, Prince George's, Talbot, Worcester. Magnolia sp., Ulmus sp. Associated with Toumeyella liriodendri (Gmelin) (Homoptera: Coccidae). May, June, September. Five locations.

Hippodamia convergens Guérin: Anne Arundel, Baltimore, Carroll, Frederick, Harford, Kent, Montgomery, Prince George's, St. Mary's, Washington, Wicomico, Worcester. Acer rubrum, Cornus florida, Elageanus umbellata, Euonymus alata, Fraxinus sp., Hedera helix, Hibiscus syriacus L., Hydrangea sp., Juglanus sp., Malus sp., Picea abies, Prunus sp., Pyracantha sp., Quercus sp., Rosa sp., Viburnum sp. Feeding on aphids. April, May, June, August.

VOLUME 92, NUMBER 2 313

This was the third most common species collected (39 locations).

H. glacialis (Fab.): Kent, Somerset. July. Two locations.

Mulsantia picta (Randall): Caroline, Howard, Kent. May, June. Three specimens.

Neoharmonia venusta venusta (Melsheimer): Anne Arundel. Salix sp. May. One specimen.

## Psylloborini

Psyllobora vigintimaculata (Say): Charles, Kent. May, September. Three locations.

#### ACKNOWLEDGMENTS

We would like to thank K. M. Berry, S. M. Bohlen, T. S. Creel, J. Fishback, K. Hogsten, D. C. Laughlin, L. C. Miles, S. C. Parry, J. P. Smokonich, S. J. Stevenson, J. A.

Wolinski, and M. E. Zastrow for their assistance in collecting specimens in the field.

We thank J. J. Drea, R. M. Hendrickson, and the rest of the staff of the Beneficial Insects Lab, USDA for supplying the *Chilocorus kuwanae* released in Maryland nurseries.

W. F. Gimpel, Jr., Maryland Department of Agriculture, and J. A. Davidson, University of Maryland, commented on earlier drafts of this manuscript.

Maryland Department of Agriculture Contribution Number 60-89.

#### LITERATURE CITED

Bailey, L. H. and E. Z. Bailey. 1976. Hortus Third. Macmillian Publ. Co., New York. 1290 pp.
Gordon, R. D. 1985. The Coccinellidae (Coleoptera) of America north of Mexico. J. New York Entomol. Soc. 93: 1–912.