



ZOOTAXA


3148

Animal biodiversity:

An outline of higher-level classification and survey of taxonomic richness

ZHI-QIANG ZHANG (ED.)

*New Zealand Arthropod Collection, Landcare Research, Private Bag 92170, Auckland, New Zealand;
zhangz@landcareresearch.co.nz*



Magnolia Press
Auckland, New Zealand

ZHI-QIANG ZHANG (ED.)

Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness
(*Zootaxa* 3148)

237 pp.; 30 cm.

23 Dec. 2011

ISBN 978-1-86977-849-1 (paperback)

ISBN 978-1-86977-850-7 (Online edition)

FIRST PUBLISHED IN 2011 BY

Magnolia Press

P.O. Box 41-383

Auckland 1346

New Zealand

e-mail: zootaxa@mapress.com

<http://www.mapress.com/zootaxa/>

© 2011 Magnolia Press

All rights reserved.

No part of this publication may be reproduced, stored, transmitted or disseminated, in any form, or by any means, without prior written permission from the publisher, to whom all requests to reproduce copyright material should be directed in writing.

This authorization does not extend to any other kind of copying, by any means, in any form, and for any purpose other than private research use.

ISSN 1175-5326 (Print edition)

ISSN 1175-5334 (Online edition)

Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness

ZHI-QIANG ZHANG (ED.)

Table of contents

- 7 **Animal biodiversity: An introduction to higher-level classification and taxonomic richness**
ZHI-QIANG ZHANG
- 13 **Phylum Porifera Grant, 1826. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness***
JOHN N.A. HOOPER, ROB W.M. VAN SOEST & ANDRZEJ PISERA
- 19 **Class Anthozoa Ehrenberg, 1834. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness***
ANDREA L. CROWTHER
- 24 **Phylum Xenacoelomorpha Philippe, *et al.*, 2011. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness***
SETH TYLER AND STEPHEN SCHILLING
- 26 **Pisces. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness***
WILLIAM N. ESCHMEYER & JON D. FONG
- 39 **Class Amphibia Gray, 182. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness***
DAVID C. BLACKBURN & DAVID B. WAKE
- 56 **Class Mammalia Linnaeus, 1758. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness***
D.E. WILSON & D.M. REEDER
- 61 **Order Testudines Batsch, 1788. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness***
UWE FRITZ
- 63 **“Phylum Nematoda Cobb 1932. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness***
MIKE HODDA
- 96 **Phylum Tardigrada Doyère, 1840. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness***
ROBERTO GUIDETTI & ROBERTO BERTOLANI
- 98 **Phylum Onychophora Grube, 1853. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness***
GEORG MAYER & IVO DE SENA OLIVEIRA
- 99 **Phylum Arthropoda von Siebold, 1848 *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness***
ZHI-QIANG ZHANG

- 104 **Class Trilobita Walch, 1771. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness***
JONATHAN M. ADRAIN
- 110 **Class Pycnogonida Latreille, 1810. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness***
ROGER BAMBER
- 112 **Order Opiliones Sundevall, 1833. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness***
ADRIANO B. KURY
- 115 **Order Scorpiones C.L. Koch, 1850. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness***
LORENZO PRENDINI
- 118 **Order Solifugae Sundevall, 1833. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness***
LORENZO PRENDINI
- 119 **Order Pseudoscorpiones de Geer, 1778. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness***
MARK HARVEY
- 121 **Order Palpigradi Thorell, 1888. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness***
LORENZO PRENDINI
- 122 **Order Ricinulei Thorell, 1876. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness***
LORENZO PRENDINI
- 123 **Superorder Parasitiformes Reuter, 1909. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness***
FRÉDÉRIC BEAULIEU
- 129 **Order Trombidiformes Reuter, 1909. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness***
ZHI-QIANG ZHANG, QING-HAI FAN, VLADIMIR PESIC, HARRY SMIT, ANDRE V. BOCHKOV, A. A. KHAUSTOV, ANNE BAKER, ANDREAS WOHLTMANN, TINGHUAN WEN, JAMES W. AMRINE, P. BERON, JIANZHEN LIN, GRZEGORZ GABRYS & ROBERT HUSBAND
- 139 **Suborder Endeostigmata Reuter, 1909. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness***
DAVID EVANS WALTER, SAMUEL BOLTON, MATTI UUSITALO & ZHI-QIANG ZHANG
- 141 **Suborder Oribatida van der Hammen, 1968. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness***
HEINRICH SCHATZ
- 149 **Order Araneae Clerck, 1757. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness***
JASON A. DUNLOP AND DAVID PENNEY

- 154 **Order Amblypygi Thorell, 1883. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness***
LORENZO PRENDINI
- 155 **Order Thelyphonida Latreille, 1804. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness***
LORENZO PRENDINI
- 156 **Order Schizomida Petrunkevitch, 1945. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness***
LORENZO PRENDINI
- 157 **Class Chilopoda, Class Symphyla and Class Pauropoda. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness***
ALESSANDRO MINELLI
- 159 **Class Diplopoda de Blainville in Gervais, 1844. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness***
WILLIAM SHEAR
- 165 **Subphylum Crustacea Brünnich, 1772. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness***
SHANE T. AHYONG, JAMES K. LOWRY, MIGUEL ALONSO, ROGER N. BAMBER, GEOFFREY A. BOXSHALL, PETER CASTRO, SARAH GERKEN, GORDAN S. KARAMAN, JOSEPH W. GOY, DIANA S. JONES, KENNETH MELAND, D. CHRISTOPHER ROGERS & JÖRUNDUR SVAVARSSON
- 192 **Class Collembola Lubbock, 1870. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness***
FRANS JANSSENS AND KENNETH A. CHRISTIANSEN
- 195 **Order Orthoptera Oliver, 1789. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness***
SIGFRID INGRISCH
- 198 **Order Phasmida Leach. 1815. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness***
PAUL D. BROCK & JUDITH MARSHALL
- 199 **Order Blattodea Brunner von Wattenwyl, 1882. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness***
GEORGE W. BECCALONI & PAUL EGGLETON
- 201 **Order Thysanoptera Haliday, 1836. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness***
LAURENCE A. MOUND
- 203 **Order Coleoptera Linnaeus, 1758. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness***
S. A. SLIPINSKI, R. A. B. LESCHEN, AND J. F. LAWRENCE
- 209 **Order Trichoptera Kirby, 1813. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness***
RALPH W. HOLZENTHAL, JOHN C. MORSE & KARL M. KJER

- 212 **Order Lepidoptera Linnaeus, 1758. In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness**
ERIK J. VAN NIEUKERKEN, LAURI KAILA, IAN J. KITCHING, NIELS P. KRISTENSEN, DAVID C. LEES, JOËL MINET, CHARLES MITTER, MARKO MUTANEN, JEROME C. REGIER, THOMAS J. SIMONSEN, NIKLAS WAHLBERG, SHEN-HORN YEN, REZA ZAHIRI, DAVID ADAMSKI, JOAQUIN BAIXERAS, DANIEL BARTSCH, BENGT Å. BENGTSSON, JOHN W. BROWN, SIBYL RAE BUCHELI, DONALD R. DAVIS, JURATE DE PRINS, WILLY DE PRINS, MARC E. EPSTEIN, PATRICIA GENTILI-POOLE, CEES GIELIS, PETER HÄTTENSCHWILER, AXEL HAUSMANN, JEREMY D. HOLLOWAY, AXEL KALLIES, OLE KARSHOLT, AKITO Y. KAWAHARA, SJAAK (J.C.) KOSTER, MIKHAIL V. KOZLOV, J. DONALD LAFONTAINE, GERARDO LAMAS, JEAN-FRANÇOIS LANDRY, SANGMI LEE, MATTHIAS NUSS, KYU-TEK PARK, CARLA PENZ, JADRANKA ROTA, ALEXANDER SCHINTLMEISTER, B. CHRISTIAN SCHMIDT, JAE-CHEON SOHN, M. ALMA SOLIS, GERHARD M. TARMANN, ANDREW D. WARREN, SUSAN WELLER, ROMAN V. YAKOVLEV, VADIM V. ZOLOTUHIN, ANDREAS ZWICK
- 222 **Order Diptera Linnaeus, 1758. In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness**
THOMAS PAPE, VLADIMIR BLAGODEROV & MIKHAIL B. MOSTOVSKI
- 230 **Phylum Phoronida Hatschek, 1888. In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness**
CHRISTIAN C. EMIG
- 231 **Phylum Rotifera Cuvier, 1817. In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness**
H. SEGERS
- 234 **Phylum Acanthocephala Kohlruther, 1771 In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness**
SCOTT MONKS AND DENNIS J. RICHARDSON



Animal biodiversity: An introduction to higher-level classification and taxonomic richness

ZHI-QIANG ZHANG

Landcare Research, Private Bag 92170, Auckland 1142, New Zealand; ZhangZ@landcareresearch.co.nz

Abstract

For the kingdom Animalia, 1,552,319 species have been described in 40 phyla in a new evolutionary classification. Among these, the phylum Arthropoda alone represents 1,242,040 species, or about 80% of the total. The most successful group, the Insecta (1,020,007 species), accounts for about 66% of all animals. The most successful insect order, Coleoptera (387,100 species), represents about 38% of all species in 39 insect orders. Another major group in Arthropoda is the class Arachnida (112,201 species), which is dominated by the mites and ticks (Acari 54,617 species) and spiders (43,579 species). Other highly diverse arthropod groups include Crustacea (66,914 species), Trilobitomorpha (19,606 species) and Myriapoda (11,885 species). The phylum Mollusca (117,358 species) is more diverse than other successful invertebrate phyla Platyhelminthes (29,285 species), Nematoda (24,783 species), Echinodermata (20,509 species), Annelida (17,210 species) and Bryozoa (10,941 species). The phylum Craniata, including the vertebrates, represents 64,832 species (for Recent taxa, except for amphibians); among these 7,694 described species of amphibians, 31,958 species of “fish” and 5,750 species of mammals.

Introduction

Discovering and describing how many species inhabit the Earth remains a fundamental quest of biology, even when we are entering the “phylogenomic age” in the history of taxonomy. With so many important issues facing us—invasive species, climate change, habitat destruction and loss of biodiversity in particular, the need for authoritative taxonomic information is higher than ever. *Zootaxa* has been a major force in describing world’s biodiversity in the last 10 years (Zhang 2011a). Most of the papers published in *Zootaxa* are descriptions of new taxa and taxonomic revisions interesting mainly to specialists. However, to reach a broader readership, we present, in this special volume, basic taxonomic data (classifications and diversity estimates) that are very important to the any user of biodiversity information. Extensions of results of taxonomic research to users, not only other taxonomists, but also all others, are important to achieve a greater impact for taxonomy.

The idea of this special volume was conceived in 2010—The International Year of Biodiversity. This volume is intended to be a collaborative effort by hundreds of taxonomists, each contributing a section on his/her group to the overall outline of the current, most-accepted taxonomy of the animal kingdom. The bulk of the volume will be a series of linear sequences of higher taxa (living and fossil, the latter to be indicated by a dagger (†)) from kingdom to family in a Linnaean hierarchy, using valid names following the International Code of Zoological Nomenclature. For each family, the best estimate of the number of described genera and species in the world is provided by specialists of the group. We want to present two types of important taxonomic information—how many species have been described and how they are classified—to the users of biodiversity information in an easily accessible volume, which is published for open access without cost to contributors.

Results

General account

Over 100 taxonomists contributed 43 outlines, some for phyla, some for classes and some for orders (see the outline below for notes and references to various outlines in this volume). About a quarter of the phyla were covered, with many gaps to be filled by specialists in future editions. During preparation of the volume, it quickly became evident that for many groups, there are different classifications for fossil and Recent taxa, and there is little dialogue among taxonomists working on Recent species and those on fossil. A good example is the outline on Porifera by Hooper *et al.* (2011), who tried to bring these together. The diversity estimates for many groups are deficient in fossil counts, and the total is therefore underestimated. We invite taxonomists working on both Recent and fossil taxa to joint efforts in providing a more complete outline of animal classification in future editions.

For the kingdom Animalia, 1,552,319 species have been described in 40 phyla (see the list below). Among these, the phylum Arthropoda alone represents 1,242,040 species, or about 80% of the total. The most successful group, the Insecta (1,020,007 species), accounts for about 66% of all animals, or 82% of arthropods. The most successful insect order, Coleoptera (387,100 species), represents about 38% of all species in 39 insect orders. Another major group in Arthropoda is the class Arachnida (112,201 species), which is dominated by the mites and ticks (Acari 54,617 species) and spiders (43,579 species). Other highly diverse arthropod groups include Crustacea (66,914 species), Trilobitomorpha (19,606 species) and Myriapoda (11,885 species).

The phylum Mollusca (117,358 species) leads, by a significant margin, among other invertebrate phyla in diversity. Significant groups include: Platyhelminthes (29,285 species), Nematoda (24,783 species), Echinodermata (20,509 species), Annelida (17,210 species) and Bryozoa (10,941 species).

The phylum Craniata, including the vertebrates, represents 64,832 species (for Recent taxa, except for amphibians). Blackburn & Wake (2011; this volume) presented a new consensus classification for amphibians, with an estimate of 7,694 described species. Eschmeyer & Fong (2011) summarised data from the “Catalog of Fish” (31,958 species). Wilson & Reeder (2011, this volume) updated their list of 5,750 species of Mammalia.

An evo-Linnaean classification of animal phyla

There are different schemes of presenting a classification or translating a phylogeny into a classification. Without any phylogenetic information, the simplest way is to list all phyla by alphabetical order. The Catalogue of Life¹, for example, lists animal phyla alphabetically, with complete loss of phylogenetic information. Traditionally, taxonomists also list taxa of equal rank using a self-chosen sequence (if not alphabetical), with basal taxa listed first and most derived one at the end. With explicit phylogenetic information available, Hennig (1965) showed that a linear sequence using a combined successive alphabet/number prefix can fully represent phylogenetic relationships. A cladogram can be presented in a hierarchy with inclusions of less inclusive members into sets of more inclusive taxa, which are given higher ranks (e.g., Hennig 1966: fig. 18 and also Dubois 2006: fig. 3). Wiley (1979) proposed an annotated Linnaean hierarchy, with comments on natural taxa and competing systems, and this is applied, for example, to the list of Diptera in a hierarchical sequence (Pape *et al.* 2011, this volume), with taxa arranged more or less in phylogenetic sequence from the primitive (oldest) to the most advanced (youngest) taxa, and some clades not ranked. I, however, found the latter not the best way, in terms of both nomenclatural aspects and also the level of details in providing phylogenetic information at the level of terminal taxa.

The phylogeny of the animal phyla is in a flux (Edgecombe *et al.* 2011), with five competing hypotheses among five major basal taxa: Ctenophora, Porifera, Placozoa, Cnidaria and Bilateria. Thus any nomenclature and classifications at the levels of subkingdom to superphylum will be unstable due to the problems at the most basal positions. Supporters of the PhyloCode (De Queiroz & Gauthier, 1992) tend to name all hierarchical nodes of a tree and the shortage of Linnaean ranks to deal with this is one reason for the rankless systems of nomenclature and their uses in classification. However, not all nodes of such trees need to be named (Dubois 2006, 2007), and nomenclatural ranks can be used consistently to express the positions of taxonomic categories in a hierarchical classification. Herein, I used a new method to present the “evo-Linnaean” classification of animal phyla: (1) The essence of this system is to use commonly used Linnaean ranks as much as possible and combine each of these ranks with a successive numbering system². This system can easily cope with all levels in a phylogenetic tree.

1. Catalogue of Life: 5th December 2011 <http://www.catalogueoflife.org/> (accessed on 18 Dec. 2011)

Dubois (2006) proposed a detailed standard nomenclatural hierarchy which altogether allows for 209 potential ranks by using additional prefix beyond what is commonly used (e.g. super, sub and infra-). However, between two major ranks (e.g. Kingdom and Phylum), the number of ranks is limited and not sufficient to express all possible levels of relationships among phyla. (2) I refer to this as the “evo-Linnaean” classification to indicate that this classification aims at reflecting evolutionary relationship as much as possible to set it apart from the traditional Linnaean classification, which has a typological connotation. Taxa should be arranged from basal-most to the most derived group following the relationship in a chosen hypothesis or reference tree. (3) New names for hypothesised taxa that lack consistent support are minimised. (4) To follow, as much as possible, the Codes of nomenclature. Although, names of taxa above the family-group are not fully regulated in the ICZN, it is a good practice to use existing names for the same taxa (Dubois 2006). This system is also applied to an evo-Linnaean classification of Arthropoda (Zhang, 2011b, this volume).

Kingdom **Animalia** Linnaeus, 1758 (1,552,319 species; of which †24,659)³

Phylum 1 **Ctenophora** Eschscholtz, 1829 (242 species)⁴

Phylum 2.1 **Porifera** Grant, 1826 (8,346 species)⁵

Phylum 2.2 1 **Placozoa** Grell, 1971 (1 species)

Phylum 2.2.2.1 **Cnidaria** Hatschek, 1888 (10,105 species)⁶

Phylum 2.2.2.2 **Myxozoa** Grassé, 1970 (2,402 species)⁷ ⁸

Phylum 2.2.2.3.1.1 **Xenoturbellida** Westblad, 1949 (2 species)⁹

Phylum 2.2.2.3.1.2 **Acoelomorpha** Ehlers, 1985 (393 species)

Phylum 2.2.2.3.2.1 **Orthonectida** Giard, 1877 (43 species)¹⁰

Phylum 2.2.2.3.2.2 **Rhombozoa** van Beneden, 1876 (123 species)¹¹

Phylum 2.2.2.3.3.1.1.1 **Cephalochordata** Owen, 1846 (33 species)¹²

-
2. The International standard ISO 2145, which defines a typographic convention for the "numbering of divisions and subdivisions in written documents". It applies to any kind of document, including manuscripts, books, journal articles, and standards. It is commonly used in Table of Contents of books to express hierarchical structure. For example, Rasnitsyn & Quicke (2002) used the numbering system in the table of contents (pages vii–viii) to express the phylogenetic tree of Insecta (page 2).
 3. This evo-Linnaean classification of animal phyla general follows consensus phylogenetic relationships summarised in Edgecombe *et al.* (2011). The poorly known extinct phylum †Trilobozoa is of uncertain placement. Only Recent species were counted for many insect orders and all of Myriapoda; most vertebrates and many invertebrate phyla; so the total number of species should be considered incomplete, as diversity of fossil taxa is underestimated.
 4. Ctenophora was placed as sister group to the remaining animals (Porifera (Placozoa (Cnidaria, Bilateria))) in phylogenomic analyses by Dunn *et al.* (2008) and Hejnol *et al.* (2009). Diversity estimate based on WoRMS (2011). Ctenophora. In: Nicolas Bailly (2011). FishBase. Accessed through: World Register of Marine Species at <http://www.marinespecies.org/aphia.php?p=taxdetails&id=1248> on 2011-12-18.
 5. Hooper *et al.* (2011, this volume), but number of fossil species unknown, although an attempt was able to integrate the classification of Recent and fossil Porifera.
 6. Daly *et al.* (2007) included 5 classes: (1) Anthozoa with an estimate of 7,500 extant species, now updated by Crowther (2011, this volume) to 6,142 species; (2) Cubozoa 36 species, here updated to 42 species; (3) Hydrozoa 3500 species, updated to 3,643 species; (4) Scyphozoa 216 species, updated by 228 species (all updates by Zhang based on Zoological Record for new species since 2008); (5) Staurozoa 50 species.
 7. Based on Lom & Dykova (2006), with updates of new species described from 2006 using *Zoological Record*.
 8. Zrzavy *et al.* (1998) placed it within Cnidaria based on morphological and 18S ribosomal DNA evidence, recently, Evans *et al.* (2010), using phylogenomic and ribosomal data sets, showed existence of two relatively stable placements for myxozoans: within Cnidaria or the alternative hypothesis at the base of Bilateria. Cnidarian taxonomists currently do not consider myxozoans as cnidarians (Daly *et al.* 2007).
 9. See Tyler & Schilling (2011, this volume), who treat this phylum as a subphylum of Xenacoelomorpha, following Philippe *et al.* (2011), who proposed the name Xenacoelomorpha for Xenoturbellida + Acoelomorpha.
 10. Edgecombe *et al.* (2011) recognised two subgroups of Mesozoa, Orthonectida and Rhombozoa, as phyla but did not include them in the phylogenetic tree. The placement of Mesozoa here follows Noordijk *et al.* (2010). Diversity estimates based on Furuya, H & van der Land, J. (2011). Orthonectida. Accessed through: World Register of Marine Species at <http://www.marinespecies.org/aphia.php?p=taxdetails&id=14220> on 2011-12-17.
 11. Diversity estimates based on WoRMS (2011) Rhombozoa. Accessed through: World Register of Marine Species at <http://www.marinespecies.org/aphia.php?p=taxdetails&id=14219> on 2011-12-17.
 12. Based Chapman (2009).

Phylum 2.2.2.3.3.1.1.2.1 **Tunicata** Lamarck, 1816 (2,792 species)¹³
 Phylum 2.2.2.3.3.1.1.2.2 **Craniata** Linnaeus, 1758 (64,832 species)¹⁴
 Phylum 2.2.2.3.3.1.2 **Echinodermata** (20,509 species; of which †13,000)¹⁵
 Phylum 2.2.2.3.3.1.2.2 **Hemichordata** Bateson, 1885 (120 species)¹⁶
 Phylum 2.2.2.3.3.2.1 **Chaetognatha** Leuckart 1854 (186 species; of which †7)¹⁷
 Phylum 2.2.2.3.3.2.2.1.1 **Nematoda** Cobb, 1932 (24,783 species, of which †10)¹⁸
 Phylum 2.2.2.3.3.2.2.1.2 **Nematomorpha** Vejdovsky, 1886 (351 species)¹⁹
 Phylum 2.2.2.3.3.2.2.2 **Tardigrada** Doyère, 1840 (1,157 species)²⁰
 Phylum 2.2.2.3.3.2.2.3.1 **Onychophora** Grube, 1853 (182 species, †3)²¹
 Phylum 2.2.2.3.3.2.2.3.2 **Arthropoda** von Siebold, 1848 (1,242,040 species; †6,182)²²
 Phylum 2.2.2.3.3.2.2.4.1 **Priapulida** Théel, 1906 (19 species)²³
 Phylum 2.2.2.3.3.2.2.4.2 **Loricifera** Kristensen, 1983 (30 species)²⁴
 Phylum 2.2.2.3.3.2.2.4.3 **Kinorhyncha** Reinhard, 1881 (179 species)²⁵
 Phylum 2.2.2.3.3.2.3.1.1 **Bryozoa** Ehrenberg, 1831 (10,941 species; of which †5,455)²⁶
 Phylum 2.2.2.3.3.2.3.1.2.1 **Entoprocta** Nitsche 1869 (169 species)²⁷
 Phylum 2.2.2.3.3.2.3.1.2.2 **Cycliophora** Funch & Kristensen, 1995 (2 species)²⁸
 Phylum 2.2.2.3.3.2.3.2.1.1 **Annelida** Lamarck, 1809 (17,210 species)²⁹
 Phylum 2.2.2.3.3.2.3.2.1.2 **Spincula** Rafinesque, 1814 (1,507 species)³⁰
 Phylum 2.2.2.3.3.2.3.2.1.3 **Echiura** Newby, 1940 (236 species)³¹
 Phylum 2.2.2.3.3.2.3.2.2 **Mollusca** Linnaeus, 1758 (117,358 species)³²
 Phylum 2.2.2.3.3.2.3.2.3 **Nemertea** Schultzze, 1851 (1,200 species)³³
 Phylum 2.2.2.3.3.2.3.2.4.1 **Brachiopoda** Duméril, 1806 (443 species)³⁴
 Phylum 2.2.2.3.3.2.3.2.4.2 **Phoronida** Hatschek, 1888 (10 species)³⁵

13. Also as Urochordata. Diversity estimates based on Chapman (2009) with updates.
14. Total for Recent taxa, except for amphibians; (1) 31,958 “fish” species (Eschmeyer & Fong, 2011, this volume); (2) 7,694 species of Amphibia (Blackburn & Wake, 2011; this volume); (3) 5,750 species of Mammalia (Wilson & Reeder 2011, this volume); (4) 9,990 species of birds (Chapman 2009); (5) 9,413 species of reptiles based on <http://www.reptile-database.org/db-info/SpeciesStat.html> (1 August update); it should be noted that this database listed 327 species of Testudines, whereas only 317 species of Testudines in Fritz (2011, this volume).
15. Updated from Pawson (2001), who estimated 7,000 living and 13,000 fossil species.
16. Based on Chapman (2009) with updates.
17. Based on Noordijk *et al.* (2010) with updates of 2010 and 2011 new names in Zoological Record.
18. See Hodda (2011, this volume).
19. Based on Poinar (2007) with updates.
20. Guidetti & Bertolani (2011, this volume): Heterotardigrada 444 species, Eutardigrada 712 species, Mesotardigrada 1 species.
21. See Mayer & Oliveira (2011, this volume).
22. See Zhang (2011b, this volume).
23. After Neuhaus, B. (2011). Priapulida. Accessed through: World Register of Marine Species at <http://www.marinespecies.org/aphia.php?p=taxdetails&id=101063> on 2011-12-17.
24. Based on Chapman (2009) & Gad (2009a,b).
25. Based on Neuhaus *et al.* (2011).
26. Phil Bock (personal communication, 17 Aug. 2011).
27. Also as Kamptozoa; diversity estimates based on WoRMS (2011) Entoprocta. Accessed through: World Register of Marine Species at <http://www.marinespecies.org/aphia.php?p=taxdetails&id=1271> on 2011-12-09.
28. Based on Funch & Kristensen (1995), Obst *et al.* (2006).
29. Based on Chapman (2009) with updates of 2009–2011 new names in Zoological Record.
30. There are recent evidence that this phylum, along with, Echiura and Siboglinidae, is part of Annelida (Struck *et al.* 2011). Diversity estimate based on WoRMS (2011). Sipuncula. In: Saiz, J. (2011) World Sipuncula database. Accessed through: World Register of Marine Species at <http://www.marinespecies.org/aphia.php?p=taxdetails&id=1268> on 2011-12-17.
31. Diversity estimates based on WoRMS (2011). Echiura. Accessed through: World Register of Marine Species at <http://www.marinespecies.org/aphia.php?p=taxdetails&id=1269> on 2011-12-17.
32. Based on Noordijk *et al.* (2010) with updates of 2010 and 2011 new names in Zoological Record.
33. Based on Chapman (2009).
34. Based on WoRMS (2011). Brachiopoda. In: Emig, C.C. (Ed) (2011). World Brachiopoda database. Accessed through: World Register of Marine Species at <http://www.marinespecies.org/aphia.php?p=taxdetails&id=1803> on 2011-12-18.
35. See Emig (2011, this volume).

- Phylum 2.2.2.3.3.2.3.3.1 **Gastrotricha** Metschnikoff, 1864 (790 species)³⁶
 Phylum 2.2.2.3.3.2.3.3.2 **Platyhelminthes** Gegenbaur, 1859 (29,285 species)³⁷
 Phylum 2.2.2.3.3.2.3.3.3.1 **Gnathostomulida** Riedl, 1969 (109 species)³⁸
 Phylum 2.2.2.3.3.2.3.3.3.2 **Micrognathozoa** Kristensen & Funch, 2000 (1 species)³⁹
 Phylum 2.2.2.3.3.2.3.3.3.3.1 **Rotifera** Cuvier, 1817 (1,583 species)⁴⁰
 Phylum 2.2.2.3.3.2.3.3.3.3.2 **Acanthocephala** Koelreuther, 1771 (1,194 species, of which †2)⁴¹

Acknowledgements

I thank all the contributors of this special volume for their collaborative efforts and my colleague Dr Rich Leschen (Landcare Research) for review and comments. The author was supported by the New Zealand Foundation for Research, Science and Technology through backbone funding of the “Defining New Zealand’s Land Biota” programme.

References

- Blackburn, D.C. & Wake, D.B. (2011) Class Amphibia Gray, 1825. In: Zhang, Z.-Q. (Ed.) *Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness*. *Zootaxa*, 3148, 39–55.
- Chapman, A.D. (2009) Numbers of living species in Australia and the world. Second edition. – Australian Biodiversity Information Services, Toowoomba.
- Daly, M., M.R. Brugler, P. Cartwright, A.G. Collins, M.N. Dawson, S.C. France, C.S. McFadden, D.M. Opresko, E. Rodriguez, S. Romano & J. Stake. (2007) The phylum Cnidaria: A review of phylogenetic patterns and diversity 300 years after Linnaeus. In: Zhang, Z.-Q. & Shear, W.A. (Eds) (2007) *Linnaeus Tercentenary: Progress in Invertebrate Taxonomy*. *Zootaxa*, 1668, 127–182.
- De Queiroz, K. & Gauthier, J. (1994) Toward a phylogenetic system of biological nomenclature. *Trends in Ecology & Evolution*, 9, 27–31.
- Dubois, A. (2006) Proposed Rules for the incorporation of nomina of higher-ranked zoological taxa in the International Code of Zoological Nomenclature. 2. The proposed Rules and their rationale. *Zoosystema*, 28, 165–258.
- Dubois, A. (2007) Phylogeny, taxonomy and nomenclature: the problem of taxonomic categories and of nomenclatural ranks. *Zootaxa*, 1519, 27–68.
- Dunn, C.W., Hejnol, A., Matus, D.Q., Pang, K., Browne, W.E., Smith, S.A., et al. (2008). Broad phylogenomic sampling improves resolution of the animal tree of life. *Nature*, 452, 745–749.
- Edgecombe, G.D., Giribet, G., Dunn, C.W., Hejnol, A., Kristensen, R.M., Neves, R.C., Rouse, G.W., Worsaae, K. & Sørensen, M.V. (2011) Higher-level metazoan relationships: recent progress and remaining questions. *Organism Diversity and Evolution*, 11, 151–172.
- Emig, C.C. (2011) Phylum Phoronida Hatschek, 1888. In: Zhang, Z.-Q. (Ed.) *Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness*. *Zootaxa*, 3148, 230–230.
- Eschmeyer, W. & Fong, J. (2011) Pisces. In: Zhang, Z.-Q. (Ed.) *Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness*. *Zootaxa*, 3148, 26–38.
- Evans NM, Holder MT, Barbeitos MS, Okamura B. & Cartwright, P. (2010) The phylogenetic position of Myxozoa: exploring conflicting signals in phylogenomic and ribosomal data sets. *Mol Biol Evol.* 27(12), 2733–46
- Fritz, U. (2011) Order Testudines Batsch, 1788. In: Zhang, Z.-Q. (Ed.) *Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness*. *Zootaxa*, 3148, 61–62.
- Funch, P. & Kristensen, R.M. (1995). Cyclophora is a new phylum with affinities to Entoprocta and Ectoprocta. *Nature*, 378 (6558), 711–714.
- Gad, G. (2009a) A clearly identifiable postlarva in the life cycle of a new species of *Pliciloricus* (Loricifera) from the deep sea of the Angola Basin. *Zootaxa*, 2096, 50–81.

36. Based on Schwank & Bartsch (1990) with updates of new ones described since 1990.

37. Seth Tyler (Personal communication)

38. Based on WoRMS (2011). Gnathostomulida. Accessed through: World Register of Marine Species at <http://www.marinespecies.org/aphia.php?p=taxdetails&id=14262> on 2011-12-17.

39. Based Kristensen & Funch (2000), who originally proposed it as a class.

40. Segers (2011, this volume).

41. Edgecombe *et al.* (2011) subsumed this phylum into Rotifera, however, specialists of Rotifera and Acanthocephala have not yet fully accepted this (Segers 2011 & Monks 2011, this volume). Diversity estimate and classification, see Monks (2011).

- Gad, G. (2009b) *Culexiregiloricus*, a new genus of Nanaloricidae (Loricifera) from the deep sea of the Guinea Basin (Southeast Atlantic). *Zootaxa*, 2096, 33–49.
- Grimaldi, D. & Engel, M.S. (2005) *Evolution of the Insects*. Cambridge University Press, Cambridge, New York.
- Guidetti, R. & Bertolani, R. (2011) Phylum Tardigrada Doyère, 1840. In: Zhang, Z.-Q. (Ed.) *Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness*. *Zootaxa*, 3148, 96–97.
- Hejnol, A., Obst, M., Stamatakis, A., Ott, M., Rouse, G. W., Edgecombe, G. D., et al. (2009). Assessing the root of bilaterian animals with scalable phylogenomic methods. *Proceedings of the Royal Society, Series B*, 276, 4261–4270.
- Hennig, W. (1965) Phylogenetic systematics. *Annual Review of Entomology*, 10, 97–116.
- Hennig, W. (1966) *Phylogenetic Systematics*, translated by D. Davis and R. Zangerl.: University of Illinois Press, Urbana.
- Hodda, M. (2011) Phylum Nematoda Cobb, 1932. In: Zhang, Z.-Q. (Ed.) *Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness*. *Zootaxa*, 3148, 63–95.
- Hooper, J.N.A., Van Soest, R.W.M. & Pisera, A. (2011) Phylum Porifera Grant, 1826. In: Zhang, Z.-Q. (Ed.) *Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness*. *Zootaxa*, 3148, 13–18.
- Kristensen, R.M. & Funch, P. (2000) Micrognathozoa: A new class with complicated jaws like those of Rotifera and Gnathostomulida. *Journal of Morphology*, 246, 1–49.
- Linnaeus, C. (1758) *Systema Naturae per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis. Editio decima, reformata. Tomus I. Laurentii Salvii, Holmiae*, 828 pp.
- Lom, J. & Dykova, I. (2006) Myxozoan genera: definition and notes on taxonomy, life-cycle terminology and pathogenic species. *Folia Parasitologica*, 53, 1–36.
- Mayer, G. & Oliveira, I.S. (2011) Phylum Onychophora Grube, 1853. In: Zhang, Z.-Q. (Ed.) *Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness*. *Zootaxa*, 3148, 98–98.
- Monks, S. & Richardson, D.J. (2011) Phylum Acanthocephala Kohlreuther, 1771. In: Zhang, Z.-Q. (Ed.) *Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness*. *Zootaxa*, 3148, 234–237.
- Neuhaus, B., Higgins, R.P. & Paavo, B. (2011) Ten Phylum Kinoryncha. In: Gordon, D.P. (ed.), *New Zealand inventory of biodiversity. Volume 2. Kingdom Animalia. Chaetognatha, Ecdysozoa, ichnofossils*. Canterbury University Press, Christchurch, New Zealand, pp. 50–89.
- Noordijk, J., van Loon, A.J., Kleukers, R.M.J.C. & Nieuwerkerken, E.J. van (2010) De Nederlandse biodiversiteit. *Nederlandse Fauna*, 10, 1–460.
- Obst, M., Funch, P. & Kristensen R.M. (2006) A new species of Cyclophora from the mouthparts of the American lobster, *Homarus americanus* (Nephropidae, Decapoda). *Organisms Diversity and Evolution*, 6 (2), 83–97.
- Pape, T., Blagoderov, V. & Mostovski, M.B. (2011, this volume) Order Diptera Linnaeus, 1758. In: Zhang, Z.-Q. (Ed.) *Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness*. *Zootaxa*, 3148, 222–229.
- Pawson, D.L. (2007) Phylum Echinodermata. *Zootaxa*, 1668, 749–764.
- Philippe, H., Brinkmann, H., Copley, R.R., Moroz, L.L., Nakano, H., Poustka, A.J., Wallberg, A., Peterson, K.J. & Telford, M.J. (2011) Acoelomorph flatworms are deuterostomes related to *Xenoturbella*. *Nature*, 470, 255–258.
- Poinar Jr., G. (2008) Global diversity of hairworms (Nematomorpha: Gordiaceae) in freshwater. *Hydrobiologia*, 595 (1), 79–83.
- Rasnitsyn, A.P. & Quicke D.L.J. (eds) (2002) *History of Insects*. Kluwer Academic Publishers, Dordrecht, Boston, London, 517 pp.
- Schwank, P. & Bartsch, I. (1990) Gastrotricha und Nemertini. *Süßwasserfauna von Mitteleuropa* 3/1+2, 1–258.
- Segers, H. (2011) Phylum Rotifera Cuvier, 1817. In: Zhang, Z.-Q. (Ed.) *Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness*. *Zootaxa*, 3148, 231–233.
- Struck, T.H., Paul, C., Hill, N., Hartmann, S., Hösel, C., Kube, M., Lieb, B., Meyer, A., Tiedemann, R., Purschke, G. & Bleidorn, C. (2011) Phylogenomic analyses unravel annelid evolution. *Nature*, 471 (7336), 95–98.
- Tyler, S. & Schilling, S. (2011b) Phylum Xenacoelomorpha Philippe, et al., 2011. In: Zhang, Z.-Q. (Ed.) *Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness*. *Zootaxa*, 3148, 24–25.
- Wiley, E.O. (1979) An annotated Linnaean hierarchy, with comments on natural taxa and competing systems. *Systematic Zoology*, 28, 308–337.
- Wilson, D.E. & Reeder, D.A. (2011) Class Mammalia Linnaeus, 1758. In: Zhang, Z.-Q. (Ed.) *Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness*. *Zootaxa*, 3148, 56–60.
- Zhang, Z.-Q. (2011a) Accelerating biodiversity descriptions and transforming taxonomic publishing: the first decade of *Zootaxa*. *Zootaxa*, 2896, 1–7.
- Zhang, Z.-Q. (2011b) Phylum Arthropoda von Siebold, 1848. In: Zhang, Z.-Q. (Ed.) *Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness*. *Zootaxa*, 3148, 99–103.
- Zrzavy, J., Mihulka, S., Kepka, P., Bezdek, A. & Tietz, D. (1998) Phylogeny of the Metazoa based on morphological and 18S ribosomal DNA evidence. *Cladistics*, 14, 249–285.

Phylum **Arthropoda** von Siebold, 1848¹ (1,242,040 species, of which †27,745)²
 Subphylum 1 †**Trilobitomorpha** Størmer, 1944 (19,606 species)³
 Subphylum 2 **Chelicerata** Heymons, 1901 (113,894 species, of which 1,957)⁴
 Class 1 **Pycnogonida** Latreille, 1810 (1330 species, of which †8 species)⁵
 Class 2 †**Aglaspidida** Walcott, 1911 (†11 species)⁶
 Class 3.1 †**Xiphosura** Latreille, 1802 (†98 species)⁷
 Class 3.2.1 †**Eurypterida** Burmeister, 1843 (†246 species)⁸
 Class 3.2.2 †**Chasmataspida** Caster & Brooks, 1956 (†8 species)⁹
 Class 3.3 **Arachnida** Cuvier, 1812 (112,201, of which †1,586)¹⁰
 Order 1.1 **Opiliones** Sundevall, 1833 (6,519 species, of which †35 species)¹¹
 Order 1.2 **Scorpiones** C.L. Koch, 1851 (2,068 species, of which †121 species)¹²
 Order 2.1 **Solifugae** Sundevall, 1833 (1,116 species, of which †3 species)¹³
 Order 2.2 **Pseudoscorpiones** de Geer, 1778 (3,494 species, of which †40 species)¹⁴
 Order 3 **Palpigradi** Thorell, 1888 (83 species, of which †1 species)¹⁵
 Order 4 †**Phalangiotarbida** Haase, 1890 (†31 species)¹⁶
 Order 5 **Ricinulei** Thorell, 1876 (73 species, of which †15 species)¹⁷
 Order 6.1 **Opilioacarida** Zakhvatkin, 1952 (37 species, of which †2 species)¹⁸
 Order 6.2 **Holothyrida** Thon, 1905 (27 species)
 Order 6.3 **Ixodida** Leach, 1815 (896 species, of which †5 species)¹⁹
 Order 6.4 **Mesostigmata** G. Canestrini, 1891 (11,424 species)
 Order 7.1 **Trombidiformes** Reuter, 1909 (25,821 species, of which †24 species)²⁰
 Order 7.2 **Sarcoptiformes** Reuter, 1909 (16,412 species, of which †113 species)²¹
 Order 8.1 †**Trigonotarbita** Petrunkevitch, 1949 (†65 species)²²
 Order 8.2.1.1 †**Uraraneida** Selden & Shear in Selden *et al.*, 2008 (†2 species)²³
 Order 8.2.1.2 **Araneae** Clerck, 1757 (43,579 species, of which †1,106 species)²⁴
 Order 8.2.2.1 †**Haptopoda** Pocock, 1911 (†1 species)²⁵

1. **BY** Zhi-Qiang Zhang (for full author addresses, see **Author name and address** after **References**). The title of this contribution should be cited as “Phylum Arthropoda von Siebold, 1848 *In*: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness”. This classification at the subphylum level generally reflects the high level relationship summarized by Giribet & Edgecombe (2012: Fig. 5), with modification and updates as indicated (see notes on Crustacea).
 Fossil Marellomorpha not included in the phylogenetic relationships focused on Recent taxa in Giribet & Edgecombe (2012). Grimaldi & Engel (2005) listed three genera.
2. Only Recent species were counted for many insect orders and all of Myriapoda; so the total number of species should be considered incomplete, as diversity of fossil taxa is underestimated.
3. See Adrain (2011, this volume).
4. There are broad consensus of high level relationship within Chelicerata (Dunlop 2010; Giribet & Edgecombe 2011); this classification at the class level generally reflects the high level relationship summarized by Giribet & Edgecombe (2011: Fig. 5), with modification and updates as indicated.
 Grimaldi & Engel (2005) listed three poorly known arthropod taxa (†Sidneyida, †Emeraldellida Størmer, 1944 and †Sanctacarida); their placement is uncertain in the current system (e.g. Dunlop 2010; Giribet & Edgecombe 2012).
5. See Bamber (2011, this volume).
6. Data from J. Ortega-Hernandez (personal communication, 15 Dec. 2011).
7. Fossil horseshoe crabs; based on Dunlop *et al.* (2011).
8. Fossil sea scorpions; based on Dunlop *et al.* (2011).
9. Based on Dunlop *et al.* (2011).
10. Arachnid classification reflects mostly the phylogenetic hypothesis by Shultz (2007), but his Acaromorpha is now disputed (see Dunlop 2010).
11. See Kury (2011, this volume).
12. See Prendini (2011a, this volume).
13. See Prendini (2011b, this volume).
14. See Harvey (2011, this volume).
15. See Prendini (2011c, this volume).
16. Based on Dunlop *et al.* (2011).
17. See Prendini (2011d, this volume).
18. See Beaulieu *et al.* (2011, this volume), with a list of families and diversity estimates for each family within the mite superorder Parasitiformes, including Opilioacarida, Holothyrida, Ixodida and Mesostigmata.
19. Based on Guglielmone *et al.* (2010).
20. See Zhang *et al.* (2011, this volume). The second mite superorder, Acariformes, consists of Trombidiformes and Sarcoptiformes.
21. This order includes two suborders: Oribatida and Endeostigmata. According to Schatz *et al.* (2011, this volume), Recent Oribatida (including Astigmata) currently include 249 families, 2399 genera and 16197 species; exclusively fossil taxa include 2 families, 20 genera and 108 species; Walter *et al.* (2011, this volume) estimated 10 families, 27 genera, 108 species for Endeostigmata, of which 3 genera and 5 species are based on fossil.
22. Based on Dunlop *et al.* (2011).
23. Selden *et al.* (2008).
24. See Dunlop & Penney (2011, this volume).

- Order 8.2.2.2.1 **Amblypygi** Thorell, 1883 (170 species, of which †9 species)²⁶
 Order 8.2.2.2.2.1 **Thelyphonida** Latreille, 1804 (119 species, of which †9 species)²⁷
 Order 8.2.2.2.2.2 **Schizomida** Petrunkevitch, 1945 (264 species, of which †4 species)²⁸
 Subphylum 2.1 **Myriapoda** Latreille, 1802 (11,885 species)²⁹
 Class 1 **Chilopoda** Latreille, 1817 (3,100 species)³⁰
 Class 2.1 **Symphyla** Ryder, 1880 (197 species)³¹
 Class 2.2.1 **Pauropoda** Lubbock, 1868 (835 species)
 Class 2.2.2 **Diplopoda** de Blainville in Gervais, 1844 (7,753 species)
 Subphylum 2.2.1 **Crustacea** Brünnich, 1772 (66,914 species)³²
 Subphylum 2.2.2 **Hexapoda** Latreille, 1825 (1,029,741 species, of which †6,182 species)³³
 Class 1.1 **Collembola** Lubbock, 1870 (8,130 species)³⁴
 Class 1.2.1 **Protura** Silvestri, 1907 (804 species)³⁵
 Class 1.2.2 **Diplura** Börner, 1904 (800 species)³⁶
 Class 2 **Insecta** Linnaeus, 1758 (1,020,007 species, of which †6,182 species)³⁷
 Order 1 **Archaeognatha** (513 species)³⁸
 Order 2.1 **Zygentoma** Börner, 1904 (561 species, of which †1 species)³⁹
 Order 2.2.1.1 **Ephemeroptera** Hyatt & Arms, 1890 (3,240 species)⁴⁰
 Order 2.2.1.2.1.1 †**Geroptera** Brodsky, 1994 (†2 species)⁴¹
 Order 2.2.1.2.1.2.1 †**Protodonata** (†42 species)⁴²
 Order 2.2.1.2.1.2.2 **Odonata** Fabricius, 1792 (5,899 species)⁴³
 Order 2.2.1.2.2.1.1 †**Palaeodictyoptera** Goldenberg, 1877 (†115 species)⁴⁴
 Order 2.2.1.2.2.2.1 **Mischoptera** Handlirsch, 1906 (†100 species)⁴⁵
 Order 2.2.1.2.2.2.2 †**Diaphanopteroidea** Handlirsch, 1906 (†50 species)⁴⁶
 Order 2.2.2.1 †**Paoliida** Handlirsch, 1906 (†12 species)⁴⁷
 Order 2.2.2.2.1 †**Caloneuroidea** Martynov, 1938 (†14 species)⁴⁸
 Order 2.2.2.2.2 †**Titanoptera** Sharov, 1968 (†15 species)⁴⁹

25. Based on Dunlop *et al.* (2011).
 26. See Prendini (2011e, this volume).
 27. See Prendini (2011f, this volume).
 28. See Prendini (2011g, this volume).
 29. Recent species only; many fossil species were described but were not included in this edition due to lack of time, so the total number of species should be considered incomplete.
 30. See Shear (2011, this volume).
 31. See Minelli (2011, this volume), which also covers Pauropoda and Diplopoda.
 32. See Ahyong *et al.* (2011, this volume), which provided a classification of all families and an estimate of 1,003 families, 9,522 genera and 66,914 species for Recent Crustacea. Crustacea is paraphyletic according to Giribet & Edgecombe (2011). Here traditional consensus classification presented in Ahyong *et al.* (2011) is followed.
 33. Classification reflects consensus phylogenetic relationship in Trautwein *et al.* (2012); inclusion of fossil orders follows Grimaldi & Engel (2005).
 34. See Janssens & Christiansen (2011, this volume).
 35. Based on Szeptycki (2007), and updated using Zoological Records from 2007.
 36. This number is cited by Chapman (2009) and also Tree of Life Web Project. 1995. Diplura. Version 01 January 1995 (temporary). <http://tolweb.org/Diplura/8204/1995.01.01> in The Tree of Life Web Project, <http://tolweb.org/>.
 37. Diversity estimates for fossil insects are not available for many orders and thus total diversity is underestimated.
 38. Based on Footitt & Adler (2009) and updated using Zoological Records from 2008.
 39. Alternative name Thysanura; estimates based Footitt & Adler (2009) and updated using Zoological Records from 2008. Fossil taxa are represented in at least 1 family (Data from Paleobiology Database on 15 December, 2011 using “Taxon Count” search)—note that these data may be incomplete.
 40. Based on Footitt & Adler (2009) and updated using Zoological Records from 2008; Footitt & Adler listed 3,046 species.
 41. Fossil taxa are represented in 2 genera and 1 family (Data from Paleobiology Database on 15 December, 2011 using “Taxon Count” search)—note that these data may be incomplete; Grimaldi & Engel (2005) mentioned “a few species” for this order.
 42. Fossil taxa are represented in 27 genera and 5 families (Data from Paleobiology Database on 15 December, 2011 using “Taxon Count” search)—note that these data are incomplete. Zoological Record lists a total of 48 species/subspecies names.
 43. Based on Footitt & Adler (2009) and updated using Zoological Records from 2008; Footitt & Adler listed 5,680 species; Trueman (2007) mentioned “around 6,000 species” for Odonata.
 44. Fossil taxa are represented in at least 82 genera and 25 families (Data from Paleobiology Database on 15 December, 2011 using “Taxon Count” search)—note that these data are incomplete.
 45. Concept and estimates follow Rasnitsyn & Quicke (2002); this order includes two orders listed in Grimaldi & Engel (2005): †Dipliptera (=Archodonata Martynov, 1932) and †Megasecoptera Brongniart, 1885.
 46. Based on Rasnitsyn & Quicke (2002).
 47. Based on Prokop & Nel (2007).
 48. Fossil taxa are represented in at least 11 genera and 7 families (Data from Paleobiology Database on 15 December, 2011 using “Taxon Count” search)—note that these data are incomplete; a total of 25 genera 9 families in David Eades. Polyneoptera Species File Online. Version 1.0/4.0. [retrieval date]. <http://Polyneoptera.SpeciesFile.org>, but Eades did not provide species for some genera.

- Order 2.2.2.2.3 **Orthoptera** Olivier, 1789 (24,276 species, of which †421 species)⁵⁰
 Order 2.2.2.3.1 **Phasmida** Leach, 1815 (3,029 species, of which †15 species)⁵¹
 Order 2.2.2.3.2 **Embioptera** Lameere, 1900 (464 species⁵², of which †1 species)
 Order 2.2.2.4.1 **Grylloblattodea** Brues & Melander, 1932 (34 species)⁵³
 Order 2.2.2.4.2 **Mantophasmatodea** Zompro, Klass, Kristensen & Adis, 2002 (21 species, of which †6 species)⁵⁴
 Order 2.2.2.5.1 **Plecoptera** Latreille 1802 (3,788 species⁵⁵, of which †45 species)⁵⁶
 Order 2.2.2.5.2 **Dermaptera** deGeer, 1773 (1,978 species)⁵⁷
 Order 2.2.2.6.1 **Zoraptera** Silvestri, 1913 (37 species)⁵⁸
 Order 2.2.2.6.2.1 **Mantodea** Burmeister 1838 (2,400 species)⁵⁹
 Order 2.2.2.6.2.2 **Blattodea** Brunner, 1882 (7,314 species)⁶⁰
 Order 2.2.3.1 †**Miomoptera** Martynov 1927 (†47 species)⁶¹
 Order 2.2.3.2.1 **Psocoptera** Shipley, 1904 (5,720 species)⁶²
 Order 2.2.3.2.2 **Phthiraptera** Haeckel, 1896 (5,102 species)⁶³
 Order 2.2.3.3.1 **Thysanoptera** Haliday, 1836 (6,019 species, of which †155 species)⁶⁴
 Order 2.2.3.3.2 **Hemiptera** Linnaeus 1758 (103,590)⁶⁵
 Order 2.2.4 †**Glosselytrodea** Martynov 1938 (†30 species)⁶⁶
 Order 2.2.5.1 **Hymenoptera** Linnaeus 1758⁶⁷ (116,861 species)⁶⁸
 Order 2.2.5.2.1.1.1 **Strepsiptera** Kirby 1813 (609 species)⁶⁹
 Order 2.2.5.2.1.1.2 **Coleoptera** Linnaeus, 1758 (387,100 species, of which †600 species)⁷⁰
 Order 2.2.5.2.1.2.1 **Neuroptera** Linnaeus, 1758 (5,868 species)⁷¹
 Order 2.2.5.2.1.2.2.1 **Megaloptera** Latreille, 1802 (354 species)⁷²
 Order 2.2.5.2.1.2.2.2 **Raphidioptera** Navás, 1916 (254 species)⁷³
 Order 2.2.5.2.2.1.1 **Trichoptera** Kirby, 1813 (14,999 species, of which †608 species)⁷⁴
 Order 2.2.5.2.2.1.2 **Lepidoptera** Linnaeus, 1758 (157,424 species, of which †86 species)⁷⁵
 Order 2.2.5.2.2.2.1 **Diptera** Linnaeus, 1758 (159,294 species, of which †3,817 species)⁷⁶
 Order 2.2.5.2.2.2.2.1 **Siphonaptera** Latreille, 1825 (2,075 species)⁷⁷
 Order 2.2.5.2.2.2.2.2 **Mecoptera** Packard, 1886 (757 species)⁷⁸

49. Based on Shcherbakov (2011).
 50. See Ingrisch (2011, this volume).
 51. See Brock and Marshall (2011, this volume) for Recent taxa only. Fossil taxa are represented in at least 14 genera and 9 families (Data from Paleobiology Database on 15 December, 2011 using “Taxon Count” search)—note that these data are incomplete.
 52. Based on Foottit & Adler (2009) and updated using Zoological Records from 2008.
 53. Grimaldi & Engel (2005) mentioned 26 species placed in 5 genera in 1 family.
 54. Updated with Arillo & Engel (2006), Eberhard *et al.* (2011). Note that this order and Grylloblattodea were placed in the order Notoptera by Arillo & Engel (2006).
 55. Based on Foottit & Adler (2009) and updated using Zoological Records from 2008; Foottit & Adler listed 3497 species.
 56. Fossil taxa are represented in at least 22 genera and 10 families (Data from Paleobiology Database on 15 December, 2011 using “Taxon Count” search)—note that these data are incomplete.
 57. Based on Foottit & Adler (2009) and updated using Zoological Records.
 58. Based on Foottit & Adler (2009) and updated using Zoological Records.
 59. Based on Otte, D., Spearman, L. & Martin, B.D.S. Mantodea Species File Online. Version 1.0/4.0. [retrieval date 15 Dec. 2011]. <<http://Mantodea.SpeciesFile.org>>.
 60. See Beccaloni & Eggleton (2011, this volume), without fossils; including 2,692 species of termites (no longer an order).
 61. Data from Paleobiology Database on 15 December, 2011 using “Taxon Count” search; this should be viewed as incomplete.
 62. Based on Foottit & Adler (2009) and updated using Zoological Records from 2008.
 63. Based on Foottit & Adler (2009) and updated using Zoological Records from 2008.
 64. See Mound *et al.* (2011, this volume).
 65. Based on Foottit & Adler (2009) and updated using Zoological Records from 2008.
 66. Based on Hong (2007) with updates.
 67. Holometabola for 11 orders from Hymenoptera to Mecoptera.
 68. Based on Hymenoptera Online <http://hol.osu.edu/> (accessed 15 Dec. 2011), which placed these species in 7,745 genera and 118 families. Fossil taxa are represented in at least 789 genera and 89 families (Data from Paleobiology Database on 15 December, 2011 using “Taxon Count” search)—note that these data are incomplete. Foottit & Adler (2009), citing Huber in that volume, mentioned 144,695 species.
 69. Based on Foottit & Adler (2009) and updated using Zoological Records from 2008.
 70. See Slipinski *et al.* (2011, this volume).
 71. Based on Foottit & Adler (2009) and updated using Zoological Records.
 72. Based on Foottit & Adler (2009) and updated using Zoological Records from 2008; Foottit & Adler listed 328 species.
 73. Based on Foottit & Adler (2009) and updated using Zoological Records from 2008.
 74. See Holzenthal *et al.* (2011, this volume).
 75. See Nieuwerkerken *et al.* (2011, this volume).
 76. See Pape *et al.* (2011, this volume), but 5,969 dubious species not included in total count.
 77. Based on Foottit & Adler (2009) and updated using Zoological Records.
 78. Based on Foottit & Adler (2009) and updated using Zoological Records from 2008.

Acknowledgements

I thank Dr Rich Leschen (Landcare Research) and Dr Qing-Hai Fan (Biosecurity New Zealand) for reviews and comments. The author was supported by the New Zealand Foundation for Research, Science and Technology through backbone funding of the “Defining New Zealand’s Land Biota” programme.

References

- Adrain, J.M. (2011) Class Trilobita Walch, 1771. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. Zootaxa*, 3148, 104–109.
- Ahyong, S.T., Lowry, J.K., Alonso, M., Bamber, R.N., Boxshall, G.A., Castro, P., Gerken, S., Karaman, G.S., Goy, J.W., Jones, D.S., Meland, K., Rogers, D.C. & Svavarsson, J. (2011) Order Scorpiones C.L. Koch, 1850. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. Zootaxa*, 3148, 165–191.
- Arillo, A. & Engel, M.S. (2006) Rock crawlers in Baltic amber (Notoptera: Mantophasmatodea). *American Museum Novitates*, 3539, 1–10.
- Aristov, D.S. (2009) A new family of the order Grylloblattida (Insecta) from the Middle Permian of Russia. *Paleontologicheskii Zhurnal*, (2), 59–63.
- Beccaloni, G.W. & Eggleton, P. (2011) Order Blattodea Brunner von Wattenwyl, 1882. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. Zootaxa*, 3148, 199–220.
- Bamber, R.N. (2011) Class Pycnogonida Latreille, 1810. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. Zootaxa*, 3148, 110–111.
- Beaulieu, F., Dowling, A.P.G., Klompen, H., Moraes, G.J. de & Walter, D.E. (2011) Superorder Parasitiformes Reuter, 1909. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. Zootaxa*, 3148, 123–128.
- Brock, P. & Marshall, J. (2011) Phasmida Leach, 1815. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. Zootaxa*, 3148, 198–198.
- Chapman, A.D. (2009) Numbers of living species in Australia and the world. Second edition. – Australian Biodiversity Information Services, Toowoomba.
- Eberhard, M.J.B., Picker, M.D. & Klass, K.-D. (2011) Sympatry in Mantophasmatodea, with the description of a new species and phylogenetic considerations. *Organisms Diversity & Evolution*, 11(1), 43–59.
- Dunlop J.A. (2010) Geological history and phylogeny of Chelicerata. *Arthropod Structure & Development*, 39, 124–142.
- Dunlop J.A. & Penney, D. (2011) Order Araneae Clerck, 1757. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. Zootaxa*, 3148, 149–153.
- Dunlop J.A., Penney D. & Jekel D. (2011) A summary list of fossil spiders and their relatives. *In: Platnick, N. I. (ed.) The world spider catalog, version 12.0. American Museum of Natural History, online at <http://research.amnh.org/entomology/spiders/catalog/index.html> DOI: 10.5531/db.iz.0001.*
- Footitt, R & Adler, P. (eds) (2009) *Insect Biodiversity: Science and Society*. Blackwell Publishing, London, 623 pp.
- Giribet, G. & Edgecombe, G.D. (2012) Reevaluating the Arthropod Tree of Life. *Annual Review of Entomology*, 57, 167–186.
- Grimaldi, D. & Engel, M.S. (2005) *Evolution of the Insects*. Cambridge University Press, Cambridge, New York.
- Guglielmone, A.A., Robbins, R.G., Apaneskevich, D.A., Petney, T.N., Estrada-Peña, A., Horak, I.G., Shao, R. & Barker, S. (2010) The Argasidae, Ixodidae and Nuttalliellidae (Acari: Ixodida) of the world: a list of valid species names. *Zootaxa*, 2528, 1–28.
- Harvey, M.S. (2011) Order Pseudoscorpiones de Geer, 1778. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. Zootaxa*, 3148, 119–120.
- Hong, Y.C. (2007) Discovery of the fossil glosselytrods (Insecta: Glosselytrodea) from Shaanxi, China. *Acta Entomologica Sinica*, 50(3), 271–280.
- Holzenthal, R.W., Morse, J.C. & Kjer, R.J., (2011) *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. Zootaxa*, 3148, 209–211.
- Ingrisch, S. (2011) Order Orthoptera Oliver, 1789. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. Zootaxa*, 3148, 195–197.
- Janssens, F. & Christiansen, K.A. (2011) Class Collembola Lubbock, 1870. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. Zootaxa*, 3148, 192–194.
- Kury, A.B. (2011) Order Opiliones Sundevall, 1833. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. Zootaxa*, 3148, 112–114.
- Mendes, L.F. (2002) Taxonomy of Zygentoma and Microryphia: historical overview, present status and goals for the new millennium. Proc. Xth International Colloquium on Apterygota, České Budějovice 2000: Apterygota at the Beginning of the Third Millennium. Elsevier GmbH.
- Minelli, A. (2011a) Class Trilobita, Class Symphyla and Class Pauropoda. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. Zootaxa*, 3148, 157–158.
- Mound, L.A. (2011) Order Thysanoptera Haliday, 1836. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. Zootaxa*, 3148, 201–202.
- Nieukerken, E.J. van, Kaila, L., Kitching, I.J., Kristensen, N.P., Lees, D.C., Minet, J., Mitter, C., Mutanen, M., Regier, J.C., Simonsen, T.J., Wahlberg, N., Yen, S.-H., Zahiri, R., Adamski, D., Baixeras, J., Bartsch, D., Bengtsson, B.Å., Brown, J.W., Bucheli, S.R.,

- Davis, D.R., De Prins, J., De Prins, W., Epstein, M.E., Gentili-Poole, P., Gielis, C., Hättenschwiler, P., Hausmann, A., Holloway, J.D., Kallies, A., Karsholt, O., Kawahara, A.Y., Koster, S. (J.C.), Kozlov, M.V., Lafontaine, J.D., Lamas, G., Landry, J.-F., Lee, S., Nuss, M., Park, K.-T., Penz, C., Rota, J., Schintlmeister, A., Christian Schmidt, B., Sohn, J.-C., Solis, M.A., Tarmann, G.M., Warren, A.D., Weller, S., Yakovlev, R.V., Zolotuhin, V.V. & Zwick, A. (2011) Order Lepidoptera Linnaeus, 1758. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. Zootaxa*, 3148, 212–221.
- Pape, T., Blagoderov, V. & Mostovski, M.B. (2011, this volume) Order Diptera Linnaeus, 1758. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. Zootaxa*, 3148, 222–229.
- Prendini, L. (2011a) Order Scorpiones C.L. Koch, 1850. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. Zootaxa*, 3148, 115–117.
- Prendini, L. (2011b) Solifugae Sundevall, 1833. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. Zootaxa*, 3148, 118–118.
- Prendini, L. (2011c) Palpigradi Thorell, 1888. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. Zootaxa*, 3148, 121–121.
- Prendini, L. (2011d) Ricinulei Thorell, 1876. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. Zootaxa*, 3148, 122–122.
- Prendini, L. (2011e) Amblypygi Thorell, 1883. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. Zootaxa*, 3148, 154–154.
- Prendini, L. (2011f) Thelyphonida Latreille, 1804. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. Zootaxa*, 3148, 155–155.
- Prendini, L. (2011g) Schizomida Petrunkevitch, 1945. *In: Zhang, Z.-Q. (Ed.) Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. Zootaxa*, 3148, 156–156.
- Prokop, J. & Nel, A. (2007) An enigmatic Palaeozoic stem-group: Paoliida, designation of new taxa from the Upper Carboniferous of the Czech Republic (Insecta: Paoliidae, Katerinkidae fam. n.). *African Invertebrates*, 48 (1), 77–86
- Rasnitsyn, A.P. & Quicke D.L.J. (eds) (2002) *History of Insects*. Kluwer Academic Publishers, Dordrecht, Boston, London, 517 pp.
- Selden, P.A., Shear, W.A. & Sutton, M.D. (2008) Fossil evidence for the origin of spider spinnerets, and a proposed arachnid order. *Proceedings of the National Academy of Sciences of the USA*, 105, 20781–20785.
- Schatz, H., Behan-Pelletier, V.M., OConnor, B.M. & Norton, R.A. (2011) Suborder Oribatida van der Hammen, 1968. *In: Zhang, Z.-Q. (Ed.), Animal biodiversity: an outline of higher-level classification and survey of taxonomic richness. Zootaxa*, 3148, 141–148.
- Shcherbakov, D.E. (2011) The alleged Triassic palaeodictyopteran is a member of Titanoptera. *Zootaxa*, 3044, 65–68.
- Shear, W. (2011) Class Diplopoda de Blainville in Gervais, 1844. *In: Zhang, Z.-Q. (Ed.), Animal biodiversity: an outline of higher-level classification and survey of taxonomic richness. Zootaxa*, 3148, 159–164.
- Shultz, J.W. (2007) A phylogenetic analysis of the arachnid orders based on morphological characters. *Zoological Journal of the Linnean Society*, 150, 221–265.
- Slipinski, S.A., Leschen, R.A.B. & Lawrence, J.F. (2011) Order Coleoptera Linnaeus, 1758. *In: Zhang, Z.-Q. (Ed.), Animal biodiversity: an outline of higher-level classification and survey of taxonomic richness. Zootaxa*, 3148, 203–208.
- Szeptycki, A. (2007) Catalogue of the world Protura. Wydawnictwa Instytutu Systematyki i Ewolucji Zwierząt Polskiej Akademii Nauk, Kraków.
- Trautwein, M.D., Wiegmann, B.M., Beutel, R., Kjer, K.M. & Yeates, D.K. (2012) Advances in insect phylogeny at the dawn of the postgenomic era. *Annual Review of Entomology*, 57, 449–68.
- Trueman, J.W.H. (2007) A brief history of the classification and nomenclature of Odonata. *Zootaxa*, 1668, 381–394.
- Walter, D.E., Bolton, S., Uusitalo, M. & Zhang, Z.-Q. (2011) Suborder Endeostigmata Reuter, 1909. *In: Zhang, Z.-Q. (ed.) Animal biodiversity: an outline of higher-level classification and survey of taxonomic richness. Zootaxa*, 3148, 139–140.
- Zhang, Z.-Q., Fan, Q.-H., Pesic, V., Smit, H., Bochkov, A.V., Khaustov, A.A., Baker, A., Wohltmann, A., Wen, T.-H., Amrine, J.W., Beron, P., Lin, J.-Z., Gabrys, G. & Husband, R. (2011) Order Trombidiformes Reuter, 1909. *In: Zhang, Z.-Q. (ed.) Animal biodiversity: an outline of higher-level classification and survey of taxonomic richness. Zootaxa*, 3148, 129–138.

Author name and address

Zhi-Qiang Zhang, New Zealand Arthropod Collection, Landcare Research, Private Bag 92170, Auckland, New Zealand; zhangz@landcareresearch.co.nz

Order **Coleoptera** Linnaeus, 1758 (4 suborders)^{1,2}

†Suborder **Protocoleoptera** Tillyard, 1934 (3 superfamilies)

†Superfamily **Tshekardocoleoidea** Rohdendorf, 1944 (3 families)

†Family **Tshekardocoleidae** Rohdendorf, 1944 (13 genera, 18 species)

†Family **Labradorocoleidae** Ponomarenko, 1969 (1 genus, 1 species)

†Family **Oborocoleidae** Kukalová, 1969 (2 genera, 2 species)

†Superfamily **Permocupedoidea** Martynov, 1933 (2 families)

†Family **Permocupedidae** Martynov, 1933 (11 genera, 30 species)

†Family **Taldycupedidae** Rohdendorf, 1961 (6 genera, 18 species)

†Superfamily **Permosynoidea** Tillyard, 1924 (2 families)

†Family **Ademosynidae** Ponomarenko, 1968 (12 genera, 29 species)

†Family **Permosynidae** Tillyard, 1924 (3 genera, 14 species)

Suborder **Archostemata** Kolbe, 1908 (8 families)

Family **Crowsoniellidae** Iablokoff-Khnzorian, 1983 (1 genus, 1 species)

Family **Cupedidae** Laporte, 1836 (53(†44) genera, 120(†89) species)

Family **Micromalthidae** Barber, 1913 (2(†1) genera, 2(†1) species)

Family **Ommatidae** Sharp and Muir, 1912 (10(†8) genera, 120(†114) species)

Family **Jurodidae** Ponomarenko, 1985 (2(†1) genera, 3(†2) species)

†Family **Triadocupedidae** Ponomarenko, 1966 (8 genera, 16 species)

†Family **Magnocoleidae** Hong, 1998 (1 genus, 1 species)

†Family **Obrieniidae** Zherikhin and Gratshev, 1994 (5 genera, 7 species)

Suborder **Myxophaga** Crowson, 1955 (5 superfamilies)

†Superfamily **Asiocoloidea** Rohdendorf, 1961 (2 families)

†Family **Asiocolidae** Rohdendorf, 1961 (1 genus, 1 species)

†Family **Tricoleidae** Ponomarenko, 1969 (3 genera, 6 species)

†Superfamily **Rhombocoleoidea** Rohdendorf, 1961 (1 family)

†Family **Rhombocoleidae** Rohdendorf, 1961 (5 genera, 10 species)

Superfamily **Schizophoroidea** Ponomarenko, 1968 (3 families)

†Family **Schizophoridae** Ponomarenko, 1968 (20 genera, 36 species)

†Family **Catiniidae** Ponomarenko, 1968 (5 genera, 6 species)

†Family **Schizocoleidae** Rohdendorf, 1961 (4 genera, 30 species)

Superfamily **Lepiceroidea** Hinton, 1936 (1882) (2 families)

Family **Lepiceridae** Hinton, 1936 (1882) (1(†1) genus, 2(†1) species)³

Superfamily **Sphaeriuoidea** Erichson, 1845 (3 families)

Family **Torridincolidae** Steffan, 1964 (7 genera, 60 species)

Family **Hydroscaphidae** LeConte, 1874 (3 genera, 22 species)⁴

Family **Sphaeriusidae** Erichson, 1845 (2 (†1) genera, 20 (†1) species)

Suborder **Adephaga** Schellenberg, 1806 (17 families)

†Family **Tritarsusidae** Hong, 2002 (1 genus, 1 species)

Family **Gyrinidae** Latreille, 1810 (25 (†13) genera, 1000 (†118) species)

Family **Trachypachidae** C. G. Thomson, 1857 (22 (†21) genera, 57 (†51) species)

Family **Meruidae** Spangler and Steiner, 2005 (1 genus, 1 species)

Family **Rhysodidae** Laporte, 1840 (40 genera, 350 species)

Family **Carabidae** Latreille, 1802 (1500 genera, 40000 species)

1. **BY** S. A. Slipinski, R. A. B. Leschen, and J. F. Lawrence (for full contact details, see the list after **References**). The title of this contribution should be cited as "Order Coleoptera Linnaeus, 1758. In: Zhang, Z.-Q. (Ed.) *Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness*".
2. Recent Coleoptera currently includes 176 families, 29500 genera and 386500 species, fossil taxa includes 31 families, 230 genera and 600 species. The classification is based on Lawrence and Newton (1995) and Bouchard *et al.* (2011). Numbers for fossil taxa were derived from Kirejtshuk *et al.* (2010) and only numbers of fossils for extinct Protocoleoptera and Coleoptera families are provided, apart from extant primitive Polyphaga. Numbers for extant taxa were taken from the *Handbook of Zoology* chapters in Beutel & Leschen (2005) and Leschen *et al.* (2010) except where noted. Because up-to-date catalogues do not exist, numbers in the *Handbook* had included estimates and here we cross-referenced these estimates with recent catalogues (e.g., Löbl & Smetana 2007) and web-based information (e.g., Hallan 2010). Cut off date for the information contained here was 10 February 2011.
3. See Ge *et al.* (2010).
4. See Fikáček and Šípková (2009).

- Family **Haliplidae** Aubé, 1836 (5 (†1) genera, 220 (†2) species)
†Family **Triaplidae** Ponomarenko, 1977 (1 genus, 2 species)
†Family **Colymbotethidae** Ponomarenko, 1994 (2 genera, 2 species)
†Family **Parahygrobiidae** Ponomarenko, 1977 (1 genus, 1 species)
†Family **Coptoclavidae** Ponomarenko, 1961 (14 genera, 32 species)
†Family **Liadytidae** Ponomarenko, 1977 (2 genera, 6 species)
Family **Noteridae** C. G. Thomson, 1859 (14 genera, 250 species)
Family **Amphizoidae** LeConte, 1853 (1 genus, 5 species)
Family **Aspidytidae** Ribera et al., 2002 (1 genus, 2 species)
Family **Hygrobiidae** Régimbart, 1879 (1837) (1 genus, 5 species)
Family **Dytiscidae** Leach, 1815 (190 (†6) genera, 4100 (†85) species)
- Suborder **Polyphaga** Emery, 1886 (7 series)
- Series **Staphyliniformia** Latreille, 1802 (2 superfamilies)
- Superfamily **Hydrophiloidea** Latreille, 1802 (4 families)
- Family **Hydrophilidae** Latreille, 1802 (200 genera, 3400 species)
Family **Sphaeritidae** Shuckard, 1839 (1 genus, 5 species)
Family **Synteliidae** Lewis, 1882 (1 genus, 7 species)
Family **Histeridae** Gyllenhal, 1808 (350 genera, 4300 species)
- Superfamily **Staphylinoidea** Latreille, 1802 (6 families)
- Family **Hydraenidae** Mulsant, 1844 (42 genera, 1600 species)
Family **Ptiliidae** Erichson, 1845 (80 genera, 650 species)
Family **Agyrtidae** C. G. Thomson, 1859 (8 genera, 70 species)
Family **Leiodidae** Fleming, 1821 (342 genera, 3700 species)
Family **Silphidae** Latreille, 1807 (15 genera, 200 species)
Family **Staphylinidae** Latreille, 1802 (3500 genera, 56000 species)
- Series **Scarabaeiformia** Crowson 1960 (1 superfamily)
- Superfamily **Scarabaeoidea** Latreille, 1802 (14 families)
- Family **Pleocomidae** LeConte, 1861 (2 genera, 50 species)
Family **Geotrupidae** Latreille, 1802 (70 genera, 920 species)
Family **Belohinidae** Paulian, 1959 (1 genus, 1 species)
Family **Passalidae** Leach, 1815 (64 genera, 800 species)
Family **Trogidae** W. S. MacLeay, 1819 (5 genera, 300 species)
Family **Glaresidae** Kolbe, 1905 (3(†2) genera, 60 (†3) species)
Family **Diphyllostomatidae** Holloway, 1972 (1 genus, 3 species)
Family **Lucanidae** Latreille, 1804 (120(†6) genera, 1500 (†11) species)
Family **Ochodaecidae** Mulsant and Rey, 1871 (15 genera, 110 species, †2/3)
Family **Hybosoridae** Erichson, 1847 (77(†3) genera, 580 (†7) species)
Family **Glaphyridae** W. S. MacLeay, 1819 (11(†1) genera, 210(†6) species)
Family **Scarabaeidae** Latreille, 1802 (1900 genera, 27000 species)
†Family **Coprinisphaeridae** Genise, 2004 (4 genera, 9 species, ichnotaxon)
†Family **Pallichnidae** Genise, 2004 (2 genera, 3 species, ichnotaxon)
- Series **Scirtiformia** Fleming, 1821 (1 superfamily)
- Superfamily **Scirtoidea** Fleming, 1821 (6 families)
- Family **Scirtidae** Fleming, 1821 (35 genera, 800 species)
Family **Decliniidae** Nikitsky et al., 1994 (1 genus, 2 species)
Family **Eucinetidae** Lacordaire, 1857 (11 genera, 53 species)
Family **Clambidae** Fischer von Waldheim, 1821 (6 genera, 170 species)
†Family **Elodophthalmidae** Kirejtshuk and Azar, 2008 (1 genus, 2 species)
†Family **Mesocinetidae** Kirejtshuk and Ponomarenko, 2010 (4 genera, 7 species)
- Series **Elateriformia** Crowson 1960 (4 superfamilies)
- Superfamily **Dascilloidea** Guérin-Méneville, 1843 (1834) (2 families)
- Family **Dascillidae** Guérin-Méneville, 1843 (1834) (15 genera, 80 species)
Family **Rhipiceridae** Latreille, 1834 (7 genera, 70 species)
- Superfamily **Buprestoidea** Leach, 1815 (1 family)

- Family **Buprestidae** Leach, 1815 (470 genera, 14700 species)
- Superfamily **Byrrhoidea** Latreille, 1804 (12 families)
- Family **Byrrhidae** Latreille, 1804 (38 genera, 430 species)
- Family **Elmidae** Curtis, 1830 (149 genera, 1500 species)
- Family **Dryopidae** Billberg, 1820 (1817) (33 genera, 300 species)
- Family **Lutrochidae** Kasap and Crowson, 1975 (1 genus, 11 species)
- Family **Limnichidae** Erichson, 1846 (37 genera, 390 species)
- Family **Heteroceridae** W. S. MacLeay, 1825 (15 genera, 300 species)
- Family **Psephenidae** Lacordaire, 1854 (35 genera, 290 species)
- Family **Cneoglossidae** Champion, 1897 (1 genus, 10 species)
- Family **Ptilodactylidae** Laporte, 1836 (34 genera, 500 species)
- Family **Chelonariidae** Blanchard, 1845 (3 genera, 250 species)
- Family **Eulichadidae** Crowson, 1973 (2 genera, 30 species)
- Family **Callirhipidae** Emden, 1924 (9 genera, 150 species)
- Superfamily **Elateroidea** Leach, 1815 (20 families)
- Family **Rhinorhipidae** Lawrence, 1988 (1 genus, 1 species)
- Family **Artematopodidae** Lacordaire, 1857 (8 genera, 45 species)
- Family **Brachypsectridae** LeConte and G. H. Horn, 1883 (1 genus, 5 species)
- Family **Cerophytidae** Latreille, 1834 (3 genera, 21 species)
- Family **Eucnemidae** Eschscholtz, 1829 (200 genera, 1500 species)
- Family **Throscidae** Laporte, 1840 (5 genera, 150 species)
- Family **Elateridae** Leach, 1815 (400 genera, 10000 species)
- Family **Plastoceridae** Crowson, 1972 (1 genus, 2 species)
- Family **Drilidae** Blanchard, 1845 (6 genera, 120 species)
- Family **Omalisidae** Lacordaire, 1857 (3 genera, 8 species)
- Family **Lycidae** Laporte, 1836 (160 genera, 4600 species)
- Family **Telegeusidae** Leng, 1920 (3 genera, 10 species)
- Family **Phengodidae** LeConte, 1861 (31 genera, 250 species)
- Family **Rhagophthalmidae** E. Olivier, 1907 (6 genera, 30 species)⁵
- Family **Lampyridae** Rafinesque, 1815 (110 genera, 2200 species)
- Family **Omethidae** LeConte, 1861 (8 genera, 33 species)
- Family **Cantharidae** Imhoff, 1856 (1815) (160 genera, 5100 species)
- Family **Podabrocephalidae** Pic, 1930 (1 genus, 1 species)
- †Family **Berentimiridae** Winkler, 1987 (1 genus, 1 species)
- †Family **Lasiosynidae** Kirejtshuk, Chang, Ren & Kun, 2010 (4 genera, 7 species)
- Series **Derodontiformia** LeConte, 1861 (1 superfamily)
- Superfamily **Derontoidea** LeConte, 1861 (3 families)
- Family **Derodontidae** LeConte, 1861 (4 genera, 30 species)
- Family **Nosodendridae** Erichson, 1846 (1 genus, 50 species)
- Family **Jacobsoniidae** Heller, 1926 (3 genera, 20 species)
- Series **Bostrichiformia** Forbes, 1926 (1 superfamily)
- Superfamily **Bostrichoidea** Latreille, 1802 (4 families)
- Family **Dermestidae** Latreille, 1804 (50 genera, 1200 species)
- Family **Endecatomidae** LeConte, 1861 (1 genus, 4 species)
- Family **Bostrichidae** Latreille, 1802 (90 genera, 570 species)
- Family **Ptinidae** Latreille, 1802 (230 genera, 2200 species)
- Series **Cucujiformia** Lameere, 1938 (4 superfamilies)
- Superfamily **Lymexyloidea** Fleming, 1821 (1 family)
- Family **Lymexylidae** Fleming, 1821 (10 genera, 70 species)
- Superfamily **Cleroidea** Latreille, 1802 (12 families)
- Family **Phloiophilidae** Kiesenwetter, 1863 (1 genus, 1 species)
- Family **Trogossitidae** Latreille, 1802 (50 genera, 600 species)

5. See Löbl & Smetana (2007).

- Family **Chaetosomatidae** Crowson, 1952 (4 genera, 12 species)
 Family **Metaxinidae** Kolibáč, 2004 (1 genus, 1 species)
 Family **Thaneroceridae** Chapin, 1924 (7 genera, 30 species)
 Family **Cleridae** Latreille, 1802 (200 genera, 3400 species)
 Family **Acanthocnemidae** Crowson, 1964 (1 genus, 1 species)
 Family **Phycosecidae** Crowson, 1952 (1 genus, 4 species)
 Family **Prionoceridae** Lacordaire, 1857 (3 genera, 160 species)
 Family **Mauroniscidae** Majer, 1995 (5 genera, 26 species)
 Family **Melyridae** Leach, 1815 (300 genera, 6000 species)
 †Family **Parandrexidae** Kirejtshuk, 1994 (3 genera, 5 species)
- Superfamily **Cucujoidea** Latreille, 1802 (37 families)
- Family **Boganiidae** Sen Gupta and Crowson, 1966 (5 genera, 11 species)
 Family **Byturidae** Gistel, 1848 (7 genera, 24 species)
 Family **Helotidae** Chapuis, 1876 (5 genera, 107 species)
 Family **Protocucujidae** Crowson, 1954 (1 genus, 7 species)
 Family **Sphindidae** Jacquelin du Val, 1860 (9 genera, 59 species)
 Family **Biphyllidae** LeConte, 1861 (7 genera, 200 species)
 Family **Erotylidae** Latreille, 1802 (260 genera, 3500 species)
 Family **Monotomidae** Laporte, 1840 (33 genera, 250 species)
 Family **Hobartiidae** Sen Gupta and Crowson, 1966 (2 genera, 6 species)
 Family **Cryptophagidae** Kirby, 1826 (60 genera, 600 species)
 Family **Agapythidae** Sen Gupta and Crowson, 1969 (1 genus, 1 species)
 Family **Priasilphidae** Crowson, 1973 (3 genera, 11 species)
 Family **Phloeostichidae** Reitter, 1911 (5 genera, 14 species)
 Family **Silvanidae** Kirby, 1837 (58 genera, 500 species)
 Family **Cucujidae** Latreille, 1802 (4 genera, 44 species)
 Family **Myraboliidae** Lawrence and Britton, 1991 (1 genus, 13 species)
 Family **Cavognathidae** Sen Gupta and Crowson, 1966 (1 genus, 9 species)
 Family **Lamingtoniidae** Sen Gupta and Crowson, 1969 (1 genus, 3 species)
 Family **Passandridae** Blanchard, 1845 (9 genera, 109 species)
 Family **Phalacridae** Leach, 1815 (51 genera, 640 species)
 Family **Propalticidae** Crowson, 1952 (2 genera, 30 species)
 Family **Laemophloeidae** Ganglbauer, 1899 (37 genera, 430 species)
 Family **Tasmosalpingidae** Lawrence and Britton, 1991 (1 genus, 2 species)
 Family **Cyclaxyridae** Gimmel et al., 2009 (1 genus, 2 species)
 Family **Kateretidae** Kirby, 1837 (14 genera, 95 species)
 Family **Nitidulidae** Latreille, 1802 (350 genera, 4500 species)
 Family **Smicripidae** G. H. Horn, 1880 (1 genus, 6 species)
 Family **Bothrideridae** Erichson, 1845 (38 genera, 400 species)
 Family **Cerylonidae** Billberg, 1820 (52 genera, 450 species)
 Family **Alexiidae** Imhoff, 1856 (1 genus, 50 species)
 Family **Discolomatidae** G. H. Horn, 1878 (16 genera, 400 species)
 Family **Endomychidae** Leach, 1815 (130 genera, 1800 species)
 Family **Coccinellidae** Latreille, 1807 (360 genera, 6000 species)
 Family **Corylophidae** LeConte, 1852 (30 genera, 200 species)
 Family **Latridiidae** Erichson, 1842 (28 genera, 1000 species)
 Family **Akalypsoischiidae** Lord et. al, 2010 (1 genus, 24 species)⁶
 †Family **Sinisilvanidae** Hong, 2002 (1 genus, 1 species)
- Superfamily **Tenebrionoidea** Latreille, 1802 (28 families)
- Family **Mycetophagidae** Leach, 1815 (18 genera, 130 species)
 Family **Archeocrypticidae** Kaszab, 1964 (10 genera, 60 species)
 Family **Pterogeniidae** Crowson, 1953 (7 genera, 26 species)

6. See Lord et al. (2010).

- Family **Ciidae** Leach, 1819 (42 genera, 650 species)
 Family **Tetratomidae** Billberg, 1820 (13 genera, 150 species)
 Family **Melandryidae** Leach, 1815 (60 genera, 420 species)
 Family **Mordellidae** Latreille, 1802 (100 genera, 1500 species)
 Family **Rhiphoridae** Gemminger, 1870 (1853) (38 genera, 400 species)
 Family **Zopheridae** Solier, 1834 (190 genera, 1700 species)
 Family **Ulodidae** Pascoe, 1869 (14 genera, 30 species)
 Family **Promecheilidae** Lacordaire, 1859 (7 genera, 20 species)
 Family **Chalcodryidae** Watt, 1974 (3 genera, 15 species)
 Family **Trachelostenidae** Lacordaire, 1859 (1 genus, 2 species)
 Family **Tenebrionidae** Latreille, 1802 (2300 genera, 20000 species)
 Family **Prostomidae** C. G. Thomson, 1859 (2 genera, 30 species)
 Family **Synchroidae** Lacordaire, 1859 (3 genera, 8 species)
 Family **Stenotrachelidae** C. G. Thomson, 1859 (7 genera, 19 species)
 Family **Oedemeridae** Latreille, 1810 (100 genera, 500 species)
 Family **Meloidae** Gyllenhal, 1810 (120 genera, 3000 species)
 Family **Mycteridae** Oken, 1843 (29 genera, 160 species)
 Family **Boridae** C. G. Thomson, 1859 (3 genera, 4 species)
 Family **Trictenotomidae** Blanchard, 1845 (2 genera, 13 species)
 Family **Pythidae** Solier, 1834 (7 genera, 23 species)
 Family **Pyrochroidae** Latreille, 1807 (30 genera, 167 species)
 Family **Salpingidae** Leach, 1815 (45 genera, 300 species)
 Family **Anthricidae** Latreille, 1819 (100 genera, 3000 species)
 Family **Aderidae** Winkler, 1927 (50 genera, 900 species)
 Family **Scraptiidae** Gistel, 1848 (35 genera, 500 species)
 Superfamily **Chrysomeloidea** Latreille, 1802 (7 families)
 Family **Vesperidae** Mulsant, 1839 (17 genera, 75 species)
 Family **Oxypeltidae** Lacordaire, 1868 (2 genera, 3 species)⁷
 Family **Disteniidae** J. Thomson, 1861 (32 genera, 336 species)
 Family **Cerambycidae** Latreille, 1802 (5232 genera, 30079 species)⁸
 Family **Megalopodidae** Latreille, 1802 (30 genera, 350 species)⁹
 Family **Orsodacnidae** C. G. Thomson, 1859 (3 genera, 40 species)⁹
 Family **Chrysomelidae** Latreille, 1802 (2114 genera, 32500 species)
 Superfamily **Curculionoidea** Latreille, 1802 (8 families)¹⁰
 Family **Nemonychidae** Bedel, 1882 (20 genera, 70 species)
 Family **Anthribidae** Billberg, 1820 (372 genera, 3900 species)
 Family **Belidae** Schoenherr, 1826 (38 genera, 375 species)
 Family **Attelabidae** (150 genera, 2500 species)
 Family **Caridae** Thompson, 1992 (4 genera, 6 species)
 Family **Brentidae** Billberg, 1820 (400 genera, 4000 species)
 Family **Curculionidae** Latreille, 1802 (4600 genera, 51000 species)
 †Family **Ulyanidae** Zherikhin, 1993 (2 genera, 3 species)

Acknowledgements

We thank Petr Svacha (Vesperidae), Miguel Monne (Cerambycidae), Den Heffern (Cerambycidae), Antonio Santos-Silva (Disteniidae), and Chris Reid (Chrysomelidae) for assistance with the numbers of chrysomeloid families. RABL

7. See Kuschel (1955).

8. Number of genera was obtained by tripling the number of neotropical genera (1744) supplied by M. Monne (pers. com). Species number was obtained from Anonymous (2010).

9. See Hallan (2010).

10. See Oberprieler et al. (2007).

was supported in part by Foundation for Research, Science and Technology through the Defining New Zealand's Land Biota OBI.

Literature Cited

- Anonymous. (2010) Biolib.cz. <http://www.biolib.cz/en/taxon/id10991/> [accessed 6 January 2011].
- Beutel, R. G., & Leschen, R.A.B. (2005) *Handbook of Zoology, Volume IV Arthropoda: Insecta, Part 38, Evolution and Systematics Coleoptera (Archostemata, Adephaga, Myxophaga, Polyphaga part.)*. Walter de Gruyter, Berlin. 567 p.
- Bouchard, B., Bousquet, Y., Davies, A.E., Alonso-Zarazaga, M.A., Lawrence, J.F., Lyal, C.H.C., Newton, A.F., Reid, C.A.M., Schmitt, M., Ślipiński, S.A., Smith, A.B.T. (2011) Family-group names in Coleoptera. *Zookeys*, 88, 1–972.
- Fikáček, M., Šípková H. (2009) New Asian *Hydroscapha*, with comments on male-female association of co-occurring species (Coleoptera, Myxophaga, Hydroscaphidae) *Zootaxa*, 2286, 31–48.
- Ge, S.-Q., Friedrich, F. & Beutel, R. G. (2010) On the systematic position and taxonomic rank of the extinct myxophagan †*Haplocheilus* (Coleoptera). *Insect Systematics & Evolution*, 41, 329–338.
- Hallan, J. (2010) *Synopsis of the described Coleoptera of the world*. <http://insects.tamu.edu/research/collection/hallan/test/Arthropoda/Insects/Coleoptera/Family/Coleoptera1.htm> [accessed 6 January 2011].
- Kirejtshuk, A.G., Ponomarenko, F.G. & Zherikhin, V.V. (2010) *Catalogue of fossil Coleoptera*. <http://www.zin.ru/animalia/coleoptera/eng/paleosys.htm>. [Accessed November 2010].
- Kuschel, G. (1955) Una nueva especie de *Cheloderus* Castelnau. *Revista Chilena de Entomología*, 4, 251–254.
- Lawrence, J. F. & Newton, A. F. (1995) Families and subfamilies of Coleoptera (with selected genera, notes, references and data on family-group names). In: Pakaluk, J. & Ślipiński, S. A. (Eds), *Biology, Phylogeny, and Classification of Coleoptera: Papers Celebrating the 80th Birthday of Roy A. Crowson*. Muzeum Instytut Zoologii PAN, Warszawa, pp. 779–1006.
- Leschen, R.A.B., Beutel, R.G., & Lawrence, J.F. (2010) *Handbook of Zoology, Coleoptera Volume 2: Morphology and Systematics (Elateroidea, Bostrichiformia, Cucujiformia partim)*. Walter de Gruyter, Berlin. 786 p.
- Löbl, I. & Smetana A. (2007) *Catalogue of Palaearctic Coleoptera. Volume 4. Elateroidea, Derodontoidea, Bostrichoidea, Lymexyloidea, Cleroidea and Cucujoidea*. Apollo Books, Stenstrup. 935 p.
- Lord, N.P., Hartley, C.S., Lawrence, J.F., McHugh, J.V., Whiting, M.F., & Miller, K.B. (2010) Phylogenetic analysis of the Minute Brown Scavenger Beetles (Coleoptera: Latridiidae), and recognition of a new beetle family, Akalyptoischionidae, fam. n. (Coleoptera: Cucujoidea). *Systematic Entomology*, 35, 753–763.
- Oberprieler, R.G., Marvaldi, A.E. & Anderson, R.S. (2007) Weevils, weevils, weevils everywhere. In: Zhang, Z.-Q. & Shear, W.A. (Eds.) *Linnaeus Tercentenary: Progress in Invertebrate Taxonomy*. *Zootaxa*, 1668, 491–520.

Author names and contact addresses

S.A. Slipinski, CSIRO Ecosystem Sciences, GPO Box 1700, Canberra, ACT, 2601, Australia

R.A.B. Leschen, New Zealand Arthropod Collection, Private Bag 92170, Auckland, New Zealand (corresponding author; leschenr@landcareresearch.co.nz)

J.F. Lawrence, CSIRO Ecosystem Sciences, GPO Box 1700, Canberra, ACT, 2601, Australia