

Type specimens of some Palaearctic Miridae in the Zoological Museum, Helsinki (Heteroptera)

I.M. Kerzhner

Kerzhner, I.M. 1997. Type specimens of some Palaearctic Miridae in the Zoological Museum, Helsinki (Heteroptera). *Zoosystematica Rossica*, 6(1/2): 115-121.

Notes on some type specimens of Miridae in the Zoological Museum, Helsinki are given. Lectotypes of 140 species-group taxa are designated. Type localities are indicated for some holotypes and lectotypes designated by H. Lindberg and E. Wagner. *Dichrooscytus josifovi* sp. n. is established for *D. albidovirens* sensu Josifov, 1974 (non Reuter, 1896). New synonymy: *Deracocoris ventralis ventralis* Reuter, 1904 = *D. scutellaris* var. *alboscutellatus* Reuter, 1891 (nomen oblitum).

I.M. Kerzhner, Zoological Institute, Russian Academy of Sciences, Universitetskaya nab. 1, St.Petersburg 199034, Russia.

Introduction

In May 1997 the author had an opportunity to work 3 days in the collection of the Zoological Museum, Helsinki. The collection includes type specimens of numerous species and varieties described by famous Finnish hemipterists O.M. Reuter, B. Poppius, C.R., R.F. and J. Sahlberg, H. Lindberg and by some of their contemporaries. Lindberg (1951) published a list of type specimens of Palaearctic species in the collection of O.M. Reuter forming part of the collection of the Museum, but his list contains many errors and omissions.

This paper includes species both described from the Palaearctic Region and those described from other regions but later found in the Palaearctic or in adjacent territories. It is not a complete list of types preserved in the Museum. Lectotype designations are made selectively, mostly for taxa which were apparently described from one specimen, or for which syntypes in other museums are badly damaged or lost, but some lectotype designations are made for other reasons. For some holotypes and lectotypes designated by E. Wagner and H. Lindberg the missing information on type locality is published here. Special attention is paid to types of Kirkaldy, Signoret and some other authors for which the current deposition in the Helsinki Museum is more or less unexpected. Some mistakes in Lindberg (1951) are corrected.

Two peculiarities of labelling in Reuter's collection should be taken into consideration: (1) he often replaced exact original labels with his own labels indicating a wide area (e.g. "Ross. mer." instead Sarepta or "Sibiria" instead Irkutsk); (2) he often did not label varieties, especially those distinguished by Greek letters only.

The names of taxa for which lectotypes are designated in this paper are asterisked. The following ab-

breviations are used: gc – glued on card, gt – glued on triangle, pn – pinned, mpn – micropinned, genit. diss. – genitalia dissected, HT – holotype, LT – lectotype, PLT – paralectotype(s), ST – syntype, STS – syntypes. Labels are separated from each other by oblique stroke. Handwritten labels are marked [h], with addition of the following abbreviations when the handwriting is known: H – Horváth, Hn – Handlirsch, J – Jakovlev, K – Kirkaldy, Kir – Kiritschenko, L – Lindberg, P – Poppius, R – Reuter, S – J. Sahlberg, Seid – Seidenstücker, Sg – Signoret. Text in Cyrillic characters is noted [Cc], printed text before and within handwritten text is marked [p]. Printed label "Coll. O.M. Reuter" is cited as "cR". Of the blue type labels of the museum "Mus. Zool. H:fors Spec. typ. No. (number) (species name)" only the number of the specimen is cited.

List of species

**acanthoides* J. Sahlberg, 1875 (*Platypsallus*). LT: ♂, gc, Ponoi/ J. Sahlbg./ J. Sahlb. ind. typ./ No. 3087. PLT: 1 ♀, as LT but No. 3086; 1 ♀, gc, Lapp. ross. [hR]/ Reuter/ cR/ Spec. typ. J. Sahlberg/ No. 3088. Currently *Chlamydatus* id.

**aenescens* Reuter, 1899 (*Pachytomella*). LT: ♂ macropt., gc, Dr Schmiedeknecht, Oran, '95, Djebel Tessalah/ Spec. typ. Reuter/ cR/ No 3490. PLT: 1 ♀ macropt., 1 ♀ brachypt., as LT, but No. 3491 and 3492. Synonym of *Dasysscytus sordidus* Fieber, 1864.

**albicans* Reuter, 1901 ("*Psallus*?"). LT: ♀, gc, Bekljär-bek/ J. Sahlb./ 215 [h]/ *Psallus albicans* Reut. Typ. [hR]/ No. 11256. PLT: 1 ♀, gc, Bekljärbek/ J. Sahlb./ No. 11251; 1 ♀, Bekljär-bek/ J. Sahlb./ Spec. typ. Reuter/ cR/ No. 3294. Synonym of *Plagiognathus hipunctatus* Reuter, 1883.

**albicans* Stichel, 1956 (*Amblytylus tarsalis* f.). LT: ♀, Madrid [hR]/ Spec. typ. Reuter/ No. 3321. – Note.

The name was given to var. β of Reuter (1894: 142) based on female(s) from Madrid.

**albidovirens* Reuter, 1896 (*Dichrooscytus pseudosabinae* var.). LT: ♀ (abdomen broken, glued on card), pn, Turkestan [hR]/ Spec. typ. Reuter/ cR/ No. 3654. – Note. The specimen was identified as belonging to var. *albidovirens* by comparison with the original description. It is a pale specimen of *D. pseudosabinae*, and it does not belong to the species for which the name *Dichrooscytus albidovirens* Reuter, 1896 was used by Josifov (1974: 155). Accordingly, the name *Dichrooscytus josifovi* sp. n. is given to the species described by Josifov. Holotype: ♂. Sary-Chelek Lake, Chatkal Range, 22.VI.1966 (Loganova), Zoological Institute, St. Petersburg. Paratypes: specimens listed by Josifov (1974) as material examined of *D. albidovirens*.

**albidus* Reuter, 1903 (*Megalocoleus*). LT: ♂ (genit. diss.), gc, Aschabad/ Ahnger/ Spec. typ. Reuter/ cR/ No. 3335. PLT: 1 ♂, 1 ♀, as LT, but No. 3333 and 3334, and ♂ with "560"; 1 ♂, gc, Aschabad/ Ahnger/ 552/ *Megalocoleus albidus* Reut. Typ. [hR]. Currently *Lopidodenus* id.

**albifrons* Reuter, 1895 (*Fulvius*). LT: ♀, gc, Aden, Simon [h]/ *Rhinofulvius albifrons* Reut. [h]/ No. 12347/ Holotypus [added by Schmitz]. Currently *Rhinofulvius* id.

alboscutellatus Reuter, 1891 (*Deraeocoris scutellaris* var.). HT: ♀, gc, Minusinsk, K. Ehnberg/ 38/ Spec. typ. Reuter/ cR/ No. 3572. – Note. Examination of the HT allows me to establish definitely the following synonymy earlier already suspected by Josifov & Kerzhner (1972: 155): *Deraeocoris ventralis ventralis* Reuter, 1904 = *D. scutellaris* var. *alboscutellatus* Reuter, 1891 (nomen oblitum), syn. n.

**algericus* Reuter, 1896 (*Phytocoris*). LT: ♂ (genit. diss.), gc, Edough [h]/ Spec. typ. Reuter/ No. 3713/ cR.

alluaudi Poppius, 1912 (*Deraeocoris*). HT in Helsinki Museum.

**amurensis* Reuter, 1883 (*Plagiognathus*). LT: ♀, gc, Amur [hR]/ Spec. typ. Reuter/ No. 3258.

**annulicornis* R.F. Sahlberg, 1848 (*Capsus*). LT: ♂, pn, Yläne/ Sahlbg/ Spec. typ./ No. 3055. PLT: 1 ♂, as LT but No. 3056. Synonym of *Adelphocoris quadripunctatus* (Fabricius, 1794).

**annulipes* Reuter, 1908 (*Mermittelocerus*). LT: ♂, 3.68/ N 2 [h]/ *Mermittelocerus annulipes* Reut. n. sp. [hR] O.M. Reuter det./ No. 12278.

**antennalis* Reuter, 1891 (*Pachytomella phoenicea* var.). LT: ♀ brachypt., gc/ Jerusalem, Reitter/ Spec. typ. Reuter/ cR/ No. 3490/ var. *antennalis* Reut. [h].

**anthocoroides* Reuter, 1875 (*Teratodella*). LT: ♂ (genit. diss.), gc, Rouen [hR]/ *Fulvius brevicornis* Reut. [hR]/ No. 9989/ Paratype [added by G. Schmitz]/ genital prep. 633-71 [added by Schmitz]. Currently *Fulvius* id.

**anticus* J. Sahlberg, 1920 (*Phytocoris dimidiatus* var.). LT: ♂, gc, Jyväskylä/ J. Sahlbg./ in Alno glutinoso 8 Juli 1914 [hS]/ var. *anticus* n. var. [hS].

**apicale* Reuter, 1882 (*Megacoelum*). LT: ♂, gc, Addah [h]/ Reuter/ Spec. typ. PLT: 1 ♀, as LT but with additional label "*Megacoelum apicale* Reut." [hR].

**artemisiae* J. Sahlberg, 1878 (*Orthotylus*). LT: ♂ (genit. diss.), gc, Kantaika/ Spec. typ. J. Sahlberg/

cR/ No. 3428. PLT: 1 ♂, 2 ♀, as LT but 3426, 3427, 3429; 1 ♂, 1 ♀, gc, Kantaika [hS]/ J. Sahlbg/ J. Sahlb. ind. typ.

**atlanticus* Wagner, 1951 (*Dicyphus boliviari* ssp.). HT collected at Santa Cruz, Tenerife.

**bicolor* Reuter, 1878 (*Maurodactylus*). LT: ♂, gt, Astrachan [hR]/ *bicolor* Reut. (♂)/ Spec. typ. Reuter/ cR/ No. 3176. Synonym of *M. albidus* (Kolenati, 1845).

**bimaculicollis* Lindberg, 1948 (*Adelphocoris*). HT collected at Galata.

**bimaculatus* Reuter, 1894 (*Calocoris hispanicus* var.). LT: ♀, pn, Sevilla [h]/ Spec. typ. Reuter/ cR/ No. 3683/ v. *bimaculatus* Reut. [hR]. Synonym of *Calocoris nemoralis* (Fabricius, 1787).

**bispinulatus* Reuter, 1891 (*Calocoris lineolatus* var.). LT: ♂, gc, Blagov. [hR]/ Spec. typ. Reuter/ cR/ No. 3699/ v. *bispinulatus* Reut. [hR]. PLT: 1 specimen without abdomen, pn, Amur [hR]/ Spec. typ. Reuter/ cR/ No. 3700; 1 ♂, pn, Mongolia/ Fl. Jenisej/ K. Ehnberg/ cR. Currently *Adelphocoris* id.

**boliviari* Reuter, 1880 (*Orthocephalus*). LT: ♂, gc, Hispania [hR]/ Spec. typ. Reuter/ cR/ No. 3533. Currently *Plagiophylax* id.

**brachymerus* Reuter, 1877 (*Phytocoris*). The species was described from Philippeville. Two specimens in the Helsinki Museum, from L'Edough and Palesko, were erroneously listed as types by Lindberg (1951) and the first of them invalidly designated as LT by Wagner (1967: 278), though the latter mentioned a true syntype kept in the Noualhier's collection (Paris Museum).

**caucasicus* Horváth, 1927 (*Myrmecophyes*). LT: ♀, gc, Kaukasus, A. Kiritshenko/ *Myrmecophyes* sp.? [hKir] A. Kiritshenko det./ Typus [p] *Myrmecophyes caucasicus* Horv. [hH]/ cR/ No. 11321. Junior synonym of *M. alacer* Horváth, 1927.

**caucasicus* Poppius, 1912 (*Calocoris*). LT: ♂ (damaged), gc, Dzhul'fa [Cc], 12.V.14 [hKir]/ cR/ No. 9551/ *Calocoris caucasicus* n. sp. [hP] B. Poppius det. Of the second syntype (♀?) only middle legs and a hind femur remained.

**caucasicus* Poppius, 1912 (*Globiceps*). LT: ♂ (genit. diss.), pn, Caucasus occ., Krasnaja Poljana, 07.V/ No. 9541. PLT: 1 ♀, gc, Caucasus occ., Krasnaja Poljana, 07.15.VI, Kiritshenko/ 24 [hKir]/ *Globiceps caucasicus* n. sp. [hP] B. Poppius det./ No. 9542.

**chinensis* Signoret, 1858 (*Pachypeltis*). LT: ♀, pn, China [hHn] coll. Signoret/ *chinensis* [hHn] det. Signoret/ *Pachypeltis chinensis* mihi, China [hSg].

**coelestialium* Kirkaldy, 1902 (*Olympiocapsus*). LT: ♂ (genit. diss.), gc, Ngan-Hoei, Ho-chan, Joannis [h]/ *Olympiocapsus caelestialium* [sic] Kirk. type [hK]/ *Eurystylus caelestialium* (Kirk.) [hP] B. Poppius det. PLT: 1 ♀, on the same card. Currently *Eurystylus* id.

**concolor* Reuter, 1890 (*Plagiorrhama*). HT: ♀, gc, Petr. [h, Cc]/ Spec. typ. Reuter/ cR/ No. 3565. Currently *Hallopodus* id. – Note. Lindberg (1951) incorrectly indicated as syntype a specimen from Bekljarn-Bek, and Wagner (1970: 71) invalidly designated it as LT.

**confluens* Reuter, 1896 (*Calocoris hispanicus* var.). LT: ♀, pn, Attica/ Spec. typ. Reuter/ cR/ No.

3682/ v. *confluens* Reut. [hR]. Synonym of *Calocoris nemoralis* (Fabricius, 1787).

convexus Reuter, 1875 (*Psallus*). ST?: ♀, gc, Corse/ cR. Synonym of *P. mollis* (Mulsant & Rey, 1852).

**croceus* J. Sahlberg, 1888 (*Orthotylus*). LT: ♀, gc, Dv. [h]/ Dvoretz/ Günther/ *croceus* J. Sahlb. [hS]. Synonym of *O. ericetorum* (Fallén, 1807).

eruciatus R.F. Sahlberg, 1848 (*Lopus*). HT: ♂, pn, ♂ [h]/ Yläne/ F. Sahlb./ Spec. typ./ No. 3092. Currently *Macrotylus* id.

**cunealis* Reuter, 1901 (*Psallus*). LT: ♂ (destroyed, only some legs remained; data on sex taken from original description), gc, Dschilarik/ J. Sahlb./ 400 [h]/ *Psallus cunealis* Reut. Typ. [hR]/ cR/ No. 11249. Synonym of *Sacculifer rufinervis* (Jakovlev, 1880).

debilicornis Reuter, 1875 (*Atractotomus*). HT: ♀, Yläne/ J. Sahlb./ *Ps. debilicornis* Reut. sp. typ. [hS]/ 187/ No. 3100. Synonym of *A. kolenatti* (Flor, 1860).

**discolor* J. Sahlberg, 1878 (*Orthotylus*). LT: ♂ (genit. diss.), gc, Kantaika [hS]/ Spec. typ. J. Sahlberg/ cR/ No. 3433. PLT: 2 ♀, as LT but from Dudinka and No. 3431, 3432; 1 ♂, gc, Kantaika [hS]/ J. Sahlbg./ J. Sahlb. ind. typ./ *Orthotylus discolor* J. Sahlb. [h, bottom label]; 1 ♀, gc, Tolstoinos [hS]/ J. Sahlbg./ J. Sahlb. ind. typ.

**dispneta* Reuter, 1903 (*Dicyphus geniculatus* var.). LT: ♂, Bejar, Spain, G.C.C./ Spec. typ. Reuter/ cR/ No. 3467. – Note. A female from the same sample is also labelled as type, but the variety was described by Reuter from males.

**dissimilis* Reuter, 1896 (*Poecilocyptus*). LT: ♀, pn, Turkestan [hR]/ Spec. typ. Reuter/ cR/ No. 3590. Currently *Polymerus* id.

**distinguendus* Reuter, 1875 (*Lygus*). LT: ♀, pn, Siber. or. [hR]/ Spec. typ. Reuter/ cR/ No. 3598. Currently *Salignus* id. – Note. The type locality was stated as Irkutsk.

doriae Reuter in Ferrari, 1884 (*Orthocephalus*). Data on a type in the Helsinki Museum (Lindberg, 1951) are erroneous and refer to *O. nitens*.

**dubia* Rey, 1890 (*Camptobrochis lutescens* var.). LT: ♂, gc, Provence [hR]/ v. *dubius* Rey [hR]/ Spec. typ. C. Rey/ cR. PLT: 1 ♂, as LT, but "Gallia m." instead "Provence". – Note. These specimens (No. 3611 and 3613) were listed by Lindberg (1951) in error as types of *Phytocoris lutescens* Schilling, 1837.

**dubius* Reuter, 1901 (*Globiceps*). LT: ♂ (genit. diss.), gc, Issyk-kul/ J. Sahlb./ 487 [h]/ *Globiceps dubius* Reut. Typ. [hR]. Synonym of *G. fulvicollis* Jakovlev, 1877. – Note. In the original description as type locality was indicated Dzhilaryk.

**elegans* Jakovlev, 1867 (*Psallus*). LT: ♂ (apex of abdomen missing), gc, Astrachan [hR]/ Spec. typ. Jakovleff/ No. 3149. PLT: 1 ♀, as LT but No. 3150. Currently *Tuponia* id.

**elegans* Reuter, 1877 (*Calocoris*). LT: ♂, pn, Syria [hR]/ *Calocoris elegans* Reut. [hR]/ Spec. typ. Reuter/ cR/ No. 3666. Synonym of *Grypocoris amoenus* (Douglas & Scott, 1868).

**elegantulum* Reuter, 1899 (*Compsidolon*). LT: ♀, gc, Jerico/ J. Sahlb./ 573 [h]/ *Compsidolon elegantulum* Reut. Typ. [hR]/ No. ... [not noted].

**ephedrae* Reuter, 1902 (*Nasocoris*). LT: ♀ (at the left side of the plate), gc, Albarracin, Spain, G.C.C./ Spec. typ. Reuter/ cR/ No. 3398. PLT: 1 ♀, at the

right from the LT; 1 ♂ (without abdomen), 1 ♀, gc (at one plate), as LT but No. 3397.

**epilobii* Reuter, 1883 (*Dicyphus*). LT: ♂, gc, Hast[ings]/ Spec. typ. Reuter/ cR. PLT: 1 ♂, as LT.

**femoralis* Reuter, 1901 (*Psallopis*). LT: ♀, gc, Biskra/ J. Sahlb./ 1939 [h]/ cR/ *Psallopis femoralis* Reut. n. g. et sp. [hR].

**flaveolus* Reuter, 1870 (*Tytthus*). LT: ♂, gc, Par-gas/ Reuter/ Spec. typ. Reuter/ No. 3076. PLT: 1 ♂, 2 ♀, gc, Pargas/ Reuter/ No. 3077, 3078, and 3079. Currently *Fieberocapsus* id. – Note. Described from several localities in the Pargas parish.

**flavosparsus* C.R. Sahlberg, 1841 (*Phytocoris*). LT: ♂, pn, ♂ [h]/ Yläne/ C. Sahlberg/ Spec. typ./ J. Sahlb./ No. 3094. PLT: 1 ♀, as LT, but ♀ [h] and No. 3093. Currently *Orthotylus* id.

**flori* J. Sahlberg, 1870 (*Teratocoris*). LT: ♂, gc, golden square/ Tschuja/ Car. ross./ J. Sahlbg./ 404 [h]/ *Teratocoris Flori* J. Sahlb. [h]. Synonym of *T. saundersi* Douglas & Scott, 1869.

fraudatrix Reuter, 1904 (*Byrsoptera rossica* var.). The variety (now considered a separate species) was described from female(s). Three topotypical males, erroneously listed by Lindberg (1951) as types of *B. rossica* Reuter, are in the collection, but no females are found.

freyi Wagner, 1954 (*Psallus (Coniortodes)*). HT from Pico de Teyde. Currently *Compsidolon* id.

**fulvipes* Reuter, 1904 (*Orthocephalus tenuicornis* var.). LT: ♀, gc, Tarsus/ J. Sahlb./ 652/ cR/ *Orthocephalus tenuicornis* M. R. v. *fulvipes* Reut. [hR] O.M. Reuter det.

**fusca* Stichel, 1930 (*Alloeotomus gothicus* f.). LT: ♂, pn, Bohemia [hR]/ cR. PLT: 1 ♂, 1 ♀, as LT. – Note. The name was given by Stichel to var. β of Reuter (1896). Reuter described this variety from Bohemia (Chodau, now Chodov, leg. v. Stein). The specimens belong to *A. gothicus*, not to the later described *A. germanicus* Wagn.

fuscovenosus Fieber, 1864 (*Psallus?*). Contrary to Lindberg (1951), specimens in Helsinki Museum designated as types (from Romania and from Sarepta, the latter received from Jakovlev) do not belong to the type series. Currently *Solenoxyphus* id.

**geocoriceps* Reuter, 1904 (*Eurycranella*). LT: ♀, gc, Egypt. sup./ Deschena/ U. Sahlb./ 104/ Spec. typ./ *Eurycranella geocoriceps* n. gen. et sp. [hR] O.M. Reuter det./ No. 9524.

**gracilis* F.R. Sahlberg, 1848 (*Globiceps*). LT: ♂ macr., pn, Karlä/ Uleåborg/ coll. Nyldr/ Spec. typ./ 12.VIII.47 [h]/ No. 3052. Currently *Myrmecoris* id.

**griseoscelis* J. Sahlberg, 1920 (*Phytocoris longipennis* var.). LT: ♀, gc, Jyväskylä/ J. Sahlb./ var. *griseoscelis* n. var. [hS].

**henschii* Reuter, 1888 (*Halticus*). LT: ♂, mpn, Görz [hR]/ Spec. typ. Reuter/ cR/ No. 3481. PLT: 1 ♂, 2 ♀, as LT, but No. 3482, 3483, 3484.

**horvathi* Reuter, 1876 (*Amblytylus*). LT: ♀, gc, Buda [hR]/ Spec. typ. Reuter/ cR/ No. 3312. Currently *Macrotylus* id.

**hyperboreus* J. Sahlberg, 1868 (*Teratocoris*). LT: ♂, gc, golden square/ Enontekis/ J. Sahlbg./ ♂/ Spec. typ./ No. 3066. PLT: 3 ♂, 4 ♀, all from Enontekis, except for 1 ♂ labelled "Lapponia/ Palmén" and 1 ♀ from Muonioniska. Synonym of *T. viridis* Douglas &

Scott, 1867. – Note. The species was described from several localities, all of them but Muonioniska [= Muonio] in the province Enontekiö Lappi (the original description uses Torneå Lappmark, the earlier Swedish name).

**inconspicua* Reuter, 1876 (*Ethelastia*). LT: ♂, mpn, Ross. mer. [hR]/ Spec. typ. Reuter/ cR/ No. 3304. Synonym of *E. liturata* (Fieber, 1858). – Note. The species was described from Sarepta.

**innotata* Reuter, 1913 (*Campylobrochis lutescens* var.). LT: ♀, mpn, Museum Paris, France, Gif (Seine-&-Oise), J. Martin, 1897.

innotata Reuter, 1870 (*Lygus*). HT: ♀, gc, Pargas/ J. Sahlbg./ Spec. typ./ No. 3057. Synonym of *Lygocoris contaminatus* (Fallén, 1807).

**insularis* Reuter, 1896 (*Calocoris sexguttatus* var.). LT: ♂, gc, Scotia/ Spec. typ. Reuter/ No. 3669/ v. *insularis* Reut. [hR]. PLT: 1 ♀, gc, Scotia/ Spec. typ. Reuter/ No. 3670/ *Calocoris (Clusterotomus) insularis* (Reut. 1896) Stys 1956 ♀. det. Stys 1956. Synonym of *Calocoris stysi* Wagner, 1968.

intermedius Reuter, 1884 (*Capsus*). LT (designated by Vinokurov, 1977: 106): ♂ (damaged, genit. diss.), pn, Sibir. occ. [hR]/ Spec. typ. Reuter/ cR/ No. 3569. PLT: 1 ♀ (damaged), as LT but No. 3568; 1 ♂, 1 ♀, pn, Leusch/ Sundman/ Spec. typ.; 1 ♀, pn, Leusch/ Sundman/ 301; 1 ♂, Satiga/ Sundman/ 90; 1 ♂, Fl. Konda/ Sundman/ 242. Synonym of *Capsus wagneri* (Remane, 1950). – Note. In the original description, 3 localities were indicated (Katyshinskaya, Leush and Sutiga), all of them are in environs of Leushi, Khanty-Mansi National District, Tyumen' Prov.

**intermedius* Reuter, 1885 (*Dichrooscytus*). LT: ♀, gc, Leipzig [hR]/ Spec. typ. Reuter/ cR/ No. 3652.

**kiritschenkoi* Poppius, 1912 (*Phytocoris*). LT: ♀, gc, Peski Muyun-Kum, oz. Kargaly-Kul', 15.V.10, Kiritshenko [Cc]/ cR/ *Phytocoris kiritshenkoi* Poppius. Typ. [hP]/ No. 9556.

kristensenii Poppius, 1914 (*Engyptatus*). HT in Helsinki Museum. Currently *Nesidiocoris* id.

**lateralis* Reuter, 1896 (*Deraeocoris cordiger* var.). LT: ♀, gc, Paris [hR]/ cR/ v. *lateralis* Reut. [hR].

**lateralis* J. Sahlberg, 1878 (*Miris virens* var.). LT: ♀, pn, Jenisejsk [hS]/ J. Sahlbg. 81 [h]/ *M. virens* L. v. *lateralis* J. Sahlbg. [h; bottom label]. Junior synonym of *Stenodema sibirica* Bergroth, 1914. – Note. Lindberg (1951) erroneously indicated as types specimens from Abak[an] sav[od] collected in 1885 by Hammarström.

lenzii Horváth, 1889 (*Chilocrates*). HT in Helsinki Museum. Synonym of *Deraeocoris patulus* (Walker, 1873).

**lineaticollis* Reuter, 1891 (*Lasiomiris*). LT: ♂, gt, Java [hR]/ Spec. typ./ *Lasiomiris lineaticollis* Reut. [hR]. PLT: 2 ♀, gt (on one pin), *Lasiomiris lineaticollis* Reut. Java [hR]. Synonym of *L. albopisulos* (Lethierry, 1888).

**lividipennis* Reuter, 1903 (*Dimorphocoris*). LT: ♂, gc, Bejar, Spain, G.C.C./ Pxt VN-m-56 [= Pinxit V. Nybom, 1956] [h]/ cR. PLT: 1 ♂, gc, Bejar, Spain, G.C.C./ cR.

**loewi* Reuter, 1896 (*Lygus rubricatus* var.). LT: ♀, gc, Austria [hR]/ Spec. typ. Reuter/ cR/ No. 3628/ v. *Loewi* Reut. [hR].

longicornis Jakovlev, 1902 (*Solenoxyphus*). HT (♂) in Helsinki Museum. Currently *Psallopoides* id.

**longicornis* Reuter, 1901 (*Miris ferrugatus* var.). LT: ♂ (antennae missing), gc, Dschilaryk/ J. Sahlb./ 392 [h]/ *Miris ferrugatus* v. *longicornis* Reut. Typ. [hR]/ No. 11234. Currently *Leptopterna longicornis*.

**longicornis* Reuter, 1903 (*Phytocoris*). LT: ♀, gc, Ahnger/ 847/ Sulukly, 12.VI [h]/ No. 11514/ *Phytocoris longicornis* Reut. n. sp. Typ. [hR]. Junior synonym of *Ph. moestus* Reuter, 1903.

**longicornis* Reuter, 1900 (*Pronototropis*). LT: ♀, gc, Syria [hR]/ Spec. typ. Reuter/ No. 3348/ pinxit [hSeid] Seidenstücker. Currently *Pleuroxonotus* id.

**luridus* Reuter, 1875 (*Orthothylus*). LT: ♀, gc, Sakkola/ J. Sahlbg./ 403/ Spec. typ./ *luridus* Reut. [hS]/ No. 3081. PLT: remnants (3 legs), gc, Pyhäjä/ J. Sahlbg./ 350/ Spec. typ./ *luridus* Reut. [hS]/ No. 3080. Synonym of *O. fuscescens* (Kirschbaum, 1856).

**maculatus* Reuter, 1896 (*Lygus lucorum* var.). LT: ♀, gc, Pargas/ Spec. typ. Reuter/ cR/ No. 3642/ v. *maculatus* Reut. [hR]. PLT: 1 ♀, gc, Dw. [h]/ Dvoretz/ No. 3058 (as var. *nigronasutus* Reut.); 1 ♀, gc, Solomina/ Car. ross./ J. Sahlbg./ 456 [h]. Currently: *Lygocoris* id.

**maculicollis* J. Sahlberg, 1920 (*Megalocoleus molliculus* var.). LT: ♂, gc, Bjerno/ J. Sahlbg./ 53/ *Megalocoleus molliculus* Fall. var.? *maculicollis* m. [hS].

maroccanus Wagner, 1951 (*Dicyphus*). HT collected at Amismiz.

**melanocerus* Reuter, 1901 (*Dicyphus*). LT: ♀, gc, Verni/ J. Sahlbg./ 545 [h]/ cR/ *Dicyphus melanocerus* Reut. Typ. [hR]/ No. 11246.

minor Reuter, 1896 (*Lygus rubicundus* ssp.). HT: ♀, gc, Felgueira, Puton, 6, 82 [h]/ cR. Currently *Agonocoris* id. – Note. In the original description, the sex was indicated as ♂ (apex of abdomen difficult to see in the carded HT) and it was suspected that the taxon represents a separate species. Comparison with *A. eduardi* Ribes, 1977 described from Spain seems desirable.

**moestus* Reuter, 1903 (*Phytocoris*). LT: ♂ (destroyed, only some legs remained), gc, Kopet Dagh/ Ahnger/ 846/ *Phytocoris moestus* Reut. n. sp. Typ. [hR]/ No. 11515.

**montandoni* Reuter, 1895 (*Allodapus*). LT: ♂ brachypt., Moldavie, vall. du Berlad, A.L. Montandon/ Spec. typ. Reuter/ cR/ No. 3560.

**montandoni* Reuter, 1894 (*Psallus quercus* var.). LT: ♂ [sic], gc, Valachie, Comana, A.L. Montandon/ cR/ No. 9532/ var. *Montandoni* Reut. [hR].

morio J. Sahlberg, 1883 (*Atractotomus*). HT: ♂ (genit. diss.), gc (reglued, apparently by Stonedahl), Jaakkima/ J. Sahlbg/ ♂/ J. Sahlbg. ind. typ./ 49/ No. 3702.

**myrmecoides* Reuter, 1901 (*Aspidacanthus*). LT: ♀, gc, Amu-Daria/ J. Sahlbg./ 161 [h]/ cR/ No. 3559/ *Aspidacanthus myrmecoides* Reut. Typ. [hR].

**nigriceps* Reuter, 1894 (*Macrotylus paykulli* var.). LT: ♀, gc, Hispania [h]/ Spec. typ. Reuter/ cR/ No. 3313/ var. *nigriceps* Reut. [h]. – Note. The variety was described from Alicante.

**nigricornis* Reuter, 1878 (*Mauroductylus*). HT: ♀, gt, Ps. *nigricornis* n. sp. Astr. unicus [hJ]/ Spec. typ. Reuter/ cR/ No. 3175. Synonym of *M. albidus* (Koleatti, 1845). – Note. According to the indication "uni-

cus" in the label, the specimen must be considered HT.

**nigricornis* Reuter, 1904 (*Pachytomella phoenicea* var.). LT: ♀ brachypt., gc/ Sarona/ J. Sahlbg./ 210/ 210/ cR/ *Pachytomella phoenicea* var. *nigricornis* Reut. [hR] O.M. Reuter det./ No. 9541.

**nigrifrons* Reuter, 1906 (*Dicyphus*). LT: ♀, gc, Szechuan', dolina Tachzhinkho, Potan. [Cc], 22.VII.93/ cR/ No. 3509/ *D. nigrifrons* Reut. Typ. [hR]. – Note. The second syntype, in the Zoological Institute, St.Petersburg, is badly damaged.

**nigripes* Reuter, 1901 (*Labops*). LT: ♀ macropt., gc, N. Mongolei, Leder, 92/ Spec. typ. Reuter/ cR/ No. 3525. Synonym of *L. burmeisteri* Stål, 1858. – Note. In the original description, the type locality was given as "Irkut", but all examined syntypes (most of which are in the Naturhistorisches Museum Wien) are labelled "N. Mongolei". Apparently, Reuter indicated the type locality in error.

**nitens* Reuter, 1894 (*Orthocephalus*). LT: ♂ brachypt., gc, Alicante [h]/ *nitens* Reut. [hR]/ Spec. typ. Reuter/ cR/ No. 3494. Synonym of *Pachytomella doriae* (Reuter, 1884).

**noualhieri* Reuter, 1902 (*Atomoscelis*). LT: ♂ (genit. diss.), gc, Museum Paris, Algérie, Biskra, coll. Noualhier, 1898/ Spec. typ. Reuter/ No. 3232. PLT: 3 ♀, as LT, but No. 3229, 3230, 3231 and with additional label "cR".

noualhieri Reuter, 1896 (*Grypocoris*). The LT designated by Wagner (1968: 52, as HT) is labelled "Madrid". Currently *Hadrodemus* id.

**oberthuri* Reuter, 1905 (*Strongylotocoris obscurus* var.). LT: ♀, mpn, Museum Paris, France, Vernettes-Bains, Pyrénées orientales, R. Oberthur, 1897/ Spec. typ. Reuter/ cR/ No. 3488. Currently *S. oberthuri*.

oculata Wagner, 1954 (*Tuponia*). HT from Puerto de la Cruz.

**olivaceus* Reuter, 1880 (*Plagiognathus*). LT: ♀, gc, S. Nevada [hR]/ Spec. typ. Reuter/ cR/ No. 3251.

**oschanini* Reuter, 1883 (*Orthotylus*). LT: ♂ (genit. diss.), pn, Turkestan [hR]/ Spec. typ. Reuter/ cR/ No. 3420.

**osiris* Kirkaldy, 1902 (*Lygus*). LT: ♀, gc, I. Reunion, S. Denis, Ms. Hacoud [h]/ *Lygus osiris* Kirk. [hK]. Synonym of *Taylorilygus apicalis* (Fieber, 1861). – Note. At the card with LT was also a male, of which only some legs remained.

**pallidipennis* J. Sahlberg, 1868 (*Plagiognathus*). LT: ♀, gc, golden square/ Valamo/ J. Sahlbg./ 244/ No. 3099. Synonym of *Plesiodesma pinetella* (Zetterstedt, 1828).

**pallidus* Reuter, 1900 (*Plagiognathus*). LT: ♀, gc, Dr. J. Vosseler, Oued Saida, Oran, 6.1894/ Spec. typ. Reuter/ cR/ No. 3248.

**palmeni* Reuter, 1875 (*Calocoris*). LT: ♀ (abdomen and one wing missing), Alp. Noric./ Nassfeld/ Palmén/ Spec. typ. Reuter/ cR/ No. 3668. Synonym of *Calocoris lineolatus* (A. Costa, 1862).

**paludum* J. Sahlberg, 1870 (*Teratocoris*). LT: ♂, gc, golden square/ Dvoretz/ Car. ross./ J. Sahlbg./ Spec. typ./ 666 (or 999) [h]/ No. 3071.

parallelus Lindberg, 1930 (*Omocoris*). The HT is a male from Tedzhen. Synonym of *O. cunealis* (Reuter, 1904).

**parvulus* Reuter, 1879 (*Orthotylus*). LT: ♂ (genit. diss.), gc, Ross. mer. [hR]/ Spec. typ. Reuter/ No. 3442. – Note. In the original description, the type locality was indicated as Astrakhan.

**pedalis* Reuter, 1904 (*Pachytomella phoenicea* var.). LT: ♂ macropt., gc, Beirut/ J. Sahlbg./ cR/ *Pachytomella phoenicea* var. *pedalis* Reut. [hR] O.M. Reuter det.

**pictipes* Reuter, 1878 (*Excentricus punctipes* var.). LT: ♀, gc, Sibiria [h]/ Spec. typ. Reuter/ cR/ No. 3265. Currently *Excentricoris pictipes*. – Note. According to the original description, the type locality is Irkutsk. The second specimen cited by Lindberg (1951) is collected in N Mongolia and does not belong to the type series.

**picturatus* Reuter, 1896 (*Poecilonotus*). LT: ♂, Smyrna/ Spec. typ. Reuter/ cR/ No. 3667.

**pilifer* Reuter, 1880 (*Phytocoris*). LT: ♂, pn, Asia min. 22.5.72/ Spec. typ. Reuter/ No. 3121.

**pilipes* Reuter, 1879 (*Camptobrochis*). LT: ♀, pn, Peyshambe [Cc]/ Spec. typ. Reuter/ cR/ No. 3575. PLT: 1 ♀, pn, Turkestan [hR]/ Spec. typ. Reuter/ cR/ No. 3574. Currently *Deraeocoris* id.

**plagiathus* Reuter, 1876 (*Plagiognathus*). LT: ♂, gc, Lesio [hR]/ Spec. typ. Reuter/ cR/ No. 3249. PLT: 1 ♀, as LT but. 3250. – Note. Contrary to Wagner (1975), the species was described from North Italy, not from Caucasus.

**plagigera* Reuter, 1908 (*Phytocoris niveatus* var.). LT: ♂, ge, Elisbp. [hJ]/ 118 [h, on blue paper]/ cR.

**poppiusi* Reuter, 1911 (*Notostira*). LT: ♂, gc, Aleksandrov. khrebet, Syr-Dar'inskaya obl. [p, Cc], 9870', per. Makbal, 3.VI.10 [hKir, Cc]. A. Kiritshenko [p, Cc]. PLT: 1 ♂ (head and pronotum missing), gc, and 3 ♀, pn, same label, one female with additional label "*N. poppiusi* Kir. ♀" [h].

**prasinus* Reuter, 1908 (*Calocoris*). LT: ♀, pn, Khappusi, 17.VI.01, P. Schmidt [h, Cc]/ cR/ No. 3676/ *Calocoris prasinus* n. sp. [hR] O.M. Reuter det. PLT: 1 ♀, as LT but No. 3677 and without the last label; 1 ♀, gt, Mauka, 13.VI.01, P. Schmidt [cC]/ cR/ No. 3678. Synonym of *Mermiteilocerus annulipes annulipes* Reuter, 1908.

**propinquus* Reuter, 1883 (*Orthotylus*). LT: ♂, gc, Hispania [hR]/ *propinquus* [sic] Reut. [hR]/ Spec. typ. Reuter/ No. 3410. Synonym of *O. ochrotrichus* Fieber, 1864.

**psammaeicolor* Reuter, 1885 (*Trigonotylus ruficornis* var.). LT: ♂, gc, Scotia/ Spec. typ. Reuter/ cR/ No. 3782. PLT: 2 ♀, as LT but No. 3783, 3784. Currently *T. psammaeicolor*.

**pseudosabinae* Reuter, 1896 (*Dichrooscytus*). LT: ♀, pn, Turkestan [hR]/ Spec. typ. Reuter/ cR/ No. 3653.

**pullus* Reuter, 1870 (*Agaliastes*). LT: ♀, gc, Par-gas/ Reuter/ Spec. typ./ No. 3103. Currently *Chlamydatus* id.

**punctiger* Reuter, 1894 (*Oncotylus*). LT: ♀ (head and pronotum missing), Escorial, 29 Junio [h] Boli-var/ Spec. typ. Reuter/ cR/ No. 3368.

**punicus* Ferrari, 1884 (*Calocoris sexpunctatus* var.). LT: ♂, pn, Tunis, dint. IV. 1882, G. e. L. Dorial v. *punicus* Ferr. [hR]/ Spec. typ. Ferrari/ cR/ No. 3686/ v. *thoracica* Put. [hR]. Synonym of *Caloco-ris nemoralis* (Fabricius, 1787).

- ***putonii** Reuter, 1875 (*Oncotylus*). LT: ♀, gc, La Nouvelle/ cR. Currently *Pastocoris* id. – Note. Lindberg (1951) cited in error two specimens from “Attica” not belonging to the type series.
- quadrimaculatus** Reuter, 1903 (*Rhopaliceschatus*). STS: 2 ♀, pn, Museum Paris, Mou-Pin, A. David, 1870/ Thibet (Mou-Pin), A. David, 1869-70/ *Rhopaliceschatus* *4maculatus* Reut. [hR?; under one of the specimens]
- ***quadripunctata** Reuter, 1902 (*Calocoris hispanicus* var.). LT: ♀, gc, Tunis/ J. Sahlb./ *Calocoris hispanicus* v. *4maculata* [sic] Reut. Typ. [hR]/ No. 11237. PLT: 1 ♂, gc, Tunis/ J. Sahlb./ Spec. typ. Reuter/ No. 3681/ cR/ v. *4maculata* [sic] [hR]. Synonym of *Calocoris nemoralis* (Fabricius, 1787).
- ***quercicola** Reuter, 1904 (*Sthenarus*). LT: ♀, gc, Fl. Hermos/ J. Sahlb./ 718/ Spec. typ. Reuter/ cR/ No. 3193/ *Psallus quercicola* n. sp. [hR] O.M. Reuter det. Currently *Psallus* id.
- ***ravana** Kirkaldy, 1902 (*Shana*). LT: ♂, gc, Birmania, Chan Yoma, Gribodo [h]/ *Shana ravana* Kirk. ♀ type [hK]. Synonym of *Deraeocoris patulus* (Walker, 1873).
- ***reuteri** Reuter, 1879 (*Oncotylus*). LT: ♂, pn, Turkestan [hR]/ Spec. typ. Reuter/ cR/ No. 3372. – Note. The species was described from Tashkent.
- ***rubicundus** Reuter, 1904 (*Calocoris*). LT: ♀, gc, Bulgar Dagh/ U. Sahlb./ 644/ spec. typ./ cR/ No. 3687/ *Calocoris rubicundus* n. sp. [hR] O.M. Reuter det.
- ***rubidus** Reuter, 1896 (*Poeciloscytus*). LT: ♀, pn, Sibiria [hR]/ Spec. typ. Reuter/ cR/ No. 3589. Currently *Polymerus* id. – Note. In an earlier note (Reuter, 1879: 55, as *vulneratus* var.), the type locality was stated as Irkutsk.
- rubrinervis** Reuter, 1880 (*Ischnoscelis*). HT: ♂, pn, 165 [h]/ *Ischnoscelis rubrinervis* Reut. n. g. et sp. [hR]/ Type [red label]. Currently *Ischnoscelicoris* id.
- ***ruficeps** Reuter, 1880 (*Megacoelum*). LT: ♂, pn, Pelopones/ Spec. typ. Reuter/ cR/ No. 3705/ var. *ruficeps* Reut. [hR]. Synonym of *Megacoelum heckeri* (Fieber, 1870).
- ***rufipes** Reuter, 1902 (*Calocoris hispanicus* var.). LT: ♂, gc, Carthago/ J. Sahlb./ 996 [h]/ *Calocoris hispanicus* v. *rufipes* Reut. Typ. [hR]/ No. 11838. Synonym of *Calocoris nemoralis* (Fabricius, 1787).
- ***sahlbergi** Reuter, 1901 (*Psallus criocorooides* var.). LT: ♂, gc, Tunis/ J. Sahlb./ *criocorooides* v. *sahlbergi* Reut. [hR]/ Spec. typ. Reuter/ cR/ No. 3272. PLT?: 1 ♂, gc, Tebessa/ J. Sahlb./ 926 [h]/ cR/ *Psallus criocorooides* v. *sahlbergi* Reut. Typ. [hR]/ No. 11348. Synonym of *P. siculus* Reuter, 1875. – Note. The locality Tebessa was not mentioned in the original description.
- ***salicicola** Reuter, 1880 (*Globiceps*). LT: ♂, gc, Pargas/ cR/ Spec. typ. Reuter/ No. 3074. PLT: 1 ♀, as LT but No. 3075.
- ***samojedorum** J. Sahlberg, 1878 (*Calocoris*). LT: ♂, gc/ Dudinka [hS]/ J. Sahlb./ J. Sahlb. ind. typ./ 239 [h]/ *Calocoris samojedorum* J. Sahlb. [h; bottom label]. PLT: 1 ♀, pn, Dudinka [hR]/ Spec. typ. J. Sahlberg/ cR/ No. 3675.
- sanguinolentus** Reuter, 1879 (*Lygus* (*Orthops*)). The LT designated by Wagner (1955: 151) is collected at Isfaram. Currently *Orthops* id.
- ***saundersi** Reuter, 1896 (*Lygus*). LT: ♀, gc. Amur [hR]/ Spec. typ. Reuter/ No. 3630.
- ***saundersi** Reuter, 1896 (*Phytocoris*). LT: ♂, mpn, Algérie, Edough [h]/ Spec. typ. Reuter/ cR/ No. 3710.
- ***schoberiae** Reuter, 1876 (*Orthotylus*). LT: ♂, gc, Hungaria [hR]/ Spec. typ. Reuter/ cR/ No. 3434. PLT: 1 ♀, as LT but No. 3435.
- ***scirtetoides** Reuter, 1891 (*Mecistoscelis*). LT: ♂, gt, *Mecistoscelis scirtetoides* Reut. Java [hR]/ *Mecistoscelis scirtetoides* n. sp. [hR] O.M. Reuter det.
- ***scitulus** Reuter, 1908 (*Phytocoris*). LT: ♀, gc. Adzik. [h]/ cR/ No. 3742/ *scitulus* Reut. Typ. [hR]. PLT: 2 ♀, as LT, but No. 3740 and 3741, and one with additional label “Spec. typ. Reuter”.
- ***scotti** Reuter, 1877 (*Orthotylus*). LT: ♀, gc, Wimb., 14.8.76 [h]/ Spec. typ. Reuter/ cR/ No. 3407. Synonym of *O. prasinus* (Fallén, 1826). – Note. The second specimen considered by Lindberg (1951) as type is dated 1877 and apparently does not belong to the type series (the original description was published in August 1877 and Reuter indicated that he examined only one female received from Saunders; it is not an evidence for HT, because Reuter assigned to this species also specimens misidentified in the works of British authors).
- ***scutellaris** Reuter, 1891 (*Camptothrochis*). LT: ♀, gt, Java [hR]/ *Camptothrochis scutellaris* n. sp. [hR] O.M. Reuter det./ No. 9965. Junior synonym of *Deraeocoris scutellaris* Carvalho, 1957.
- ***setosus** Reuter, 1891 (*Lahops*). LT: ♂ macropt., gc, V. Sujetuk/ Hammarstr./ Spec. typ. Reuter/ cR/ No. 3254. PLT: 3 ♂ macropt., 1 ♀ macropt., 3 ♀ brachypt., same locality and collector.
- ***sibirica** J. Sahlberg, 1878 (*Pachytoma*). LT: ♂ brachypt., gc, Krasnoj./ Spec. typ. J. Sahlberg/ cR/ No. 3523. PLT: 1 ♂ brachypt., 2 ♀ macropt. from Krasnoyarsk. Synonym of *Anapus rugicollis* (Jakovlev, 1877).
- ***signaticornis** Reuter, 1904 (*Atomoscelis*). LT: ♀, gc, Heliopolis/ J. Sahlb./ Spec. typ. Reuter/ cR/ No. 3228. Currently *Badezorus* id.
- signatus** Reuter, 1878 (*Actinocoris*). HT: ♂, gc, Pargas/ Reuter/ Spec. typ./ No. 3073.
- ***signatus** J. Sahlberg, 1868 (*Agalliaestes*). LT: ♂, gc, golden square/ Enontekis/ J. Sahlbg./ ♂/ Spec. typ./ No. 3108. PLT: 8 specimens from Enontekis. Synonym of *Chlamydatus opacus* (Zetterstedt, 1838). – Note. The species was described from several localities in the province Enontekiö Lappi (the original description uses Törneå Lappmark, the earlier Swedish name).
- ***signifer** Reuter, 1909 (*Lygus pabulinus* var.). LT: ♂, gt, Plummer's I., Md. [p], Aug. 16.06/ Heidemann collector. PLT: 1 ♀, as LT. Currently *Lygocoris* id.
- simulans** Horváth, 1877 (*Piezocranum*). HT: ♂, gc, Buda, 2.6.74/ Spec. typ. Horváth/ cR/ No. 3489.
- sinicus** Poppius, 1915 (*Phytocoris*). The HT is kept in Helsinki Museum.
- ***steganoides** J. Sahlberg, 1875 (*Stiphrosoma*). LT: ♂, pn, Kantalaiks/ J. Sahlb./ J. Sahlb. ind. typ./ No. 3082. PLT: 1 ♀, as LT, but with additional label “128” and No. 3083. Currently *Strongylocoris* id. – Note. Specimens No. 3084, 3085, 3485 were labelled as types in error.

**strigicollis* Lindberg, 1933 (*Calocoris*). LT: ♂, pn, Portugal, Pomarasam, Guadiana, IV.1921 [h]/ M. Brakenhielm ded. 30.VI.1921/ Spec. typ. [p] *Calocoris strigicollis* [hL] Lindberg/ coll. Lindberg. Synonym of *Hadrodemus nouathieri* (Reuter, 1896).

**syracus* Reuter, 1896 (*Grypocoris*). LT: ♂, gc, Aintab [h]/ Spec. typ. Reuter/ cR/ No. 3664.

**tarsalis* Reuter, 1894 (*Amblytylus*). LT: ♂, gc, Bagnuls [hR]/ Spec. typ. Reuter/ No. 3317. PLT: 2 ♀, gc, Bagnuls [hR]/ Spec. typ. Reuter/ cR/ No. 3318, 3320.

**tenebrosus* Reuter, 1875 (*Calocoris*). LT: ♀, pn, Sibiria [hR]/ Spec. typ. Reuter/ cR/ No. 3696. Currently *Adelphocoris* id. – Note. The type locality was indicated (Reuter, 1879) as environs of Irkutsk.

**testaceus* Reuter, 1879 (*Dicyphus*). LT: ♀, gc, Shagimardan [Cc]/ Spec. typ. Reuter/ cR/ No. 3461. – Note. The second syntype, in Zoological Museum of the Moscow University, is badly damaged.

**tibialis* Reuter, 1892 (*Dolichomiris*). LT: ♀, gc, Colonia Tovar, E. Simon, I.II.88/ No. 9645. Synonym of *D. linearis* Reuter, 1882.

**tibialis* Reuter, 1891 (*Halticus*). LT: ♂, gt, Java [hR]/ Spec. typ./ *Halticus tibialis* Reut. Java [hR].

**tomentosus* Reuter, 1904 (*Atomoscelis*). LT: ♀, gc, Heliopolis/ J. Sahlb./ Spec. typ. Reuter/ cR/ No. 3234. Currently *Badezorus* id.

tridens Wagner, 1954 (*Phytocoris*). The HT is from Ayios Hilarion.

**trispinosum* Reuter, 1904 (*Stenodema*). LT: ♀, gc, Batylin/ Fl. Lena m./ B. Poppius/ No. 3037/ *trispinosum* Typ. [hR] O.M. Reuter det. PLT: no less than 10 specimens from various localities.

**unicolor* Reuter, 1880 (*Phytocoris*). LT: ♀, pn, Peloponnes/ *unicolor* Reut. [hR]/ Spec. typ. Reuter/ cR/ No. 3788. Synonym of *Ph. exoletus* A. Costa, 1853.

verbena Wagner, 1954 (*Coniortodes*). Contrary to Ribes & Goula (1986: 284), the HT is in Helsinki Museum. Currently *Compsidolon* id.

**vitelinus* Reuter, 1903 (*Megalocoleus albidus* var.). LT: ♀, gc, Aschabad, Ahnger/ 621/ Spec. typ. Reuter/ cR/ No. 3336. Synonym of *Lopidodenus albidus* (Reuter, 1903). – Note. One more syntype is lost, only a pin with labels remained.

**vitticeps* Reuter, 1902 (*Halticus luteicollis* var.). LT: ♀, pn, Museum Paris, env. de Tanger, Favier, 1857/ Spec. typ. Reuter/ cR/ No. 3487/ v. *vitticeps* Reut. [hR].

**vittiger* Reuter, 1896 (*Phytocoris*). LT: ♂, pn, Espinar, Agosto de 1894, Bolivar/ Spec. typ. Reuter/ cR/ No. 3758. – Note. One more syntype (No. 3759) completely destroyed.

References

- Josifov, M. 1974. Beitrag zur Systematik der paläarktischen *Dichrooscytus*-Arten (Heteroptera, Miridae). *Reichenbachia*, **15**: 149-173.
- Josifov, M. & Kerzhner, I.M. 1972. Heteroptera aus Korea. I. Teil (Ochteridae, Gerridae, Saldidae, Nabidae, Anthocoridae, Miridae, Tingidae und Reduviidae). *Ann. zool. (Warszawa)*, **29**: 147-180.
- Lindberg, H. 1951. Verzeichnis der Typen in O.M. Reuters paläarktischer Heteropterensammlung. *Comment. biol.*, **12**(14): 1-34.
- Linnauvori, R. 1964. Hemiptera of Egypt, with remarks on some species of the adjacent Eremian region. *Ann. zool. fenn.*, **1**: 306-356.
- Reuter, O.M. 1879. De Hemipteris e Sibiria orientali nonnullis adnotaciones criticae. *Öfv. finnska Vet. Soc. Förh.*, **21**: 42-63.
- Reuter, O.M. 1894. Ad cognitionem Capsidarum. II. Capsidae palaearcticae. *Rev. Entomol.*, **13**: 128-152.
- Reuter, O.M. 1896. *Hemiptera Gymnocerata Europae. Hémiptères Gymnocérates d'Europe, du bassin de la Méditerranée et de l'Asie Russe*, **5**: 1-392. Helsingfors [also published in *Acta Soc. Sci. fenn.*, **23**(2): 1-392].
- Ribes, J. & Goula, M. 1986. Dr. E. Wagner's entomological collection: Miridae (Insecta, Heteroptera) preserved in the Zoological Museum Hamburg (FRG). *Mitt. Humb. zool. Mus. Inst.*, **8**: 243-335.
- Vinokurov, N.N. 1977. On the systematic and intra-specific variability of capsid bugs of the genus *Capsus* F. (Heteroptera, Miridae). *Entomol. Obozr.*, **56**: 103-115. (In Russian).
- Wagner, E. 1955. Neuer Beitrag zur Systematik der Gattung *Lygus* Hhn. (Hem. Het. Miridae). *Acta entomol. Mus. natn. Pragae*, **29** (1954): 149-158.
- Wagner, E. 1967. Über *Phytocoris* s. str. (Heteroptera, Miridae, *Phytocoris* Fall.). *Reichenbachia*, **8**: 271-284.
- Wagner, E. 1968. Was ist *Grypocoris noualhieri* Reuter, 1896? (Heteroptera, Miridae). *Notulae entomol.*, **48**: 49-54.
- Wagner, E. 1970. Über *Plagiorrhamma* Fieber, 1870 (Heteroptera, Miridae). *Notulae entomol.*, **50**: 67-72.
- Wagner, E. 1975. Die Miridae Hahn, 1831, des Mittelmeerraumes und der Makaronesischen Inseln (Hemiptera: Heteroptera), Teil 3. *Entomol. Abhandl. staatl. Mus. Tierk. Dresden*, **40**, Suppl.: i-ii, 1-483.

Acknowledgements

The author is thankful to Dr L. Huldén and Dr A. Jansson (Helsinki) for help in preparation of this work.

Received 12 August 1997